

DIRECTIONS FOR USE

IMPORTANT

GROUND APPLICATION Vegetable and Field Crops Citrus Cotton (On Young Fruit)  
Deciduous Fruit Crops  
Plantings  
Where Recommended for AIR APPLICATION Field Crops

MIXING DIRECTIONS

ALFALFA Grasshoppers Alfalfa Webworms Armyworms Aphids Looppers  
Leafhoppers Lygus Bugs Alfalfa Mites ALFALFA (Speed) Lygus  
Mite Suppression Mites Thrips  
By Air

BEANS (Dry or Field) SOYBEANS LIMA BEANS AND PEAS Alfalfa Looppers Aphids  
Spider Mites Lygus Leaf Miners GREEN BEANS Aphids Spider Mites Looppers Nitidulids Leaf  
hoppers Large Looppers Stink  
Bugs

BROCCOLI CABBAGE CAULIFLOWER BRUSSELS SPROUTS KALE LOLLARDS Imported Cabbage  
worms Diamond Back Caterpillars Aphids Looppers

CELERY Aphids Looppers Armyworms Salt Marsh Caterpillars Leaf Miners

COTTON Lygus Leaf Perforators Fleahoppers Armyworms  
Bollworms Looppers and for Suppression of Mites  
FOR HEAVY POPULATION OF LISTED INSECTS

CUCUMBERS SUMMER SQUASH Cucumber Beetles Aphids Spider Mites  
worms Pickleworms Leaf Miners  
By Air

EGGPLANT PEPPERS Aphids Spider Mites blister Beetles Flea Beetles

SPINACH CHARD TURNIP GREENS Aphids Looppers Armyworms Spider  
Miners

MELONS (Cantaloupe Honeydew Muskmelon Watermelon) PUMPKINS  
Armyworms Leafhoppers Spider Mites Leaf Miners

ORANGES LEMONS GRAPEFRUIT TANGERINES Aphids Soft Brown  
Culicids Leaf Rollers (Tortrix) (Calf and Aviz) Thrips

RICE Grasshoppers

TOBACCO Hornworms Flea Beetles

TOMATOES (Field) Fruit Flies (Drosophila)

Worms Hornworms Leaf Miners Flea Beetles Mites (Field and  
For Field)

GREENHOUSE VAPOR TREATMENT OF ROSES OTHER ORNAMENTALS  
CUCUMBERS Spider Mites Adult Whiteflies Aphids Leafrollers, Beetles

9859-258  
NOV 7 1975  
VICIUS  
LIFT ACT  
CUTTER-  
SUBJECTS



**FASCO**

# ROM 8 EC

INSECTICIDE

NET WT. 1.05 LBS (475 G)  
BY WEIGHT

**CAUSES EYE AND SKIN DAMAGE  
MAY BE FATAL IF SWALLOWED OR  
ABSORBED THROUGH SKIN  
OUT OF REACH OF CHILDREN**

Keep out of reach of children. Do not use in areas where children play. Do not use in areas where food is stored or prepared. Do not use in areas where food is eaten. Do not use in areas where food is stored or prepared. Do not use in areas where food is eaten. Do not use in areas where food is stored or prepared. Do not use in areas where food is eaten.

1 GALLON

Spider Mites  
Aphids  
Adult Whiteflies

**NOTE**

FRUIT FLIES: Do not use on and around food processing plants, livestock, poultry, and pet areas. Under Mites

**ENVIRONMENTAL PRECAUTIONS**

DO NOT USE IN WATERS, STREAMS, PONDS, AND WETLANDS. DO NOT USE IN AREAS WHERE BIRDS, BEES, AND OTHER POLLINATING INSECTS ARE ACTIVE. DO NOT USE IN AREAS WHERE FISH, SHEEP, GOATS, MARSHES, SWAMPS, WETLANDS, AND OTHER WILDLIFE ARE FOUND. DO NOT USE IN AREAS WHERE CHILDREN PLAY.

**KEEP OUT OF REACH OF CHILDREN**

NET WT. 1.05 LBS (475 G)

**CLOVER MITES, ROACHES, LARVAE, AND STAGBEEFLIES** (not on lawns)

**GRASSHOPPERS** (not on lawns and pastures)

**FLYING**

**NOTE TO PHYSICIAN**

**ENVIRONMENTAL PRECAUTIONS**

DO NOT USE IN WATERS, STREAMS, PONDS, AND WETLANDS. DO NOT USE IN AREAS WHERE BIRDS, BEES, AND OTHER POLLINATING INSECTS ARE ACTIVE. DO NOT USE IN AREAS WHERE FISH, SHEEP, GOATS, MARSHES, SWAMPS, WETLANDS, AND OTHER WILDLIFE ARE FOUND. DO NOT USE IN AREAS WHERE CHILDREN PLAY.

**KEEP OUT OF REACH OF CHILDREN**

NET WT. 1.05 LBS (475 G)

MANUFACTURED BY  
**KEMW-MDCEE CHEMICAL CORPORATION**  
OKLAHOMA CITY, OKLAHOMA