Kang to 3:5 - 27/ EPA REGISTRATION NO. 9859-266 U.S. ENVIRONMENTAL PROTECTION AGENCY OFFICE OF PESTICIDES PROGRAMS REGISTRATION DIVISION (WH-567) TERM OF ISSUANCE WASHINGTON, D.C. 20460 NOTICE OF PESTICIDE: REGISTRATION NAME OF PESTICIDE PRODUCT REREGISTRATION Tanco Parathion 2% Bait (Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended) NAME AND ADDRESS OF REGISTRANT (Include ZIP code) Т Марола Сэкцион скоро<mark>алу</mark> Line Dogwood Acre Tab Patie 51, 32902 L NOTE: Changes in labeling formula differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Figistration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above U.S. EPA registration number. On the basis of information furnished by the registrant, the above named pesticide is hereby Registered/Reregistered under the Federal Insecticide, Fungicide, and Rodenticide Act. A copy of the labeling accepted in connection with this Registration/Reregistration is returned herewith. Registration is in no way to be construed as an indorsement or approval of this product by this Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others. to go duct a complete to be a construction of the second of the Company of the Company Other to some or relative provides contracted to the provides and the provides to the provides of Note that the Miller area. We also when the Admin make all Tropical transport to engineer and distant * 1 (11)* of, the treatment of the Reconstration D. Her Children Communication of construction of the form the firm of the contract of the contr 3. costs to the contract of your type to be contacted label the between your consist the supposed one hipmont. Extended the A 79 Enoles we have a turther second trep of the nall respect tabeling. official terror contracts in the property of my recent with, the respect terration is visited by notes to the established by the Actor proposed with Fables and a file). When pelicase for etimos is a tito i compilator and i andethese committees. I stamped conversion the large property of vone secretar. Jav S. I Hermelder Poduct Manac∈r (* connectiondo endenomide Franci Terri tration Division (98-767) TTACHMENT IS PPLICABLE

EPA Form 8570-4 (Rev. 5-76)

SIGNA TIRE DE AF HOS POPES

Remarkable print

(Center Panel)

RESTRICTED USE PESTICIDE

/For retail sale to and use only by Certified Applicators or persons under their indirect supervision and only for those uses covered by the Certified Applicator's Acertification.

PARATHION 2% BAIT

ACTIVE INGREDIENT

Parathion (0, 0 - diethyl - 0 - p-nitrophenyl

phosphorothicate) 2.0 %

4 4

KEEP OUT OF REACH OF CHILDREN

POISON

DANGER

PELIGRO

'PRECAUCION AL USUARIO: Si usted no lee insles, no use este producto
'hasto que la etiqueta haya sido explicada ampliamente.

NOT FOR HOME USE

FIRST AID TREATMENT

"STATEMENT OF PRACTICAL TREATMENT"

Call a doctor (physician), clinic or hospital immediately. Explain that the victim has been exposed to parathion and describe his condition. If the doctor cannot come, take victim to the clinic or hospital. Move victim immediately from the area where parathion is present. If INHALED, remove victim to fresh air. If not breathing, immediately give artificial respiration, preferably mouth-to-mouth, and maintain until doctor sees victim. If breathing is difficult, give oxygen. IF IN EYES OR ON SKIN, immediately flush with plenty of water for at least 15 minutes while removing contaminated personal clothing and shoes to avoid continued possible exposure to victim or persons in attendance. IF SWALLOWED, induce vomiting immediately by giving two glasses of water and by touching back of throat with finder. DO NOT INDUCE VOMITING OR GIVE AMAGE. THING BY MOUTH TO AN UNCONSCIOUS PERSON. Have victim lie down and sheep duret.

Mfs. By Landia Chemical Company Lakeland, Florida 33802

MICEPTED
with OMMENTS
EPA Letter Dated:

JUN 1 31985

Under the Ledend Exections. Fungicide, and Rodenticide Act an amended, for the pesticide registered under EPA Res. No.

9759-266

NET WEIGHT:

RE-ENTRY STATEMENT
Do not apply this product in such a manner
as to directly or through drift expose
workers or other persons. The area being
treated must be vacated by unprotected
persons. Do not enter treated areas for 48
hours unless appropriate protective clothins
is worn. Because certain states may require
more restrictive reentry intervals for various
crops treated with this product, consult your
State Department of Agriculture for further
information.

Written or oral warnings must be given to workers who are expected to be in a treated area or in an area about to be treated with this product.

ORAL WARNINGS shall inform workers of areas or fields that may not be entered, without protective equipment and clothing specified on this label, during treatment or for 48 hours following treatment. Follow First Aid Treatment instructions shown on this label in case of accidental exposure.

When oral warnings are given; warnings shall be given in a language customarily understood by workers. Oral warnings must be given if there is reason to believe that written warnings cannot be understood by workers.

WRITTEN WARNINGS must include the following information:

DANGER-Area treated with paration on (date).

Do not enter without appropriate protective clothing for 48 hours. Follow First Aid Treatment instructions shown on this product label in case of accidental exposure.

ATTENTION

To avoid excessive residues of parathion on food or forage crops always observe the statements found under "Directions for Use," limiting the time before harvest when parathion may be applied.

Because this material is poisonous by skin contact, inhalation or swallowing it should not be used in such a manner or under weather conditions as will permit drift of the spray onto areas not intended to be treated. Do not apply or allow drift to areas occupied by unprotected humans or beneficial animals.

EPA Re≤. No. 9859-EPA Est. No. 9859-FL-1

EPA Est. No. 51036-BA-1

(LEFT PANEL)

剛。

PRECAUTIONARY STATEMENTS Hazards to Humans and Domestic Animals

Keer out of reach of children.

DANGER! PELIGRO!

PRECAUCION AL USUARIO: Si usted no lee ingles, no use este producto hasta que la etiqueta hava sido explicada ampliamente.

POISONOUS IF SWALLOWED, INHALED OR ABSORBED THROUGH SKIN.

RAPIDLY ABSORBED THROUGH SKIN.

REPEATED EXPOSURE HAY, WITHOUT SYMPTOMS, BE INCREASINGLY HAZARDOUS.

Do not breathe vapors or dust. Do not set in eyes, on skin or on clothing.

POISON SIGNS(Symptoms)

Parathion is a very dangerous poison. It rapidly enters the body on contact with all skin surfaces and eyes. Clothing wet with this material must be removed immediately. Exposed persons must receive prompt medical treatment for they may die.

Some of the signs and symptoms of poisoning are: Headaches, nausea, vomiting, cramps, weakness, blurred vision, pin-point pupils, tightness in chest, labored breathing, nervousness, sweating, watering of eyes, drooling or frothing of mouth and nose, muscle spasms and coma.

POISON

ATROPINE IS AN ANTIDOTE CONSULT PHYSICIAN FOR EMERGENCY SUPPLY

NOTE TO PHYSICIAN

Antidote-administer atropine sulfate in large doses, TWO to FOUR ms.
intravenously or intramuscularly as soon as evanosis is overcome. Repeat
at 5 to 10 minute intervals until signs of atropinization appear. 2-PAM
choloride is a pharmacologic antidote and may be administered as an adjunct
to, but not a substitute for, atropine which is a symptomatic and often life
saving antidote. DO NOT GIVE MORPHINE OR TRANQUILIZERS BECAUSE THESE ANTIDOTES MAY PRODUCE ADVERSE EFFECT ACCENTUATING THE PHARMACOLOGIC EFFECT
OF THIS PRODUCT. This product is a strong cholinesterase inhibitor affecting the central and peripheral nervous systems and producing cardiac and
respiratory depression. At first sign of pulmonary edema, the putient should
be given supplemental oxygen and treated symptomatically. Continued absorption of the poison may occur and fatal relapses have been reported after '.'
initial improvements: VERY CLOSE SUPERVISION OF THE PATIENT IS INDICATED...
FOR AT LEAST 48 HOURS.

ENVIRONMENTAL HAZARDS

This product is toxic to fish, shrimp, crab, birds, and other wildlife, Birds and other wildlife in treated areas may be killed. Keep out of lakes, gireams, ponds, tidal marshes and estuaries. Shrimp and crate may be killed at application rates recommended on this label. Do not apply

where these are important rescurces. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment, or disposal of wastes. Apply this product only as specified on this label. This product is highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your Cooperative Agricultural Extension Service.

PHYSICAL OR CHEMICAL HAZARDS

In case of:

FIRE-Use water spray, foam, dry chemical or CO2. SPILL or LEAK, keep all unprotected persons away. Cover with absorbent such as soda ash, lime, clay or saw dust. Sweep up and bury. Wash area thoroughly with strong lye solution.

WORK BAFETY RULES

Persons working with this product should have frequent blood tests of their cholinesterase levels. If the cholinesterase level falls below a critical point, no further exposure should be allowed until it has been determined by means of blood tests that the cholinesterase level has returned to normal. Before using this product, consult the Agricultural Extension Service or Experimental Station for specific recommendations regarding such blood tests. If handled indoors, provide mechanical exhaust ventilation.

Keep all unprotected persons and animals away from treated area or where there is danger of drift.

Do not rub eyes or mouth with hands. If you feel sick in any way, STOP work and set help right away. See First Aid (Practical Treatment) section.

USE ON. WHEN WEARING THE FOLLOWING PROTECTIVE EQUIPHENT AND CLOTHING

(1) Wear water-proof pants, coat, hat, rubber boots or rubber overshoes.
(2) Wear safety soddles. (3) Wear a pesticide respirator Jointly approved by the Mining Enforcement and Safety Administration (formerly the U.S. Bureau of Mines) and the National Institute for Occupational Safety and Health under the provisions of 30 CFR part 11. (4) Wear heavy duty, natural rubber sloves.

IMPORTANT! Before removins sloves, wash them with soap and water. Always wash hands, face and arms with soap and water before smokins, extins or """ drinkins.

AFTER WORK, take off all clothing, and shoes. Shower, using soap and water. Wear only clean clothes when leaving Job. Do not wear contaminated clothing. Wash protective clothing and protective equipment with soap and water after asch use. Respirator should be cleaned and filter replaced according to instructions included with respirator.

(Right F nel)

STORAGE AND DISPOSAL

1 - 10 1980

.Do not contaminate water, food, or feed by storage or disposal.

- 1. Handle container carefully so as to avoid damage and prevent spills,
- 2. Do not use or store in or around the home.
- 13. Store container in a well ventilated place.
- 4. Empty container retains vapor and product residue.
 Observe all labeled safeguards until container is cleaned, reconditioned or destroyed.
- '5. Keep out of reach of children and domestic animals. DISPOSAL:

Wastes of this pesticide are acutely hazardous. Improper disposal of excess resticide, spray mixtures, or rinsate is a violation of federal law. If these wastes cannot be disposed of according to label use instructions, contact your state pesticide or Environmental Control Agency, or the Hazardous Waste Representative at the meanest EPA Regional Office for guidance. Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or incinerate, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

DIRECTIONS FOR USE

Restricted Classification - It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

FOR AGRICULTURAL USE ONLY

Apply at the recommended rates when insects first appear. Remeat application when needed. Do not apply closer to harvest than the days indicated for each crop. For aerial application, use the pounds per acre indicated for each crop. For preplant broadcast soil treatment, incorporate into top 4-9 inches of soil immediately after application. Use lower dosade for mineral soils and higher, dosade for ordanic soils.

CROPS

BEANS: To control cutworms and wireworms apply at a rate of 30 to 40 pounds per acre as a broadcast soil application. Do not apply within 15 days of harvest.

BLACKEYED_PEAS_AND_PEAS: To control cutworms and wireworms apply at a rate of no more than 25 pounds per acre as a broadcast soil application. Bo not'Maply within 15 days of harvest.

BEEIS: To control cutworms and wireworms apply at a rate of 30 pounds per acre as a broadcast soil application. Do not apply within 15 days of harvest to the roots. If tops are to be used for food or feed, do not apply within 21 days of harvest.

/|CANIALDURES=_MELONS: To control wireworms apply at a rate of 25 pounds per

acre as a broadcast application. Do not apply within 7 days of harvest. Do not apply before vining CARROIS: To control wireworms apply at a rate of not more than 30 younds per acre as a broadcast soil application. Do not apply within 15 days of harvest. Do not use treated tors for food or feed. CELERY: To control cutworms and wireworms apply at a rate of 30 to 40 pounds per acre as a broadcast soil application. Do not apply within 30 days of · harvest. CORN: To control cutworms and wireworms apply at a rate of 30 to 40 Pounds per acre as a broadcast application. Do not apply within 12 days of harvest. .CORNISMEEIl: To control wireworms use a preplant broadcast soil treatment. Incorporate into top 6 inches of soil using 30 to 40 pounds per acre. lower dosage for organic soils. RESTRICTED TO FLORIDA UNLY. CUCUMBER__LEIIUCE__RADISHES: To control cutworms and wireworms apply at a rate of 25 pounds per acre as a broadcast application. Do not apply within 15 days of harvest on cucumbers and radishes. Do not apply within 21 days of harvest on lettuce. On the cucumbers, do not apply before vining, EGGPLANI: GARLIC: To control cutworms and wireworms apply at a rate of 25 rounds per acre as a broadcast application. Do not apply within 15 days of harvest. CNDIVE: To control cutworms and wireworms apply at a rate of 25 pounds per acre as a broadcast application. Do not apply within 21 days of harvest. KALE: To control cutworms and wireworms apply at a rate of 25 pounds per acre as a broadcast application. Do not apply within 10 days of harvest. KOHLRABI: To control cutworms and wireworms apply at a rate of 30 to 40 pounds per acre as a broadcast application. Do not apply within 21 days of harvest. MUSIARD_GREENS: To control cutworms, wireworms, and white grubs at a rate of 25 rounds mer acre as a broadcast soil application. Do not apply within 10 days of harvest. OKRA: To control cutworms and wireworms apply at a rate of 30 to 40 pounds per acre as a broadcast soil application. Do not apply within 21 days of harvest. QNIQNS: To control cutworms and wireworms apply at a rate of 30 to 40 pounds per acre as a broadcast soil application. Do not apply within 15 days of harvest. To control white grubs, apply at a rate of 30 to 40 pounds per acre PEANUIS: as a broadcast soil application. Mix into top 3 to 4 inches of soil immediately following application. Apply 2 weeks before planting. To control wireworms, apply at a rate of 50 to 60 rounds per acre at time of planting or pessing, incorporating lightly into soil. To control Southern corn rootworm apply at a rate of 30 to 40 pounds per acre in a 12 inch band by ar the row Just before ressins time, or apply as a preplant broadcast soil treatment and incorporate into top soil. To control lesser cornstalk bongs, apply at a rate of 30 to 40 pounds per acre in a 12 inch bank over the row Just before or after ressing. If applied before to ressing incorporate limto the top few inches of soil. If applied after pessins, apply before irrigation and thoroughly water into soil. 2EPPERS: To control cutworms and wireworms apply at a rate of 30 to 40 mounds Do not apply within 15 days of harvest. per acre as a broadcast application. PDIAIDES: To control cutworms and wireworms apply at a rate of 30 to 40 pounds per acre as a broadcast application. Do not apply within 5 days of harvest. 2PINACH: To control cutworms and wireworms apply at a rate of 25 pounds per

acre as a broadcast application. Do not apply within 14 days of hervest.

 $^{\circ}$ SQUASH: To control cutworms, wireworms, and white grubs, apply at a rate of 25 pounds per acre as a broadcast soil application. Do not apply within 15 days of harvest.

SIRAMBERRIES: To control cutworms and wireworms apply at a rate of 30 to 40 pounds per wore as a broadcast soil application. Bo not apply within 14 days of harvest. As a preplant broadcast soil treatment apply 2-3 weeks before planting and immediately incorporate into top 6 inches of soil. Use lower dosade for mineral soils and higher dosade for organic soils.

SUGAR_BEEIS:_SHEET_POIATOES:_SUGARCANE: To control wireworms and cutworms apply as a preplant broadcast soil treatment. Apply 2-3 weeks before planting and immediately incorporate into top 6 inches of soil. Use lower dosage for inches a soil and higher dosage for organic soils.

'IDMAIDES: To control cutworms and wireworms apply at a rate of 30 to 40 pounds per acre as a broadcast application. Do not apply within 10 days of harvest.

IDBACCO: To control wireworms apply at a rate of 30 to 40 pounds per acre, Do not apply within 5 days of harvest (priming) or 15 days of harvest (by cutting). Avoid contact with plant juices when priming or cutting tobacco, when necessary to enter treated fields within 48 hours after application, protective clothing must be worn.

:IURNIES: To control cutworms, wireworms, and white grubs, apply at a rate of 25 pounds per acre as a broadcast soil application. Do not apply within 10 days of harvest.

DIHER_USES

-ORNAMENIALS: For control of wireworms on sladiolus and flowerins plants, apply at a rate of 30 to 40 pounds per acre. For use by commercial growers only. Apply in furrow over bulbs at time of planting.

LIMIT OF WARRANTY AND LIABILITY

This company warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes set forth in the Complete Directions for Use label booklet ("Birections") when used in accordance with those Directions under the conditions described herein. NO OTHER EXPRESS OR IMPLIED WARRANTY OF MERCHANTABILITY AND FITNESS FOR PARTICULAR PURPOSE IS EXCLUDED. This warranty is also subject to the conditions and limitations stated herein.

Buyer and all users shall promptly notify this company of any claims whether based in contract, negligence, strict liability, other tort or otherwise. Buyer and all users are responsible for all loss or damage from use or handling which results from conditions beyond the control of this company, including but not limited to, incompatibility with products other than those set Yorth in the Directions, and application in any manner, not explicitly set forth the Directions.

THE EXCLUSIVE REMEDY OF THE USER OF BUYER, AND THE LIMIT OF THE LIABILITY OF THIS COMPANY OR ANY OTHER SELLER FOR ANY AND ALL LOSSES, INJURIES OF DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT (INCLUDING CLAIMS BAGSE, IN CONTRACT, NEGLIGENCE, STRICT LIABILITY, OTHER TORT OR OTHERWISE) SHALL BE THE PURCHASE PRICE PAID BY THE USER OR BUYER FOR THE QUANTITY OF THIS PRODUCT INVOLVED, OR, AT THE ELECTION OF THIS COMPANY OR ANY OTHER SELLER, THE REPLACEMENT OF SUCH QUANTITY OR, IF NOT ACQUIRED BY PURCHASE, REPLACEMENT OF SUCH QUANTITY, IN NO EVENT SHALL THIS COMPANY OR ANY OTHER SELLER BE LIABLE, FOR ANY INCIDENTAL OR CONSEQUENTIAL DAMAGES.

The buwer and all users are deemed to have accepted the terms of this LIMIT OF WARRANTY AND LIABILITY which may not be varied by any verbal or written agreement.

1

y got 9