

1 of 14
Pm 25
9859-273

Center Panel

ATRAZINE 4L

For Season Long Weed Control in Corn, Sorghum and
Certain Other Crops

ACTIVE INGREDIENTS:

Atrazine (2-chloro-1-ethylamino-6-isopropylamino- s-triazine)	10.80%
Related compounds	2.20%
INERT INGREDIENTS	87.00%
TOTAL	100.00%

Contains 4 pounds active ingredient per gallon.

KEEP OUT OF REACH OF CHILDREN

CAUTION

ACCEPTED

AUG - 4 1987

Under the Federal Insecticide,
Fungicide, and Rodenticide Act,
as amended, for the pesticide
registered under
EPA Reg. No. 9859-273

STATEMENT OF PRACTICAL TREATMENT

IF SWALLOWED: Call a physician or Poison Control Center. Drink 1 or 2 glasses of water and induce vomiting by touching back of throat with finger. Do not induce vomiting or give anything by mouth to an unconscious person.

IF ON SKIN: Wash with plenty of soap and water.

IF IN EYES: Flush with plenty of water. Get medical attention if irritation persists.

IF INHALED: Remove victim to fresh air. If not breathing, give artificial respiration, preferably mouth-to-mouth. Get medical attention.

See Side Panel For Additional Precautionary Statements

EPA Acc. No. 9859-6216

EPA Est. No. 9859-273-1

Manufactured By
LAMBDA CHEMICAL COMPANY
LAKELAND, FLORIDA 33802

NET CONTENTS:

BEST AVAILABLE COPY

(Left Panel)

PRECAUTIONARY STATEMENTS

Hazards To Humans And Domestic Animals

CAUTION: Harmful if swallowed, inhaled, or absorbed through the skin. Do not breathe vapors or spray mist. Avoid contact with eyes, skin or clothing. Wash thoroughly after handling.

ENVIRONMENTAL HAZARDS

This pesticide is toxic to aquatic invertebrates. Do not apply directly to water or wetlands. Runoff and drift from treated areas may be hazardous to aquatic organisms in neighboring areas. Do not contaminate water by cleaning of equipment or disposal of wastes.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

RE-ENTRY STATEMENT

Do not enter treated areas without protective clothing until sprays have dried.

Because certain states may require more restrictive reentry intervals for various crops treated with this product, consult your State Department of Agriculture for further information.

Do not apply this product in such a manner as to directly or through drift expose workers or other persons. The area being treated must be vacated by unprotected persons.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by other procedures approved by state and local authorities.

GENERAL INFORMATION

Atrazine 4L herbicide controls most annual broadleaf and grass weeds in corn, sorghum and certain other crops.

Apply at the rates given under each crop before or after weeds emerge. Use lower rates on light soils and low organic matter soils. Use higher rates on heavy soils and high organic matter soils.

Atrazine 4L acts mainly through root absorption so rainfall or irrigation is needed to move it into the root zone. Should weeds develop, a shallow cultivation or rotary hoeing will generally result in better weed control.

BEST AVAILABLE COPY

Avoid using Atrazine 4L near adjacent desirable trees, shrubs, or plants, or injury may occur.

NOTE: Landia does not recommend Atrazine 4L in combination with other herbicides or oils except as specifically described on the label.

Atrazine 4L is non-flammable.

APPLICATION PROCEDURES

Atrazine 4L, a liquid formulation, should be mixed with water or liquid fertilizer and applied as a spray. Pour Atrazine 4L into the tank during or after filling Hydraulic (jet) or mechanical agitation is recommended during mixing and application to keep the material in suspension. All return lines to the tank must discharge below liquid level and agitation should not be so violent as to cause air bubbles to form in the liquid. Wash sprayer thoroughly after use.

Ground Application: Apply Atrazine 4L in 10-40 gallons of water per acre to assure even coverage unless otherwise specified. Use TeeJet 8002, 8003, or 8004 fan type or other appropriate nozzles with equal or greater size openings for broadcast applications. Use TeeJet 8002-E, or 8004-E or similar nozzles for band applications. Screens in nozzles and in suction and in-line strainers should be no finer than 50-mesh. Use a pump with capacity sufficient to maintain 35-40 psi operating pressure.

Aerial Application: Use only where broadcast applications are specified. Apply a minimum of 1 gallon of water for each 2 pints of Atrazine 4L to be applied per acre. Do not apply where uniform coverage cannot be obtained or where excessive spray drift may occur. Avoid application directly to animals or humans. Flamen or loaders should avoid inhalation of spray mist or contact with skin and should wash thoroughly before eating and at the end of each day's operation.

Post-emergence Weed Control with Atrazine 4L plus Emulsifiable Oil in Water: The addition of oil to Atrazine 4L in water sprays may speed the activity of Atrazine 4L and provide quicker kill of weeds. However, under certain conditions the use of oil may seriously damage corn or sorghum. To minimize the possibility of damage, follow the directions, procedures, and precautions given below.

For combining with Atrazine 4L, use a crop oil designated for use with Atrazine 4L containing at least 1% suitable emulsifier. Oils contaminated with water or other materials can cause compatibility problems and/or crop injury.

EQUIPMENT PROCEDURES FOR ATRAZINE 4L AND EMULSIFIABLE OIL:

1. Thoroughly clean sprayer prior to use. Do not use sprayer contaminated with 2,4-D or other materials as crop damage or sprayer clogging may result.
2. Fill the spray tank with water, allowing for the addition of the Atrazine 4L.

BEST AVAILABLE COPY

3. With the agitation system in operation, add the Atrazine 4L.
4. Add the emulsifiable oil last.
5. Empty the tank as completely as possible before refilling to prevent a buildup of oil in the tank. Keep the agitation system in operation to avoid a separation of the oil from the water in the spray mixture still remaining in the tank.
6. If an oil film starts to build up in the tank, drain the tank and clean out with a strong detergent solution or solvent.
7. Clean the sprayer thoroughly immediately after use by flushing the system with water containing a detergent.

NOTE:

1. Do not use oil in Atrazine 4L sprays when corn or sorghum is under stress from prolonged cold, wet weather, poor fertility, or other factors or when corn or sorghum is wet and succulent from recent rainfall as crop injury may occur.
2. Do not use oils in sprays when treating inbred lines or any breeding stock as injury may occur.
3. Adding other insecticides, herbicides, liquid fertilizers, or other materials is not recommended because they may cause compatibility problems or crop injury.
4. Store and handle emulsifiable oil carefully. Oil contaminated with even a small amount of water may not emulsify properly when added to the tank.
5. Do not make more than one application of Atrazine 4L and emulsifiable oil in water per season.
6. All return lines to the tank must discharge below liquid level.

BEST AVAILABLE COPY

(Right Panel)

CORN

One application will control most annual broadleaf grass weeds such as barnyardgrass, witchgrass (Panicum capillare), yellow foxtail, green foxtail, wild oats, large (hairy) crabgrass, giant foxtail, velvetleaf, morningglory, lambsquarters, pigweed, ragweed, nightshade, purslane, and mustard.

Atrazine 4L may be applied either before planting, at planting or after planting at the rate indicated below.

For preplant and pre-emergence applications, nitrogen solutions or complete liquid fertilizers may replace all or part of the water as a carrier of Atrazine 4L. Do not apply after corn has emerged as there is danger of liquid fertilizers causing crop injury.

Preplant: Broadcast in the spring after plowing at the rate indicated below. Apply before, during, or after final seedbed preparation. If soil is tilled or worked after application, avoid deep incorporation of Atrazine 4L. Best results have been obtained when Atrazine 4L is applied within two weeks prior to planting.

Pre-emergence: Apply during or shortly after planting prior to weed emergence at the rate indicated below.

Post-emergence: Apply before weeds exceed 1-1/2 inches in height at the rate indicated below.

Rate per acre of Atrazine 4L

Soil texture	Broadcast in 10-40 gal. of water/A	20-inch band+ in 5-20 gal. of water/A	13-inch band+ in 4-14 gal. of water/A
Light soils: Sands, loamy sands, and sandy loams	4 pints	2 pints	1-1/4 pints
Medium soils: Silt and clay loams that are low in organic matter	4-3/4 pints	2-1/2 pints	1-1/2 pints
Heavy soils: Silt and clay loams with medium to high organic matter & clays	6 pints	3 pints	2 pints
Peats, mucks, and high organic clays (Apply post-emer-	6 pints	3 pints	2 pints

BEST AVAILABLE COPY

gence only)

+Based on 40-inch row spacing. For other band or row widths, adjust the rate in proportion to the area actually treated.

Northern Nutgrass or Yellow Nutsedge (Cyperus esculentus) Control: Broadcast 8 pints per acre preplant, pre-emergence, or post-emergence before nutgrass is more than 1-1/2 inches high. If applied before nutgrass begins growth, cultivate within 2-3 weeks. If applied after nutgrass has emerged, cultivate within 10-14 days.

Quackgrass Control on Land Going Into Corn Production: Split Application: Broadcast 4 pints per acre in the fall or spring and plow 1-3 weeks later. Broadcast a second application at the rate of 4 pints per acre in the spring before, during, or after planting, but before weeds are 1-1/2 inches high.

Single Application: Broadcast 6-8 pints per acre in the fall or spring. Plow 1-3 weeks after application.

Post-emergence weed control with Atrazine 4L plus emulsifiable oil in water:

Ground Application: Broadcast 4 pints of Atrazine 4L and 1 gallon of emulsifiable oil in 20-40 gallons of water per acre after emergence but before weeds reach 1-1/2 inches in height.

Aerial Application: Broadcast 4 pints of Atrazine 4L and 1/2 gallon of emulsifiable oil in no less than 5 gallons of water per acre after weed emergence but before weeds reach 1-1/2 inches.

PRECAUTIONS

1. Do not apply more than 8 pints of Atrazine 4L to corn in any one year.
2. Land treated with Atrazine 4L should not be planted to any crop except corn until the following year or injury may occur.
3. Following harvest of a treated crop, plow (moldboard or diskplow) and thoroughly till the soil in fall or spring to minimize possible injury to rotational spring-seeded crops, regardless of the rate used.
4. If Atrazine 4L is applied after June 10, do not rotate with crops other than corn or sorghum the next year or injury may occur.
5. If Atrazine 4L is used at a broadcast rate higher than 6 pints per acre (or comparable rates in a band application), a crop of untreated corn or sorghum should precede the next rotational crop.
6. In the High Plains and Intermountain areas of the West where rainfall is sparse and erratic or where irrigation is required, use Atrazine 4L to control weeds in corn only when corn is to follow corn or a crop of untreated corn or sorghum is to precede other rotational crops.
7. In western Minnesota and eastern parts of the Dakotas, Nebraska, and Kansas, corn treated with Atrazine 4L should not be followed with soybeans if the broadcast rate applied was more than 4 pints per acre (or comparable rate in a band) or injury may occur.

- 8. Do not plant sugar beets, tobacco, vegetables (including dry beans), spring-seeded small grains, or small-seeded legumes and grasses the year following Atrazine 4L application or injury may occur.
- 9. Do not graze treated or feed treated forage to livestock for 21 days following application.

TANK MIXTURES ON CORN

Dual* 6E

Use as tank mixture for control of certain broadleaf and grass weeds in corn. Refer to Dual* 6E label for all directions, weeds controlled, precautions, and limitations.

Sim-Trol 9DF or Sim-Trol 4L

Use as tank mixture for control of many annual weeds, including carpet-weed, crabgrass, fall panicum, foxtail, lambsquarters, morningglory, pigweed, ragweed, and velvetleaf. Apply before planting, at planting, or after planting, but before crop and weeds emerge, at rates in Table 2a. Use 1:1 ratio for control of above weeds. Use 1:2 ratio for expected heavy infestations of crabgrass and fall panicum. Apply in a minimum of 20 gallons of water/A. Cultivate shallowly if weeds develop.

Preplant: Apply in spring during or after final seedbed preparation. If soil is tilled or worked after application, avoid deep incorporation. For best results, apply within two weeks before planting.

Preemergence: Apply during or shortly after planting, but before crop and weeds emerge.

Refer to Corn sections of this label and to Sim-Trol 9DF or Sim-Trol 4L label for further directions, Limitations and cautions.

T A B L E 2a
TANK MIXTURES WITH SIMAZINE ON CORN

Soil Texture	1:1 RATIO*		1:2 RATIO**	
	This Product	Sim-Trol 9DF or Sim-Trol 4L	This Product	Sim-Trol 9DF or Sim-Trol 4L
Sand, loamy sand, sandy soil	2 pt.	1.1 lb. 2 pt.	1.32 lb.	1.4 lb. 2.67 pt.
Loam, silt loam, silt, clay loam, sandy clay loam, silty clay loam, sandy clay, or silty clay with low organic matter	2.4 pt.	1.3 lb. 2.4 pt.	1.6 pt.	1.7 lb. 3.2 pt.

BEST AVAILABLE COPY

Loam, silt loam, silt, clay loam, sandy clay loam, silty clay loam, sandy clay, or silty clay with medium to high organic matter, & clay (including dark prairie soils of the Corn Belt)	3 pt.	1.5 lb.	1.92 pt.	2.1 lb.
		3 pt.		3.84 pt.

*For control of above weeds

**For control of expected heavy infestations of crabgrass and fall panicum.

Preplant and Pre-emergence Application
(In Eastern and Central Nebraska and Northeastern Kansas Only)

Broadleaf and grass weed control

Rate per acre of Atrazine 4L

Soil Texture	Broadcast in 10-40 gal. of water/A	20-inch Band+ in 5-20 gal. of water/A	13-inch Band+ in 4-14 gal. of water/A
Sand to sandy loam, eroded slopes and soils low in organ- ic matter		Do not use	
Other soils (shallow incorpor- ation may improve control)	4 to 4-3/4 pints	2 to 2-1/2 pints	1-1/4 to 1-1/2 pints

+Based on 40-inch row spacing. When making band applications, reduce the broadcast rate per acre in proportion to the area treated.

Post-emergence: For the control of broadleaves and grasses, apply post-emergence applications of Atrazine 4L in no less than ten gallons of water per acre at the rate indicated in the table below. Weeds should not exceed 1-1/2 inches in height at application.

Post-emergence Application++
(weeds should not be more than 1-1/2 inches high)

Rate per acre of Atrazine 4L

20-Inch 13-Inch

BEST AVAILABLE COPY

Weeds to be controlled and soil texture	Stage or height of sorghum at treatment	Broadcast in 10-40 gal. of water/A	band+ in 5-20 gal. of water/A	band+ in 4-14 gal. of water/A
---	---	------------------------------------	-------------------------------	-------------------------------

Broadleaf and grass weeds

Sand or loamy sand

Do not use

Sandy loam

See broadleaf weeds below

Silt loam to Silt loam to sandy clay loam

Completely emerged

4 to 4-3/4 pings

2 to 2-1/2 pints

1-1/4 to 1-1/2 pints

Olton and Pullman clay soils

At least 6 inches high

4 to 4-3/4 pint

2 to 2-1/2 pints

1-1/4 to 1-1/2 pints

Silty clay loams and heavier soils

Completely emerged

6 pints

3 pints

2 pints

Broadleaf weeds

Sandy loams and heavier soils in Oklahoma, New Mexico, Texas, West Kansas, Colorado, and desert regions of Arizona and California only

At least 6 inches high

2-1/2 pt. plus 3/4 - 1-1/2 pints surfactant

1-1/4 pt. plus 3/8 - 3/4 pint surfactant

3/4 pt. plus 1/4 - 1/2 pt. surfactant

+Based on 40-inch row spacing.

++Up to "close-in".

Post-emergence broadleaf weed control with Atrazine 4L plus emulsifiable oil in water (New Mexico, Oklahoma, and Texas only): Apply when sorghum is 6-10 inches high and before boot stage. Treat broadleaf weeds before they reach 6 inches.

Ground Application: Broadcast 2-1/2 pints of Atrazine 4L and 1/2 - 1 gallon of emulsifiable oil in 20-40 gallons of water per acre.

Aerial Application: Broadcast 2-1/2 pints of Atrazine 4L and 1/2 gallon of emulsifiable oil in no less than 5 gallons of water per acre.

PRECAUTIONS

BEST AVAILABLE COPY

1. Heavy rains immediately following application tend to result in excessive concentrations of herbicide in seed furrow, resulting in possible crop injury. Applications to furrow-planted sorghum should not be made until furrows are leveled (plowed-in). Deep planter marks or seed furrows should also be leveled before application.
2. Application made to sorghum growing under stress caused by minor element deficiency or to sorghum growing on highly calcareous soils may result in crop injury.
3. Do not plant any rotation crops other than those specified in the Suggestions For Rotational Crops Section below for 18 months following application.
4. Do not graze or feed forage from treated areas for 21 days following application.
5. Following harvest of a treated crop, plow (moldboard or disk-plow) and thoroughly till the soil in fall or spring to minimize possible injury to rotational spring-seeded crops, regardless of the rate used.

SUGGESTIONS FOR ROTATIONAL CROPS:

1. Sorghum or corn may be planted the year following treatment.
2. All other rotational crops may be planted 18 months after application.
3. When the broadcast rate used does not exceed 2-1/2 pints per acre, fall wheat may be planted 15 months after application.

TANK MIXTURES ON GRAIN SORGHUM

Igran* 80W

Use as tank mixture for control of certain broadleaf and grass weeds in grain sorghum. Refer to Igran* 80W label for all directions, weeds controlled, precautions, and limitations.

SUGARCANE

For control of many annual broadleaf and grass weeds, including amaranths, crabgrass, fireweed, Florida's paintbrush, foxtails, junglerice and wiregrass:

Broadcast 2-4 quarts in 20-50 gallons of water/A for adequate coverage of the soil surface at time of planting or ratooning, but before cane emerges. One additional application may be made over cane as it emerges, and two additional applications may be made interline after emergence as directed sprays.

For control of emerged pellitory weed (Florida only): Apply 0.8-1.2 pints in at least 40 gallons of water/A as a directed spray. Add 2 quarts of surfactant for each 50 gallons of spray. Thoroughly cover weed foliage.

Precautions: (1) Do not apply after "close-in". (2) Do not apply more than 10 quarts/A to any one crop of cane.

For control of alexandergrass, large crabgrass, pellitory (artillery) weed and spiny amaranth, use only one of the following methods at

planting or ratooning (Florida only).

1. Apply 4 quarts in 20-50 gallons of water/A, preemergence, broadcast or banded over-the-top applications as needed, postemergence to sugarcane and weeds, at 2 quarts in 20-50 gallons of water/A. Treat before weeds exceed 1.5 inches in height.
2. Apply 1-3 times as needed, at 2 quarts in 20-50 gallons of water/A, either broadcast or banded over-the-top, post-emergence to sugarcane and weeds. Treat before weeds exceed 2.5 inches in height.

CHEMICAL FALLOW

WHEAT-SORGHUM FALLOW

This treatment controls annual broadleaf and grass weeds following wheat harvest and in the following sorghum crop when grown under minimum tillage.

Apply 6 pt. to wheat stubble immediately following wheat harvest. If weeds are present, remove them with a sweep plow or other suitable implement after application. Plant sorghum into wheat stubble the following spring with minimum disturbance of the soil. Use a surface planter or a planter leaving a shallow furrow. If weeds are present at planting, remove them with a sweep plow or other suitable implement before planting.

Precautions: (1) Use only on silt loam or finer textured soil. (2) Wheat-sorghum-fallow cropping sequence must be followed. (3) Do not apply following sorghum harvest.

Note: Do not graze or feed forage from treated area to livestock. Do not plant any crop other than those on this label within 18 months following treatment.

WHEAT-CORN-FALLOW (KS, NE)

This product controls cheatgrass (downy brome, chess), kochia, mustards, pigweed, Russian thistle, wild lettuce, wild sunflower, and volunteer wheat during period after wheat harvest. Weed control may extend into following corn crop grown under minimum tillage.

Apply with ground equipment only.

Follow directions for use, notes, and precautions in the "Wheat-Sorghum-Fallow" section above, substituting corn for references to sorghum.

WHEAT-FALLOW-WHEAT (CO, KS, MT, NE, ND, SD, AND WY)

For preemergence control of cheatgrass (downy brome, chess), common lambsquarters, field pennycress, kochia, mustard, Russian thistle, wild lettuce, and suppression of volunteer wheat during fallow period of a wheat-fallow-wheat rotation, apply 1-2 pt. in 10-40 gal. of water/A for ground application on all soils except those listed under "Precautions." For control of pigweed and wild sunflower, use the higher rate. Apply to stubble ground. Treat only once during same fallow period.

If weeds are present at application and this product is used alone, use either an approved contact herbicide before or after treatment, or tillage after treatment.

Use tillage to control weeds which escape during fallow period. Till before planting. For this product applied alone or in tank mixture with Paraquat CL, plant at least 2 inches deep and 12 months or more after application.

Precautions: (1) Do not use on sand soil. (2) Do not treat eroded hillsides, caliche and rocky outcroppings, or exposed calcareous sub-soil. (3) Do not treat soils of the Rosebud and Canyon Series in western NE and adjoining counties in CO and WY. (4) Do not treat soils with calcareous surface layers. (5) Avoid spray overlap.

Note: Do not graze treated areas within 6 months after applications.

RANGELAND

This product controls certain weeds, including cheatgrass (downy brome, chess), tumble mustard, safflower, and annual or common broomweed. Use lower rate where cheatgrass is major problem weed; use higher rate for other weeds.

For ground applications, use a minimum of 10 gal. of water/A. For Aerial applications, use a minimum of 5 gal. of water/A.

See "Application Procedures" for additional information.

ESTABLISHMENT OF PERENNIAL RANGE GRASSES (CA, ID, NV, OR, UT, WA)

Apply 1.6-2 pt./A in fall after rains begin but before ground freezes. Perennial range grass may be seeded when rain begins the next fall following treatment. Plant in deep furrows such as those made by a deep furrow rangeland drill. Treated areas may be grazed 7 months after seedling. However, to provide sufficient time for establishment of desirable range grasses, treated areas should not be grazed until the fall following seedling.

Renovation of existing stands of perennial range grasses Fall treatment in Central Great Plains (CO, KS, MT, NE, ND, SD, WY): Apply 1.6-2 pt./A in fall before ground freezes.

Spring treatment in Southern Great Plains (OK, TX): Apply 2 pt/A in spring before April 30.

Do not apply to crested and intermediate wheatgrasses. Do not cut or feed range grass hay. Do not graze treated areas within 7 months following fall application or 3 months following spring application. Apply only once per year.

**Warm Season Perennial Range Grasses - Midwest
Switchgrass**

For control of many annual broadleaf grass weeds including cheatgrass, foxtails (giant, green, and yellow), lambsquarters, pigweed, ragweed,

smartweed, and velvetleaf.

For ground application use a minimum of 15 gal. of water/A.

Use only on loam, silt loam, silty clay loam, and silty clay soils with at least 1 percent organic matter.

New seeding: Apply 2 qt/A after planting but before weeds emerge. Prepare a good, firm seedbed. Plant 1/2 inch deep with a grassland drill (preferred method) or a conventional drill. If a conventional drill is used on prepared seedbeds, remove all tension from the disk openers. For best results, siltipack or roll after planting. Clip weeds that escape in July or August. Avoid clipping switch-grass seedlings.

Established stands: Apply 2 qt/A in April or early May, before weed emergence.

Rotational crops: See "Rotational crops" in Corn section

Note: Do not cut for hay. Do not graze treated areas within 4 months following application at seedling or 3 months following application to established switchgrass. Treat only once per year.

NONSELECTIVE WEED CONTROL ON NONCROPLAND

This product provides long term weed control in industrial sites, highway medians and shoulders, railroad right-of-ways, lumber yards, petroleum tank farms and in noncrop areas on farms, such as around buildings, equipment and fuel storage areas, along fences and lanes.

Apply before or soon after weeds begin growth. Make postemergence applications when weeds are young and actively growing. Use the higher rates on fine clay and muck soils. Use sufficient water to assure thorough coverage.

Annual broadleaf and grass weeds (including barnyardgrass, cheatgrass, crabgrass, lambsquarters, foxtail, ragweed puncturevine and turkey mullein). Apply 1.5-2.5 gallons per acre.

Hard-to-kill annual and perennial broadleaf and grass weeds (including bluegrass, burdock, Canada thistle, dogfennel, orchardgrass, plantain, quackgrass, purpletop, redtop, and smooth brome). Apply 2.5-5 gallons per acre.

Hard-to-kill biennial and perennial weeds (including bull thistle and sowthistle): Apply 5-7.5 gallons per acre.

For longer residual control in regions of high rainfall and a long growing season, apply 5-7.5 gallons per acre.

*Dual trademark of Ciba-Geigy for metholachlor U.S. patent No. 3,937,730.

*Iqran trademark of Ciba-Geigy for terbutryn U.S. patent No. 3,634,062.

CONDITIONS OF SALE

All statements concerning the use of this product apply only when used as directed. THE MANUFACTURER MAKES NO WARRANTIES, EXPRESSED OR IMPLIED, CONCERNING THIS PRODUCT OR ITS USE, WHICH EXTEND BEYOND THE DESCRIPTION ON THE LABEL. Read all directions carefully.