

195 22  
PM 25  
-1859-274

(Center Panel)

**ATRAZINE 60W**

For season-long weed control in corn and sorghum  
For weed control in certain other crops;  
in noncrop areas; and industrial sites

**ACTIVE INGREDIENTS:**

Atrazine (2-chloro-4-ethylamino-6-isopropylamino- s-triazine .....	76.00%
Related Compounds .....	4.00%
<b>INERT INGREDIENTS</b> .....	20.00%
<b>TOTAL</b> .....	100.00%

**KEEP OUT OF REACH OF CHILDREN**

**C A U T I O N**

**STATEMENT OF PRACTICAL TREATMENT**

**IF SWALLOWED:** Call a physician or Poison Control Center. Drink 1 or 2 glasses of water and induce vomiting by touching back of throat with finger. Do not induce vomiting or give anything by mouth to an unconscious person.

**IF ON SKIN:** Wash with plenty of soap and water.

**IF IN EYES:** Flush with plenty of water. Get medical attention if irritation persists.

**IF INHALED:** Remove victim to fresh air. If not breathing, give artificial respiration, preferably mouth-to-mouth. Get medical attention.



See Side Panel For Additional Precautionary Statements

EPA Acc. No. 9859-ETU

EPA Est. No. 51036-GA-1

Manufactured For:  
**LANDIA CHEMICAL COMPANY**  
LAKELAND, FLORIDA 33802

NET CONTENTS: \_\_\_\_\_



(Left Panel)

PRECAUTIONARY STATEMENTS

Hazards To Humans And Domestic Animals

CAUTION: Harmful if swallowed, inhaled, or absorbed through skin. Do not breathe dust or spray mist. Avoid contact with eyes, skin or clothing.

NOTE TO PHYSICIAN: There is no specific antidote for atrazine. If this product is ingested, induce emesis or lavage stomach. The use of an aqueous slurry of activated charcoal may be considered.

GROUNDWATER: Atrazine leaches readily and accepted label rates have been found to result in contamination of water supplies by way of groundwater. Therefore, users are advised to avoid use of atrazine in all-drained soils, particularly in areas having high groundwater tables.

ENVIRONMENTAL HAZARDS

This pesticide is toxic to aquatic invertebrates. Do not apply directly to water or wetlands. Runoff and drift from treated areas may be hazardous to aquatic organisms in neighboring areas. Do not contaminate water by cleaning of equipment or disposal of wastes.

RE-ENTRY STATEMENT

Do not enter treated areas without protective clothing until sprays have dried.

Because certain states may require more restrictive reentry intervals for various crops treated with this product, consult your State Department of Agriculture for further information.

Do not apply this product in such a manner as to directly or through drift expose workers or other persons. The area being treated must be vacated by unprotected persons.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal. Store in a dry place.

PESTICIDE DISPOSAL

Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL

Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

GENERAL INFORMATION

This herbicide controls many annual broadleaf and grass weeds in corn, sorghum, sugarcane, pineapple, and certain other crops specified on this label. It is also effective in noncrop areas and industrial sites for control of most annual and many perennial broadleaf and grass weeds. This product may be applied before or after weeds emerge.

BEST AVAILABLE COPY

Following many years of continuous use of this product and chemically related products, biotypes of some of the weeds listed on this label have been reported which cannot be effectively controlled by this and related herbicides. Where this is known or suspected, we recommend the use of this product in combination with other registered herbicides which are not triazines. Consult with your State Agricultural Extension Service for specific recommendations.

Since this product acts mainly through root absorption, its effectiveness depends on moisture to move it into the root zone. If weeds develop, a shallow cultivation or rotary hoeing will generally result in better weed control.

This product is noncorrosive to equipment and metal surfaces, nonflammable, and has low electrical conductivity.

Avoid using near adjacent desirable plants or in greenhouses, or injury may occur.

Where the use directions give a range of rates, use the lower rate on coarse-textured soil and soil low in organic matter; use the higher rate on fine-textured soil and soil high in organic matter.

NOTE: Micro-Flo does not recommend applications in combination with other herbicides or oils, except as specifically described on the label.

APPLICATION PROCEDURES

GROUND APPLICATION: Use conventional ground sprayers equipped with nozzles that provide accurate and uniform application. Be certain that nozzles are uniformly spaced and are the same size. Calibrate sprayer before use and recalibrate at the start of each season and when changing carriers. Unless otherwise specified, use a minimum of 10 gallons of spray mixture per acre for all preplant, preemergence, and postemergence applications (with or without soil or surfactant) with ground equipment.

Use a pump with capacity to (1) maintain 35-40 psi at nozzles, (2) provide sufficient agitation in tank to keep mixture in suspension, and (3) to provide a minimum of 20% bypass at all times. Use centrifugal pumps which provide propeller shear action for dispersing and mixing this product. The pump should provide a minimum of 10 gallon minute 100 gallon tank size circulated through a correctly positioned sparger tube or jets.

Use screens to protect the pump and to prevent nozzles from clogging. Screens placed on the suction side of the pump should be 16-mesh or coarser. Do not place a screen in the recirculation line. Use 50-mesh or coarser screens between the pump and boom, and where required, at the nozzles. Check nozzle manufacturer's recommendations.

For band applications, calculate amount to be applied per acre as follows:

BEST AVAILABLE COPY

band width in inches  
-----  
row width in inches

X

broadcast  
rate  
per acre

=

amount needed  
per acre of field

BEST AVAILABLE COPY

**BEST AVAILABLE COPY**

**AERIAL APPLICATION:** Use aerial application only where broadcast applications are specified. Apply in a minimum of 1 gallon of water for each 1 pound of Atrazine 80W applied per acre. For postemergence treatments on corn and sorghum, apply recommended rate in a minimum of 2 gallon of water/A. Avoid applications under conditions where uniform coverage cannot be obtained or where excessive spray drift may occur.

Avoid application to humans or animals. Flagmen and loaders should avoid inhalation of spray mist and prolonged contact with skin, and should wash thoroughly before eating and at the end of each day's operation.

**APPLICATION IN WATER OR LIQUID FERTILIZER:** Nitrogen solution or complete liquid fertilizer may replace all or part of the water as a carrier for preemergence or preplant ground application on corn and sorghum. Check the compatibility of this product with liquid fertilizer and/or nitrogen solution as shown below before use. Do not apply in nitrogen solution or complete liquid fertilizer after corn or sorghum emerges, except as noted under LAY-BY TREATMENT for corn, or crop injury may occur.

To determine the compatibility of Atrazine 80W, alone or in tank mixtures, with fluid fertilizer, pour the products into a small container of fluid fertilizer in the proportions shown below. For different spray volume or herbicide rate, make appropriate changes in the ingredients.

For each pound of Atrazine 80W or other dry product to be applied per acre, add 1 level teaspoon or 5 milliliters of the product to one pint of fluid fertilizer (assuming a volume of 25 gal/A).

For each pint of a liquid or flowable product to be applied per acre, add 0.5 teaspoon or 2.5 milliliters of the product to one pint of fluid fertilizer (assuming a volume of 25 gal/A).

After thorough mixing, let stand for 5 minutes. If the herbicide fertilizer combination remains mixed, or can be remixed readily, the mixture is compatible and can be sprayed. If it is not compatible, make a water slurry with the herbicide before adding fertilizer and use the mixture only if this tests shows it is compatible.

**APPLICATION IN WATER PLUS EMULSIFIABLE OIL OR OIL CONCENTRATE:** Adding emulsifiable oil or oil concentrate to postemergence water-based sprays on corn and sorghum may improve weed control. However, under certain conditions, the use of either type of oil may seriously injure the crop. To minimize this possibility, observe the following directions.

Use a crop oil containing 1-2% suitable emulsifier, or a suitable crop oil concentrate containing not more than 20% emulsifier or surfactant blend. Emulsifiable oil and oil concentrate contaminated with water or other materials can cause compatibility problems and/or crop injury.

**MIXING PROCEDURES - ALL USES:** (1) Be sure sprayer is clean and not

BEST AVAILABLE COPY

contaminated with any other materials, or crop injury or sprayer clogging may result. (2) Fill tank 1/4 full with clean water, nitrogen solution, or complete liquid fertilizer. (3) Start agitation. (4) Be certain that the agitation system is working properly and creates a rippling or rolling action on the liquid surface. (5) Pour product directly from bag into tank. (6) Continue filling tank until 90% full. Increase agitation if necessary to maintain surface action. (7) Add emulsifiable oil, oil concentrate, or tank mix herbicides after this product is thoroughly suspended. (8) Finish filling tank. (9) Empty tank as completely as possible before refilling to prevent buildup of oil or emulsifiable concentrate residue. Maintain agitation to avoid separation of materials. (10) If an oil or emulsifiable concentrate film starts to build up in tank, drain it and clean with strong detergent solution or solvent. (11) Clean sprayer thoroughly immediately after use by flushing system with water containing a detergent.

**CENTER PIVOT SPRINKLER APPLICATION:** Preemergence or postemergence: Apply Atrazine 80W alone, or in tank mixtures with other herbicides on this label which are registered for center pivot application, with irrigation water. Apply either after planting before crop and weeds emerge, or after crop emergence, but before lay-by (20-30 inches) and before weeds exceed 1.5 inches in height. Apply at rates recommended on this label. Prepare mixture with a minimum of 1 part water to 1 part product. Injecting a larger volume of a more dilute slurry per hour will assure more accurate calibration of metering equipment. Maintain sufficient agitation to keep herbicide in suspension. Meter slurry into irrigation water during entire period. Apply in 1/2-1 inch of water. Use the lower water volume on coarser textured soils, the higher volume on finer textured soils. More than 1 inch of water may reduce weed control by moving herbicide below the effective zone in the soil. Inject dilute slurry into system through a positive displacement pump.

**PRECAUTIONS:** (1) Apply only through irrigation systems containing anti-siphon and check valves to prevent contamination of well during shutdown and overflow of solution tank. (2) Inject ahead of any right angle turn in the main line to insure adequate mixing. (3) Chemical injection pumps and water pumps must have interlocking controls to insure simultaneous shutoff. (4) Application when drift may occur from windy conditions, when system joints and connections are leaking, or when nozzles are not providing uniform distribution may cause crop injury. (5) Where sprinkler distribution patterns do not overlap sufficiently, unacceptable weed control may result. Where sprinkler distribution patterns overlap excessively, crop injury may result.

ATRAZINE 80W APPLIED ALONE  
CORN OR GRAIN SORGHUM

PREPLANT INCORPORATED OR PREEMERGENCE

BEST AVAILABLE COPY

## WEEDS CONTROLLED

barnyardgrass (watergrass)	cocklebur	mustards
giant foxtail	groundcherry	nightshade
green foxtail	jimsonweed	pigweed
large (hairy) crabgrass	kochia	purslane
wild oats	lambsquarters	ragweed
witchgrass (Fanicum capillare)	annual	velvetleaf (buttonweed)
yellow foxtail	morningglory	

## POSTEMERGENCE WITH EMULSIFIABLE OIL OR OIL CONCENTRATE IN WATER

## WEEDS CONTROLLED

annual morningglory	pigweed
cocklebur	ragweed
jimsonweed	smartweed
lambsquarters	wild buckwheat
mustards	velvetleaf

## CORN

PREPLANT (BROADLEAF AND GRASS CONTROL): Broadcast in spring after plowing at rate in Table 1. Apply before, during, or after final seedbed preparation. If soil is tilled or worked after application, avoid deep incorporation. For best results, apply within two weeks prior to planting.

PREEMERGENCE (BROADLEAF AND GRASS CONTROL): Apply during or shortly after planting before weed emergence, at rate in Table 1.

POSTEMERGENCE (BROADLEAF AND GRASS CONTROL): Apply before weeds exceed 1.5 inches in height, at rate in Table 1.

TABLE 1: BROADLEAF AND GRASS WEED CONTROL ON CORN\*

SOIL TEXTURE	BROADCAST RATE/A
COARSE Sand, loamy sand, sandy loam	2.5 lb.
MEDIUM Loam, silt loam, silt, clay loam, sandy clay loam, silty clay loam, sandy clay, or silty clay with low organic matter	1 lb.
FINE Loam, silt loam, silt, clay loam,	3.75 lb.

8 OF 23

BEST AVAILABLE COPY

sandy clay loam, silty clay loam,  
sandy clay, or silty clay with medium  
to high organic matter, and clay  
(including the dark prairie soils  
of the Corn Belt)

Peat, muck, and high organic clay  
(apply postemergence only)

3.75 lb.

\*Broadleaf control (eastern CO, western KS, western NE, NM, OK Pan Handle, west TX, and eastern WY): On sand, loamy sand, sandy loam, mild to strongly alkaline soil, and all recently leveled soil, apply no more than 1.5 lb./A, either preplant or preemergence. On other soils in these areas, apply rate in Table 1 for broadleaf and grass control

LAY-BY TREATMENT (BROADLEAF AND GRASS CONTROL): Broadcast 1.25 - 2.5 lb/A before weeds are 1.5 inches tall and corn is 20-30 inches tall. When using nitrogen solutions, direct the spray to lower 3-4 inches of cornstalks to avoid corn foliage injury.

POST EMERGENCE WITH EMULSIFIABLE OIL OR OIL CONCENTRATE IN WATER: Add 1 gallon of emulsifiable oil/A for ground application or 0.5 gal/A for aerial application, or add 1 qt of oil concentrate/A for ground application.

BROADLEAF AND GRASS CONTROL: Broadcast 2.5 lb/A plus emulsifiable oil or oil concentrate after weed emergence, but before weeds reach 1.5 inches in height.

BROADLEAF CONTROL: Broadcast 1.5 lb/A plus emulsifiable oil or oil concentrate before pigweed and lambsquarters reach 6 inches in height and before all other weeds reach 4 inches in height. A cultivation may be necessary if all weeds are not controlled or if weeds regrow.

PRECAUTIONS FOR APPLICATIONS WITH EMULSIFIABLE OIL OR OIL CONCENTRATE IN WATER: (1) Do not apply when crop is under stress from prolonged cold, wet weather, poor fertility, or other factors, or when crop is wet and succulent from recent rainfall, or crop injury may occur. (2) Inbred lines or any breeding stock may be severely injured by applications with emulsifiable oil or oil concentrate. (3) Adding other insecticides, herbicides, liquid fertilizers, or other materials is not recommended, because they may cause compatibility problems or crop injury. (4) Store and handle emulsifiable oil and oil concentrate carefully. Oil contaminated with even a small amount of water may not emulsify properly when added to the tank. (5) Do not make more than one application per season except as recommended for control of yellow nutsedge, quackgrass, and Canada thistle.

#### PROBLEM WEEDS:

Yellow nutsedge and Canada thistle: This product will control yellow nutsedge (*Cyperus esculentus*) and Canada thistle (*Cirsium arvense*) when applied according to directions. For best results, apply each year



**BEST AVAILABLE COPY**

until yellow nutsedge or Canada thistle is eliminated or reaches a level of infestation where neither weed is an economic problem. If yellow nutsedge or Canada thistle regrow following last application, cultivate once.

Four alternative methods of use for control of yellow nutsedge and Canada thistle are listed below in order of preference. If other weed species, such as annual grasses, are also expected, use alternative 2 or 4.

1. Broadcast 2.5 lb/A plus 1 gal of emulsifiable oil or 1 qt of oil concentrate. Apply after crop and yellow nutsedge or Canada thistle emerge, but before yellow nutsedge reaches a height of 3 inches or Canada thistle reaches a height of 6 inches. Repeat application before lay-by (20-30 inches), 10-20 days after the first application.
2. Broadcast 2.5 lb/A preplant for control of yellow nutsedge, or broadcast 2.5 lb./A during or shortly after planting, but prior to crop or weed emergence for control of Canada thistle. Follow with an application of 2.5 lb/A plus 1 gal of emulsifiable oil or 1 qt of oil concentrate after corn and weeds emerge, but before yellow nutsedge or Canada thistle reaches a height of 6 inches.
3. Broadcast 5 lb/A plus 1 gal of emulsifiable oil or 1 qt of oil concentrate after crop emerges, but prior to lay-by (20-30 inches). Apply after yellow nutsedge and Canada thistle emerge, but before yellow nutsedge is 3 inches tall or Canada thistle is 6 inches tall.
4. Broadcast 5 lb/A before, during, or shortly after planting but before crop or weed emergence (yellow nutsedge control only).

NOTE: Do not use emulsifiable oil or oil concentrate when corn is wet or under stress, especially when using 5 lb/A of this product. See "Precautions for applications with emulsifiable oil or oil concentrate in water" for additional directions.

**QUACKGRASS CONTROL ON LAND GOING INTO CORN PRODUCTION:**  
Split application: Broadcast 2.5 lb/A in fall or spring. Plow 1-3 weeks later. Apply a second application at 2.5 lb/A in spring before, during, or after planting, but before weeds are 1.5 inches high. This split application will control quackgrass and most annual broadleaf and grass weeds.

**SINGLE APPLICATION:** Broadcast 3.75 - 5 lb/A in fall or spring. Plow 1-3 weeks later.

**TANK MIXTURES FOR CORN**

This product may be tank mixed with these herbicides for control of certain broadleaf and grass weeds in corn:

- |                              |                               |
|------------------------------|-------------------------------|
| Dual* (metolachlor)          | Lasso* or Lasso EC (alachlor) |
| Dual + paraquat              | Lasso or Lasso EC + Roundup   |
| Dual + Roundup* (glyphosate) | Lasso or Lasso EC + paraquat  |

**BEST AVAILABLE COPY**

Dual + Princep\*  
Dual + Princep + paraquat  
Dual + Princep + Roundup  
Bexton\* or Ramrod\* (propachlor)

Paraquat  
Princep  
Princep + paraquat  
Princep + Roundup  
Sutan\*

Use tank mix directions appearing on the labels of the above herbicides when tank mixing with this product. Observe all precautions and limitations on labeling of products used in a particular tank mix.

Princep 80W, Princep 4L (4LC), or Princep Caliber\* 90  
In addition to the weeds listed under Atrazine 80W Applied Alone - Corn and Grain Sorghum - Preplant or Freemergence, this combination also controls crabgrass, fall panicum, and carpetweed.

Broadcast tank mix before planting, at planting, or after planting, but before crop and weeds emerge, at rates in Table 2. Use the 1:1 ratio for control of most weeds. Use the 1:2 ratio for expected heavy infestations of crabgrass and fall panicum. Cultivate shallowly if weeds develop.

PREPLANT: Apply in spring during or after final seedbed preparation. If soil is tilled or worked after application, avoid deep incorporation. For best results, apply within two weeks prior to planting.

PREEMERGENCE: Apply during or shortly after planting, but before crop and weeds emerge.

Refer to Corn sections of this label and to Princep 80W, Princep Caliber 90, or Princep 4L (4LC) labels for further directions, limitations, and precautions.

TABLE 2: TANK MIXTURES WITH PRINCEP ON CORN

Soil Texture	Broadcast rate/A			
	1:1 Ratio*		1:2 Ratio**	
	This product	Princep 80W+	This product	Princep 80W+
Sand, loamy sand, sandy loam	1.25 lb.	1.25 lb.	0.83 lb.	1.67 lb.
Loam, silt loam, silt, clay loam, sandy clay loam, silty clay loam, sandy clay, or silty clay with low organic matter	1.5 lb.	1.5 lb.	1 lb.	2 lb.
Loam, silt loam, silt, clay loam, sandy clay				

<b>BEST AVAILABLE COPY</b>				
----------------------------	--	--	--	--

loam, silty clay loam, sandy clay, or silty clay with medium to high organic matter, and clay (including dark prairie soils of the Corn Belt)	1.8 lb.	1.8 lb.	1.2 lb.	2.4 lb.
--	---------	---------	---------	---------

\*For control of most weeds.

\*\*For control of expected heavy infestations of crabgrass and fall panicum.

+When using Princep Caliber 90 or Princep 4L (4LC), use equivalent rates: one lb. of Princep 80W equals 0.9 lb. of Princep Caliber 90 or 1.6 pt. of Princep 4L (4LC).

PRINCEP 80W, PRINCEP 4L (4LC), OR PRINCEP CALIBER 90 PLUS ROUNDUP: Use as tank mixture for preemergence and postemergence control of certain broadleaf and grass weeds where corn will be planted directly into a cover crop, established sod, or in previous crop residues. Refer to Roundup label for all directions, weeds controlled, precautions, and limitations.

PRINCEP 80W, PRINCEP 4L (4LC), OR PRINCEP CALIBER 90 PLUS PARAQUAT: Use as tank mixture with Princep and paraquat to kill existing vegetation and for residual weed control where corn will be planted directly into a cover crop, established sod, or in previous crop residues. Add this product and Princep to water in spray tank, agitating until thoroughly mixed. Then add paraquat and a nonionic surfactant, such as X-77\*. Continue agitation during application. Broadcast 1.25 - 2.5 lb. of this product plus 1.25 - 2.5 lb. Princep 80W (or 2-4 pt Princep 4L (4LC), or 1.1 - 2.2 lb. Princep Caliber 90) plus 1-2 pt paraquat in 20-60 gal of water per sprayed acre. Apply before, during, or after planting but before corn emerges. Add 0.5 pt. of a nonionic surfactant, such as X-77 per 100 gal of spray mixture. Use the 2 pt rate of paraquat if existing vegetation is 4-6 inches tall. This mixture will not control weeds taller than 6 inches.

Refer to further limitations and precautions on labels for this product, Princep, and paraquat.

PRECAUTIONS FOR ALL APPLICATIONS TO CORN: (1) Do not apply more than 5 lb/A of this product per year. (2) Following harvest, plow (moldboard or disk-plow) and thoroughly till soil in fall or spring to minimize possible injury to spring-seeded rotational crops, regardless of rate used.

NOTE for all applications to corn: Do not graze or feed forage from treated areas for 21 days following application.

ROTATIONAL CROPS: (1) Do not rotate to any crop except corn or sorghum until the following year, or injury may occur. (2) If applied after June 10, do not rotate with crops other than corn or sorghum the next year, or crop injury may occur. (3) If the broadcast rate, or

**BEST AVAILABLE COPY**

equivalent band application rate, exceeds 3.75 lb/A of Atrazine 80W or 3.75 lb/A of Atrazine 80W plus Princep 80W (or equivalent of another Princep formulation), a crop of untreated corn or sorghum should precede the next rotational crop. (4) In the High Plains and Intermountain areas of the West where rainfall is sparse and erratic or where irrigation is required, use only when corn or sorghum is to follow corn or sorghum or when a crop of untreated corn or sorghum is to precede other rotational crops. (5) In eastern parts of the Dakotas, KS, western MN, and NE, do not rotate to soybeans if the rate applied to corn or sorghum was more than 2.5 lb/A or equivalent band application rate, or soybean injury may occur. (6) Injury may occur to soybeans planted the year following application on soils having a calcareous surface layer. (7) Do not plant sugar beets, tobacco, vegetables (including dry beans), spring-seeded small grains, or small-seeded legumes and grasses the year following application, or injury may occur.

**SORGHUM AND SORGHUM-SUDAN HYBRIDS (GRAIN AND FORAGE TYPES)**

**PREPLANT (BROADLEAF AND GRASS CONTROL):** Broadcast in spring after plowing at rate in Table 3. Apply before, during, or after final seedbed preparation. If soil is tilled or worked after application, avoid deep incorporation. For best results, apply within two weeks prior to planting.

**PREEMERGENCE (BROADLEAF AND GRASS CONTROL):** Apply during or shortly after planting, but prior to weed or crop emergence at rate in Table 3.

**TABLE 3: PREPLANT AND PREEMERGENCE BROADLEAF AND GRASS WEED CONTROL IN SORGHUM\***

SOIL TEXTURE	ORGANIC MATTER	BROADCAST RATE/A
COARSE Sand, loamy sand, sandy loam	any level	DO NOT USE (except for preemer- gence use on bedded sorghum in AZ & CA
MEDIUM AND FINE Loam, silt loam, silt	less than 1%	
clay loam, sandy clay loam	1-1.5%	2-2.5 lb.
silty clay loam, sandy clay, silty clay, clay	more than 1.5%	2.5-3 lb.

\*Do not apply preplant in AL, AR, FL, GA, LA, MS, NC, NM, OK, SC, TN or TX. Do not apply preemergence in NM, OK or TX except in northeast OK and the TX Gulf Coast.

In case of planting failure sorghum or corn may be replanted. Do not make a second broadcast application, or injury may occur. If originally applied in a band and sorghum or corn is replanted in untreated row middles, this product may be applied in a band to the

second planting.

**BEST AVAILABLE COPY**

**PREEMERGENCE BROADLEAF WEED CONTROL IN FURROW IRRIGATED BEDDED SORGHUM (AZ AND CA ONLY):** For preemergence control of many broadleaf weeds, broadcast 1-1.5 lb/A after bed preparation, during or after planting, but before sorghum and weeds emerge and before the first furrow irrigation. Follow with several regular irrigations, making sure to thoroughly wet all soil.

**PRECAUTIONS FOR PREEMERGENCE APPLICATION TO FURROW IRRIGATED BEDDED SORGHUM IN AZ AND CA:** To avoid possible sorghum injury, do not use on sand or loamy sand soil or on sorghum planted in furrows. Applications to sorghum growing on alkali soils or where cuts, fills, or erosion have exposed calcareous or alkali subsoils may cause crop injury. In case of crop failure, do not replant sorghum for 8 months following application. Corn may be planted immediately.

**POSTEMERGENCE (BROADLEAF AND GRASS CONTROL):** Apply at rate in Table 4 before weeds exceed 1.5 inches in height. Apply up to "close-in".

**TABLE 4: POSTEMERGENCE BROADLEAF AND GRASS WEED CONTROL IN SORGHUM**

SOIL TEXTURE	MINIMUM SORGHUM HEIGHT AT TREATMENT	BROADCAST RATE/A
Sandy or loamy sand	DO NOT USE	
Sandy loam	See directions for postemergence broadleaf weed control	
Silt loam to sandy clay loam	Completely emerged	2.5-3 lb.
Olton and Pullman clay soil	6 inches	2.5-3 lb.
Silty clay loam and finer textured soil	Completely emerged	3.75 lb.

**POSTEMERGENCE BROADLEAF WEED CONTROL WITH EMULSIFIABLE OIL OR OIL CONCENTRATE IN WATER:** Broadcast 1.5 lb/A for control of many broadleaf weeds. Apply before pigweed and lambsquarters reach 6 inches in height and before all other weeds reach 4 inches in height. In CO, western KS, NM, OK, TX and desert regions of AZ and CA, apply when sorghum is 6-10 inches in height, but before it reaches boot stage. In all other areas, apply after sorghum reaches the 3-leaf stage. Add 1 gal of emulsifiable oil/A for ground application and 0.5 gal/A for aerial application or add 1 qt/A of oil concentrate for ground application. A cultivation may be necessary if all weeds are not controlled or if weeds regrow.

For the list of weeds controlled, see Atrazine 80W Applied Alone - Corn or Grain Sorghum - Postemergence with Emulsifiable Oil or Oil Concen-

**BEST AVAILABLE COPY**

trate in Water.

---

**PRECAUTIONS FOR APPLICATIONS WITH EMULSIFIABLE OIL OR OIL CONCENTRATE IN WATER:** See "PRECAUTIONS for applications with emulsifiable oil or oil concentrate in water" in Corn section.

---

**POSTEMERGENCE BROADLEAF WEED CONTROL WITH SURFACTANT** (CO. western KS, NM, OK, TX, and desert regions of AZ and CA only): Broadcast 1.5 lb/A plus 0.75-1.5 pt of surfactant after sorghum reaches 6 inches in height, but before weeds exceed 1.5 inches in height. Apply only on sandy loam and finer textured soil.

---

**PRECAUTIONS FOR ALL APPLICATIONS TO SORGHUM:** (1) Heavy rain immediately following application tends to cause excessive concentrations of herbicide in seed furrow, resulting in possible crop injury. Do not apply to furrow-planted sorghum until furrows are leveled (plowed in). Level deep planter marks or seed furrows before application. (2) Application to sorghum growing under stress caused by minor element deficiency or to sorghum growing on highly calcareous soil may result in crop injury. (3) Following harvest, plow (mold-board or disk-plow) and thoroughly till soil in fall or spring to minimize possible injury to spring-seeded rotational crops, regardless of rate used. (4) Injury may occur if both this herbicide, preplant or preemergence, and an at-planting systemic insecticide applied in-furrow are used.

---

**NOTE:** Do not graze or feed forage from treated areas for 21 days following application.

**ROTATIONAL CROPS:** See "Rotational crops" in Corn section.

**TANK MIXTURES FOR GRAIN SORGHUM**

**IGRAN\* 80W:** Use as tank mixture for control of certain broadleaf and grass weeds in grain sorghum. Refer to Igran 80W label for all directions, weeds controlled, precautions, and limitations.

**DUAL 8E:** Use as tank mixture with Dual 8E for control of those weeds listed on the Dual label, as well as on this label. Use this tank mixture only on-Concep\* treated sorghum seed. Refer to the Dual 8E label for all directions, precautions, and limitations.

**ATRAZINE 80W ALONE - CHEMICAL FALLOW**

**WHEAT-SORGHUM-FALLOW:** To control annual broadleaf and grass weeds following wheat harvest and in the following sorghum crop when grown under minimum tillage, broadcast 3.75 lb/A to wheat stubble immediately following wheat harvest. If weeds are present at application, remove them with a sweep plow or other suitable implement after application, or use an approved contact herbicide before or after the application of

BEST AVAILABLE COPY

Atrazine 80W. Plant sorghum into wheat stubble the following spring with minimum soil disturbance. Use a surface planter or a planter leaving a shallow furrow. If weeds are present at planting, remove them with a sweep plow or other suitable implement before planting.

For the list of weeds controlled, see Atrazine 80W Applied Alone - Corn or Grain Sorghum - Preplant Incorporated or Preemergence.

---

**PRECAUTIONS:** (1) Use only on split loam or finer textured soil. (2) Wheat-sorghum-fallow cropping sequence must be followed. (3) Do not apply following sorghum harvest.

---

**NOTE:** Do not graze or feed forage from treated area to livestock. Do not plant any crop other than those on this label within 18 months following treatment.

**WHEAT-CORN-FALLOW (KS, NE):** This product controls cheatgrass (downy brome, chess), kochia, mustards, pigweed, Russian thistle, wild lettuce, wild sunflower, and volunteer wheat during period after wheat harvest. Weed control may extend into following corn crop grown under minimum tillage.

Follow directions for use, notes, and precautions in the 'Wheat-Sorghum-Fallow' section above, substituting corn for references to sorghum.

**WHEAT-FALLOW-WHEAT (CO, KS, MT, NE, ND, SD, AND WY):** For preemergence control of cheatgrass (downy brome, chess), common lambsquarters, field pennycress, kochia, mustard, Russian thistle, wild lettuce, and suppression of volunteer wheat during fallow period of a wheat-fallow-wheat rotation, broadcast 0.6-1.25 lb/A on all soils except those listed under "Precautions". For control of pigweed and wild sunflower, use the higher rate. Apply to stubble ground. Treat only once during same fallow period.

#### TANK MIXTURES FOR CHEMICAL FALLOW

##### WHEAT-SORGHUM-FALLOW OR WHEAT-CORN-FALLOW (KS, NE)

**Igran 80W:** Use as tank mixture for control of certain broadleaf and grass weeds in wheat-sorghum-fallow or wheat-corn-fallow rotations. Refer to Igran 80W label for directions, weeds controlled, precautions, and limitations.

**Paraquat:** If weeds are present at application, a tank mix with paraquat may be used. Broadcast 3.75 lb/A of Atrazine 80W plus 1-2 pt of paraquat in 20-60 gal of water/A by ground equipment. Add 0.5-1 pt of a nonionic surfactant, such as X-77, per 100 gal of spray mixture. Add Atrazine 80W to spray tank first and thoroughly mix with water. Then add paraquat, followed by surfactant. Use the 2 pt rate of paraquat if weeds are 4-6 inches tall. This mixture will not control weeds taller than 6 inches. Apply to stubble ground. Treat only once during same fallow period. Refer to paraquat label for further directions, precautions, and limitations.

**BEST AVAILABLE COPY**

WHEAT-FALLOW-WHEAT (CO, KS, MT, NE, ND, SD, AND WY)

Igran 80W: Use as tank mixture for control of certain broadleaf and grass weeds in wheat-fallow-wheat rotations. Refer to Igran 80W label for directions, weeds controlled, precautions, and limitations.

Paraquat: If weeds are present at application, a tank mix with paraquat may be used. Broadcast 0.6-1.25 lb of Atrazine 80W plus 1-2 pt of paraquat in 20-60 gal of water/a by ground equipment. Add 0.5-1 pt of a nonionic surfactant, such as X-77, per 100 gal of spray mixture. Add Atrazine 80W to spray tank first and thoroughly mix with water. Then add paraquat, followed by surfactant. Use the 2 pt rate of paraquat if weeds are 4-6 inches tall. This mixture will not control weeds taller than 6 inches. Apply to stubble ground. Treat only once during same fallow period. Refer to paraquat label for further directions, precautions, and limitations.

If weeds are present at application and this product is used alone, use either an approved contact herbicide before or after treatment, or tillage after treatment.

Use tillage to control weeds which escape during fallow period. Till before planting. For this product applied alone or in tank mixture with paraquat, plant at least 2 inches deep and 12 months or more after application.

---

PRECAUTIONS: (1) Do not use on sand soil. (2) Do not treat eroded hillsides, caliche and rocky outcroppings, or exposed calcareous subsoil. (3) Do not treat soils of the Rosebud and Canyon Series in western NE and adjoining counties in CO and WY. (4) Do not treat soils with calcareous surface layers. (5) Avoid spray overlap.

---

NOTE: Do not graze treated areas within 6 months after application.

AERIAL APPLICATION: In order to assure that spray will be controllable within the target area when used according to label directions, make applications at a maximum height of 10 ft. using low drift nozzles at a maximum pressure of 40 psi, and restrict application to periods when wind speed does not exceed 10 mph. To assure that spray will not adversely affect adjacent sensitive nontarget plants, apply Atrazine 80W alone by aircraft at a minimum upwind distance of 400 ft from sensitive plants, or apply Atrazine 80W + Igran 80W at a minimum upwind distance of 500 ft from sensitive plants. When applying Atrazine 80W + Igran 80W by aircraft, use a minimum of 5 gal of water per acre.

**PROSO MILLET**

For preplant or preemergence control of foxtails, kochia, common lambsquarters, redroot pigweed, Russian thistle, slimleaf lambsquarters and spotted spurge, apply 0.6-1.25 lb/A on soils with less than 2% organic matter, or apply 1.25-2.5 lb/A on soils with 2% or higher organic matter.



BEST AVAILABLE COPY

PRECAUTIONS: Make only one application per year.

ROTATIONAL CROPS: See "Rotational crops" in Corn section.

#### RANGELAND

To control certain weeds, including cheatgrass (downy brome, chess), common (annual) broomweed, little barley, medusahead, sagewort, and tumble mustard. Use lower rate where cheatgrass is major problem weed; use higher rate for other weeds.

**AERIAL APPLICATION:** In order to assure that spray will be controllable within the target area when used according to label directions, make applications at a maximum height of 10 ft above vegetation, using low drift nozzles at a maximum pressure of 40 psi, and restrict application to periods when wind speed does not exceed 10 mph. To assure that spray will not adversely affect adjacent sensitive nontarget plants, apply Atrazine 80W by aircraft at a minimum upwind distance of 400 ft from sensitive plants.

**ESTABLISHMENT OF PERENNIAL RANGE GRASSES IN WEST (CA, ID, NV, OR, UT, WA):** Broadcast 1-1.25 lb/A in fall after rains begin but before ground freezes. Perennial range grass may be seeded when rain begins the next fall following treatment. Plant in deep furrows such as those made by a deep furrow rangeland drill. Treated areas may be grazed 7 months after seeding. However, to provide sufficient time for establishment of desirable range grasses, treated areas should not be grazed until the fall following seeding.

#### RENOVATION OF EXISTING STANDS OF PERENNIAL RANGE GRASSES

Central Great Plains (CO, KS, MT, NE, ND, SD, WY): Broadcast 0.6-1.25 lb/A in fall before ground freezes.

Southern Great Plains (OK, TX): Broadcast 1.25 lb/A in spring before April 30.

West (CA, ID, NV, OR, UT, WA): Do not graze area to be treated for 6 months prior to application. Broadcast 0.6-0.9 lb/A by ground equipment in fall before ground freezes.

Bluegrass and intermediate wheatgrass may be injured by this product at recommended rates.

**NOTE:** Do not cut or feed range grass hay. Do not graze treated areas within 7 months following fall application or 3 months following spring application. Apply only once per year.

#### WARM SEASON PERENNIAL RANGE GRASSES - MIDWEST

**SWITCHGRASS AND BIG BLUESTEM:** For control of those weeds listed under Atrazine 80W Applied Alone - Corn or Grain Sorghum - Preplant Incorporated or Preemergence.

**BEST AVAILABLE COPY**

Use only on loam, silt loam, silty clay loam, clay loam, and silty clay soils with at least 1 percent organic matter.

**NEW SEEDING:** Broadcast 2.5 lb/A after planting but before weeds emerge. Prepare a good, firm seedbed. Plant 1/2 inch deep with a grassland drill (preferred method) or a conventional drill. If a conventional drill is used on prepared seedbeds, remove all tension from the disk openers. For best results, cultipack or roll after planting. Clip weeds that escape in July or August. Avoid clipping switchgrass or big bluestem seedings.

**ESTABLISHED STANDS.** Broadcast 2.5 lb/A in April or early May, before weed emergence.

**NOTE:** Do not cut for hay. Do not graze treated areas within 4 months following application at seeding or 3 months following application to established switchgrass or big bluestem. Treat only once per year.

**ROTATIONAL CROPS:** See "Rotational crops" in Corn section.

**SUGARCANE**

For control of many broadleaf and grass weeds, including amaranths, crabgrass, fireweed, Flora's paintbrush, foxtails, junglerice and wiregrass, broadcast 2.5-5 lb of Atrazine 80W at time of planting or ratooning, but before sugarcane emerges. Broadcast aerially in a minimum of 5 gal of spray per acre, or broadcast or band by ground equipment in a minimum of 20 gal per acre, unless indicated otherwise. One additional application may be made over the sugarcane as it emerges, and two additional applications may be made interline after emergence as directed sprays. Repeat treatments, where needed, may be applied broadcast, band, or interline as suggested with the final application being prior to "close-in". Do not exceed the rate of herbicide suggested for any one crop of sugarcane.

**NOTE:** Where high rates of Atrazine 80W are used, apply in a minimum of 1 gal of water for each 1 lb of product applied per acre.

**AERIAL APPLICATION:** In order to assure that spray will be controllable within the target area when used according to label directions, make applications at a maximum height of 10 ft. using low drift nozzles at a maximum pressure of 40 psi, and restrict application to periods when wind speed does not exceed 10 mph. To assure that spray will not adversely affect adjacent sensitive nontarget plants, apply Atrazine 80W alone by aircraft at a minimum upwind distance of 400 ft from sensitive plants.

**FLORIDA**

For control of emerged pellitoryo weed: Apply 0.5-0.75 lb/A in at least 40 gal of water as a directed spray by ground equipment prior to "close-in". Add 4 qt of surfactant for each 100 gal of spray. Thoroughly cover weed foliage.

For control of alexandergrass, large crabgrass, pellitory (artillery)

BEST AVAILABLE COPY

weed, and spiny amaranth, use one of the following methods at planting or ratooning:

- 1. Apply 5 lb/A preemergence. Follow with one or two applications as needed, postemergence to sugarcane and weeds, at 2.5 lb/A. Treat before weeds exceed 1.5 inches in height.
- 2. Apply 1-3 times, as needed, at 2.5 lb/A postemergence to sugarcane and weeds. Treat before weeds exceed 1.5 inches in height.

TEXAS

Use Atrazine 80W for control of barnyardgrass, pigweed, purslane and sunflower, in plant or ratoon sugarcane.

Apply 5 lb/A of Atrazine 80W preemergence. Follow with one or two applications, as needed, at 3.75 lb/A postemergence to sugarcane and weeds.

For best results when weeds are emerged, add a nonionic surfactant at a concentration of 2 qt/100 gal to the spray and apply before weeds exceed 1.5 inches in height.

PRECAUTIONS: (1) injury to sugarcane may occur when under moisture stress, when soil is of low adsorptive capacity, or when land is first cropped to sugarcane. (2) Do not apply after "close-in". (3) Do not apply more than 12.5 lb/A to any one crop of sugarcane.

TURF GRASSES FOR SOD

ST. AUGUSTINEGRASS, CENTIPEDEGRASS, AND ZOYSIA GRASS

Broadcast 2.5-5 lb/A according to soil texture to control those weeds listed under Atrazine 80W Applied Alone - Corn or Grain Sorghum - Pre-plant Incorporated or Preemergence.

Muck or peat	5 lb.	Old beds: Within 2 days after lifting sod
		New beds: 3-4 days after sprigging or plugging
Sandy soil	2.5 lb.	Old beds: Within 2 days after lifting sod.
		New beds: 7-10 days after sprigging or plugging.

If weeds regrow, apply an additional 2.5 lb/A on muck or peat, or 1.25 lb/A on sandy soil.

PRECAUTIONS: (1) Do not apply within 30 days prior to cutting or

BEST AVAILABLE COPY

lifting. (2) Do not apply in combination with surfactants or other spray additives, or injury may occur. (3) Do not use north of NC.

---

#### MACADAMIA NUTS

For preemergence control of many broadleaf and grass weeds, including crabgrass, foxtail, wiregrass. Flora's paintbrush, spanishneedles, and fireweed, broadcast 2.5-5 lb/A before harvest and before weeds emerge. Repeat as necessary. Do not spray when nuts are on ground during harvest period. Do not apply by air.

#### PINEAPPLE

For control of purslane, spanishneedles, annual grasses, annual bindweed, ageratum, amaranth, rattlepod. Flora's paintbrush, fireweed, spurge, indigo, and papalo, apply up to 8 lb/A as a blanket spray immediately after planting or following harvest. Make additional blanket or interspace applications of up to 2 lb/A at 1-2 month intervals as needed, prior to differentiation. Apply in 20-40 gal of water/A to assure thorough coverage.

---

PRECAUTIONS: (1) Do not apply more than 37.5 lb/A per cycle. (2) Repeated monthly applications to plant foliage may slow plant growth and delay fruiting.

---

NOTE: Do not apply within 45 days of fruit harvest, or forage harvest if forage is to be fed to livestock.

#### GUAVA

Use only on established plantings which are at least 18 months old. Apply as a directed spray at 2.5-5 lb/A of Atrazine 80W in 20-50 gal of spray mix preemergence or early postemergence to weeds. When applying postemergence the use of a surfactant and greater spray volume (80-100 gal of spray mix per acre) may enhance weed control. Do not apply more frequently than at 4 month intervals. Do not apply more than 10 lb of Atrazine 80W per year. Controls many annual broadleaf and grass weeds, including fireweed, purslane, scarlet pimpernel, spanish needles, and sowthistle.

Do not allow spray to contact foliage or fruit.

#### CONIFERS

For control of annual broadleaf and grass weeds prior to transplanting, after transplanting or in established conifers (including Douglas, fir, grand fir, noble fir, white fir, Austrian pine, bishop pine, Jeffrey pine, knobcone pine, loblolly pine, lodgepole pine (shore pine), monterey pine, ponderosa pine, Scotch pine, slash pine, blue spruce, and Sitka spruce): Broadcast 2.5-5 lb in a minimum of 5 gal of water per acre by air or 10 gal by ground before weeds are 1.5 inches tall.

**BEST AVAILABLE COPY**

Apply to established trees between fall and early spring while trees are dormant. For new transplants, apply during or soon after transplanting. For applications prior to transplanting, allow sufficient precipitation to activate Atrazine before transplanting. In areas where spring and summer rainfall is inadequate to activate Atrazine 80W, apply during fall prior to spring transplanting.

For the list of weeds controlled, see Atrazine 80W Applied Alone - Corn or Grain Sorghum - Preplant Incorporated or Preemergence.

**QUACKGRASS CONTROL:** Broadcast 5 lb in a minimum of 5 gal of water per acre by air or 10 gal by ground between fall and early spring while trees are dormant and before quackgrass is more than 1.5 inches tall.

**NOTE:** Temporary injury to trees may occur following the use of Atrazine 80W on coarse-textured soil.

---

**PRECAUTIONS:** (1) Do not graze treated areas. (2) Do not apply to seedbeds. (3) Apply only once per year.

---

**AERIAL APPLICATION:** In order to assure that spray will be controllable within the target area when used according to label directions, make applications at a maximum height of 10 ft above vegetation, using low drift nozzles at a maximum pressure of 40 psi, and restrict application to periods when wind speed does not exceed 10 mph. To assure that spray will not adversely affect adjacent sensitive nontarget plants, apply Atrazine 80W by aircraft at a minimum upwind distance of 400 ft from sensitive plants.

**NONSELECTIVE WEED CONTROL ON NONCROP LAND**

This product provides long-term weed control on industrial sites, highway medians and shoulders, railroad rights-of-way, lumberyards, petroleum tank farms, and in noncrop areas on farms, such as around buildings, equipment and fuel storage areas, along fences, and lanes. Do not use near adjacent desirable plants or in greenhouses, or injury may occur.

Apply before or soon after weeds begin growth. Make postemergence applications when weeds are young and actively growing. Use the higher rates on fine clay and muck soils. Use sufficient water to assure thorough coverage. Use at least 1 gal of water for each pound of product; use more if practical

**NOTE:** Do not allow livestock to graze treated areas.

**ANNUAL BROADLEAF AND GRASS WEEDS** (including barnyardgrass, cheatgrass, crabgrass, lambsquarters, foxtail, ragweed, puncturevine, and turkey mullein): Broadcast 5-12.5 lb/A.

**HARD-TO-KILL ANNUAL AND MANY PERENNIAL BROADLEAF AND GRASS WEEDS** (including bluegrass, burdock, Canada thistle, dogfennel, orchardgrass,

**BEST AVAILABLE COPY**

plantain, quackgrass, purpletop, redtop, and smooth brome): Broadcast 12.5-25 lb/A.

HARD-TO-KILL BIENNIAL AND PERENNIAL WEEDS (including bullthistle and sowthistle): Broadcast 25-50 lb/A.

For longer residual control in regions of high rainfall and a long growing season, broadcast 25-50 lb/A.

For small areas, 4.5 oz. per 1,000 sq ft is equivalent to 12.5 lb/A.

**CONDITIONS OF SALE**

All statements concerning the use of this product apply only when used as directed. THE MANUFACTURER MAKES NO WARRANTIES, EXPRESSED OR IMPLIED, CONCERNING THIS PRODUCT OR ITS USE, WHICH EXTEND BEYOND THE DESCRIPTION ON THE LABEL. Read all directions carefully.

Bexton\* trademark of Dow Chemical Co. for propachlor

Caliber\* trademark of Ciba-Geigy

Concep\* trademark of Ciba-Geigy

Dual\* trademark of Ciba-Geigy for metolachlor  
U.S. Patent No. 3,937,730

Igran\* trademark of Ciba-Geigy for terbutryn  
U.S. Patent No. 3,634,062

Lasso\* trademark of Monsanto Company for alachlor

Princep\* trademark of Ciba-Geigy for simazine

Ramrod\* trademark of Monsanto company for propachlor

Roundup\* trademark of Monsanto Company for glyphosate herbicide

Sutan\* + trademark of Stauffer Chemical Co. for butylate

X-77\* trademark of Kalo Laboratories, Inc.