Active Ingredients

Pyrethrins .033%
*Piperonyl Butoxide .33 %
*(butylcarbityl) (6-propylpiperonyl) ether
Related Compounds .046%
Inert Ingredients 99.591%

Active ingredients above indicates residue after 5 months storage of IRT Sacks originally treated with 50 ± 10 mg/sq. ft. piperonyl butoxide, and 5 ± 1 mg/sq. ft. pyrethrius on the outer surface of the outer ply of basis weight 60 lb. per 3,000 sq. ft. ream. Treatment increases resistance to penetration by insects for a period of 9 to 12 months under normal shipping, handling and storage conditions. Tight end closures and specially designed longitudinal seam assist in protecting the contents, providing the sack is otherwise intact. For maximum effectiveness, IRT sacks must be filled within 5 months of the date of fabrication.

Number of sacks and weight of contents______

of_____sacks.

Date of Sack Fabrication______

USDA Registration No. 9882-1

St. Regis Paper Company New York, New York 10017

The outer ply of this shipping sack has been treated with synergized pyrethrins to increase its resistance to penetration by insects under normal shipping, handling and storage conditions. Tight end closures and specially designed longitudinal seam essist in protecting the contents, previding the each is otherwise intest. This sack has been registered with the United States Department of Agriculture under USDA Reg. No.

ACCEPTED

11-29-69.

UNDER THE FEDERAL INSECTICIDE FUNGICIDE AND FUND THE ACT FOR ECO (SIAC) INDICATE HEBISTER-ED UNDER MO. 1982-1

Active	Ingr
Py	rethri
	peron
	*(b)
Re	lated
Inert I	

storage of I ft. piperony the outer su 3,000 sq. ft. tion by inse shipping, ha and speciall the contents mum effection the date

N	U	m	D	6	r	O.	•

of	
•	

Date of Sac

USDA Regi

The outer ply of this shipping treated with synamical dispression residence to proceed and in by inner shipping, harding and the control of the proceeding and the control of the procedure of th