HAZARO TO HUMANS AND DOMESTIC ANIMALS

Harmful if swalled. Avoid breathing vapor. Avoid contact with skin, eyes, or clothing. Wear mask or respirator recommended by NIOSH for protection. Weah exposed skin with soap and water after using.

On not apply while food processing is under way. Exposed edible products should be removed, covered or stored in closed containers. Facilities and equipment in federally inspected meet and poultry plants must be washed with an effective cleaning compound followed by a potable water rines efter apraying.

Remove pets and birds, and cover fish aquariums before spraying.

STORAGE AND DISPOSAL

Do not store in direct sunlight, or near heat such se an open flame. Do not use toward an open flame. Do not conteminate water, food, or feed by storage or disposal.

CONTAINER DISPOSAL Triple rines (or equivalent). Then offer for recycling or reconditioning, or punture and dispose of in a senitary landfill . or by public disposel trucks (California).

DIRECTIONS FOR USE It is a violation of Pederal Law to use this product in a manner inconsistent with its labeling.

Space spray: for control of accessible, exposed stages of flies, including fruit flies, mosquitoes, small flying moths, gnate, wasps, hornate, chases skippers, and these common stored product peath: granary wasvils, rice weevils, confused flows bestles, baw-coothed grain bestles, apider bestles, cigaratte bestles, Drugstors bestles, Angoumois grain moths, Mediteranness flour moths, Indianness methe; Tobacco moths, meal worms, and grain mites; Close windows and doors and shut off ventilating systems. Use a fogger or vaporizor adjusted to deliver an aerosol apray (no droplets over 50 microns in diameter

CUSTOM

Food Plant Fogging

INSECTICIDE

ACTIVE INGREDIENTS:

100% 0.5%

Pyrethrine 0.5%
*Piperonyl Butoxide Technical 5.0%
Petroleum Distillate 94.5%

*Equivalent to 4.0% (butylcarbityl)
(6-propylpiperonyl) ether end to 1.0%
related compounds.

CAUTION KEEP OUT OF REACH OF CHILDREN

See Side Panel
For Additional Precautionary Statements

PHYSICAL DR CHEMICAL HAZARDS
Do not apray near or toward an open flame.

A Concentrated PYREMONE PRODUCT.

Registered trademark of: FAIRFIELD AMERICAN CORPORATION.

Mfg. Ey:

Custom Blend of California

202 W. BROADWAY, SAN GABRIEL, CALIFORNIA EPA EST. NO. 9887-CA-1 91776 EPA REG. NO. 9887-14

NET CONTENTS: GALLON

and AOS less than 3U microns). Apply at rate of 1 page for 1000 cubic feet of space. Offect the spruy toward the ceiling and upper corners of the area and berind obstructions. For best control of washe and hornets, apply directly to the nests in the evening. Keep the area closed for at least 15 minute Vacate the treated area and ventilate before reoccupying. Repeat treatment as necessary.

Space and Contact Spray: For control of accessible. exposed stones of silverfish, crickets, roaches, eniders, scornions, clover mites, cheese mites, earwigs, and ents, first using equipment which cen deliver a pin soray, direct soray into all cracks and cravices in the woodworck, wells, floors, underweith sinks, behind pipes and in all places where these insects are known to seek shelter. Contact os many insects as possible. Soray directly. all ant trails. Do not allow soray to contact foot. Then, shutting all doors, windows, and shutting off all ventilating systems, proceed with space spraying directions as given above. Has a rosson rate of 1 once per 1000 cubic feet of space. Keep the arma closed for at least 15 minutes. Vecate the treated area and ventilata before reoccuping. Represt treatment sa necessery.

In mushroom production and processing: To control mushroom flies or funges gnate in growing houses apply at a rate of 1 once per 1000 cubic feet of space. Engger should be operated 8 to 10 feet away from beds and directed alightly upward over them. Circulating fans should be left in operation, but for best results, refrigeration should be turned often packing and canning plants, use at a rate of 1 once to 2000 cubic feet of space.

Designed for use in Edible Product Areas of Official Establishments Operating Under the Meut, Poultry, Shell Egg Grading and Egg Products & Inspection Programs and in Other Food Processing Plants.

Apply with Equipment Designed and Adjusted to Deliver a True Mist.

Buyer assumes all risks of use, storage or handling of this material not in strict accordance with directions given herswith.

ACCEPTED

OCT 10 1985

Un' Act, on a natural Act, on a natural repaired in PA Rey, No.