

RED TORPEDO

RAT KILLER

CAUTION

DIRECTIONS: USE IN A MANNER TO PROTECT CHILDREN, PETS AND DOMESTIC ANIMALS. Use of tamper-proof bait boxes is recommended. Select locations where rats and mice travel, feed and water. Place bait along walls, in corners and in concealed locations where the rodents can feed unobserved rather than in an open area such as the center of a room. For rats, use one or two bait placements of $\frac{1}{4}$ to one lb. each at every point of large activity. For mice, a larger number of bait placements should be made. Tablespoon amounts ($\frac{1}{4}$ to $\frac{1}{2}$ ounces) of bait should be placed at 8 - 12 ft. intervals. Inspect daily; do not allow bait to run out. Bait until all signs of feeding have stopped. Keep an uninterrupted supply of bait for rats at least 10 days, for mice at least 15 days. If bait becomes sour, moldy or otherwise spoiled, it should be promptly replaced.

CAUTION:

If swallowed by humans, pets, or domestic animals, this material may reduce the clotting ability of the blood and cause bleeding. In such case, intravenous and oral administration of Vitamin K combined with blood transfusions may be indicated, as in the case of hemorrhage caused by overdoses of bishydroxycoumarin.

RED TORPEDO

USE AS IS

RAT KILLER

CAUTION

DIRECTIONS: USE IN A MANNER TO PROTECT CHILDREN, PETS AND DOMESTIC ANIMALS. Use of tamper-proof bait boxes is recommended. Select locations where rats and mice travel, feed and water. Place bait along walls, in corners and in concealed locations where the rodents can feed unobserved rather than in an open area such as the center of a room. For rats, use one or two bait placements of $\frac{1}{4}$ to one lb. each at every point of large activity. For mice, a larger number of bait placements should be made. Tablespoon amounts ($\frac{1}{4}$ to $\frac{1}{2}$ ounces) of bait should be placed at 8 - 12 ft. intervals. Inspect daily; do not allow bait to run out. Bait until all signs of feeding have stopped. Keep an uninterrupted supply of bait for rats at least 10 days; for mice at least 15 days. If bait becomes sour, moldy or otherwise spoiled, it should be promptly replaced.

CAUTION:

If swallowed by humans, pets, or domestic animals, this material may reduce the clotting ability of the blood and cause bleeding. In such case, intravenous and oral administration of Vitamin K combined with blood transfusions may be indicated, as in the case of hemorrhage caused by overdoses of bishydroxycoumarin.