

USE DIRECTIONS

FOR CLEANING

Lightly soiled areas - surfaces to be cleaned should be mopped or brushed with a solution containing 1 ounce of Hanophene per gallon of water. Rinse cleaned area with fresh water. Heavily soiled areas - mop or brush soiled surface with a solution containing 2 ounces of Hanophene per gallon of water. Rinse cleaned surface with fresh water.

GENERAL PURPOSE DISINFECTION

To disinfect walls, floors and other hard surfaces in homes, schools, restaurants, rest rooms and other public areas, mop or brush surface to be treated with a solution containing 1 ounce of Hanophene per gallon of water. Surfaces that are heavily soiled should be cleaned with 2 ounces of Hanophene per gallon of water and then disinfected. All treated food contact surfaces and/or equipment must be thoroughly scrubbed with detergents and rinsed with potable water prior to reuse. To disinfect locker rooms, shower stalls, walkways around pools and other areas against pathogenic fungi, mop area with a solution containing 1 ounce of Hanophene in 1 gallon of water. This solution will destroy TRICHOPHYTON MENTAGROPHYTES, a causative agent of athlete's foot.

HOSPITAL DISINFECTION

To disinfect walls, floors, furniture and other hard surfaces in hospitals and nursing homes, mop or brush surface to be treated with a solution containing 1 ounce of Hanophene per gallon of water. Those surfaces that are heavily soiled with blood, fecal material or urates should be thoroughly scrubbed with a solution containing 2 ounces per gallon of water and then rinsed with a fresh solution containing 2 ounces per gallon of water before disinfection.

NET CONTENTS

Hanco

960

Hanophene

ACCEPTED

SEP 12 1974

UNDER THE FEDERAL INSECTICIDE ACT
FUNGICIDE AND RODENTICIDE ACT
FOR ECONOMIC POISON REGISTRATION
REGISTERED UNDER NO. 9451

DISINFECTANT - CLEANER - SANITIZER -
FUNGICIDE - DEODORANT

ACTIVE INGREDIENTS

Isopropyl Alcohol	12.00%
Soap	5.00%
sodium o-phenylphenate	3.00%
sodium 4-chloro-2-cyclopentylphenate	3.00%
tetrasodium ethylene diamine tetraacetate	1.00%
Essential Oils	0.50%
INERT INGREDIENTS	75.50%

Phenol Coeff:

S. aureus—15
S. typhosa—9

DANGER

SANITIZATION

To sanitize walls, floors and equipment in food processing plants, first clean thoroughly with a solution containing 1 ounce of Hanophene per gallon of water. If surface is heavily soiled, use Hanophene at a rate of 2 ounces per gallon to insure thorough cleaning. Sanitize cleaned surface with a rinse solution containing 1 ounce of Hanophene per gallon of water. All treated food contact surfaces and/or equipment must be thoroughly scrubbed with detergents and rinsed with potable water prior to reuse.

DEODORIZATION

Deodorize surfaces during cleaning, mop or brush surfaces with a solution containing 2 ounces of Hanophene per gallon of water. Hanophene is an effective antibacterial cleaner designed for use in household, industrial and institutional cleaning. When used as directed, Hanophene will disinfect hard surfaces, such as floors, walls, counter tops and furniture. It may be relied upon to deodorize bathroom areas, garbage pails, food storage areas, kennel areas and other areas where obnoxious odors may develop. This product may be used as a general purpose antibacterial-antifungal detergent in homes, schools, restaurants, hospitals, nursing homes, public rooms, food processing plants and in other places where efficient cleaning and antimicrobial action is desired.

See left panel for use directions.

DANGER

KEEP OUT OF REACH OF CHILDREN.

Corrosive. May cause eye damage. Causes skin irritation. Protect eyes and skin when handling. Do not get in eyes, on skin or on clothing. Harmful if swallowed. Avoid contamination of food.

FIRST AID

In case of contact, immediately flush eyes or skin with plenty of water for at least 15 minutes. For eyes, call a physician. Remove and wash contaminated clothing before reuse.

If swallowed, drink PROMPTLY a large quantity of milk, egg whites, gelatin solution or if these are not available, drink large quantities of water. Avoid alcohol. Call a physician immediately.

NOTE TO PHYSICIAN

Probable Mucosal damage may contraindicate the use of gastric lavage. Measures against circulator shock, respiratory depression and convulsion may be needed.

DO NOT USE, POUR, SPILL, OR STORE NEAR OPEN FLAME.

HANCO MANUFACTURING Co., Inc.
MEMPHIS, TENNESSEE