

Bananas appears and repeat at 10 day intervals as needed.
Preharvest: against Sigatoka (*Cercospora musae*).
Dosage: up to 5 lbs./acre in sufficient water to cover.
Start application as soon as disease appears and repeat at 2 weeks intervals.
Postharvest: against crown rot, surface molds (*Gleosporium*, *Thielaviopsis*, *Fusarium* species).
Dosage: up to 4 lbs./100 gallons of water as a dip or spray. Use 5 gallons of dilute suspension per 100 trays of fruit as a spray to fruit crowns after their passage through hydrocooler or water rinse spray.

Beans against downy mildew.
Dosage: 2 lbs. 100 gallons of water.
Begin application before disease appears.
Repeat applications at 5 to 7 day intervals or as needed to maintain control.
Do not apply to Succulent Beans within 4 days of harvest.

Carrots against leafspot.
Dosage: 2 lbs. 100 gallons of water.
Sprays to be started when plants are 6 weeks old and to be continued every 7-10 days as needed.

Cranberries against fruit rot.
Dosage: 2 lbs. 100 gallons of water.
Immediately after bloom 2 sprays to be effected with an interval of 14 days. Do not apply later than 4 weeks after mid-bloom.

Cucumbers against downy mildew and anthracnose.
Dosage: 2 lbs. 100 gallons of water.
Begin application as soon as disease appears. Repeat applications at 7 to 10 day intervals as necessary for control.

Endive against downy mildew.
Dosage: 2 lbs. 100 gallons of water.

Onions against downy mildew and purple blotch.
Start spraying as soon as disease appears. Treatment to be repeated weekly.
Dosage: 2 lbs. 100 gallons of water (up to 3 lbs. per acre).

Potatoes against early and late blight.
Begin application before disease appears.
Dosage: 2 lbs. 100 gallons of water.
First treatment 2 lbs. per acre.
Second and third treatment 3 lbs. per acre.
Fourth and following treatment 4 lbs. per acre.
Applications to be repeated every 10 days.

Spinach against downy mildew and white rust.
Dosage: 2 lbs. 100 gallons of water.
Start spraying as soon as disease appears and continue every 10 days, but not later than 7 days before harvest.
Remove excess residues by washing.

Sugar beets against cercospora, up to 2 lbs. per acre.
Dosage: 2 lbs. 100 gallons of water.
Start application end July early August, to be repeated every 7-10 days for 3-5 sprays, according to local recommendations.
Do not apply later than 3 weeks before harvest.
Do not feed treated tops to livestock.

Tomatoes against early and late blight, leafspot and gray mold.
Begin application before disease appears.
Dosage: 2 lbs. 100 gallons of water.
Treatment to be repeated every 8-10 days.

CAUTION: Keep Out of Reach of Children

May cause irritation of nose, throat, eyes and skin. Do not breath dust or spray. In case of contact, flush affected area with water. For eyes get medical attention.

General instructions: The recommended amount of M4 should be washed through the filter-screen into the spray-tank with the required amount of water.

M4 can be applied in a mixed spray with DDT, both Parathion and Malathion, Aldrin, Dieldrin and with most common insecticides in powder-form. It is not compatible with lead, lime or lime-sulphur.

This product is toxic to fish. Keep out of any body of water. Apply this product only as specified on this label.

Important: Keep in cool, dry place. Do not allow to become wet during storage. This will bring about a chemical change which will reduce the effectiveness of this material, as well as generate flammable vapors. Keep container closed when not in use.

KEEP AWAY FROM FIRE AND SPARKS

Notice to buyers: All goods supplied by us are of outstanding quality and we believe them to be suitable, however, since the manufacturer cannot guarantee weather conditions, crop conditions, or other influencing factors, we cannot accept any responsibility for damage arising through the application or use of this product other than that limited to the terms of this label.

N.V. CHEFARO MAATSCHAPPIJ
Rotterdam - Holland

Postbox 6014 - Telex 22658 - Tel.: (010) 256860 - Cables: Chefaro Rotterdam

Net 50 lbs.



Distributed in U.S.A. by:

Henley & Co., Inc.

202 East 44th Street

NEW YORK N.Y. - 10017

Cables: Henleysa n New York

TELEX: 22658