

MV-4 WETTABLE POWDER

MANEB WITH ZINC OXIDE ADDED

for the control of different plant diseases


ACTIVE INGREDIENT

Maneb (manganese ethylenebisdithiocarbamate) 80.- %
(expressed as manganese metal 16.5 %)

INERT INGREDIENTS

U.S.D.A. REGISTRATION No. 9932-3

20.- %



Asparagus	against <u>rust</u> . Dosage: 2 to 3 pounds of MV-4 per acre in 50 to 200 gallons of water. In established fields begin applications right after harvest and continue at 10 day intervals. For young plantings which will not be harvested, begin applications before disease appears and repeat at 10 day intervals as needed.	Start application as soon as disease appears. First sprays to be effected in the plantbeds (100 gallons of spray per acre) and further sprays in the field, every 8-10 days. Remove residues by washing. Do not apply within 7 days of harvest.
Bananas	Preharvest: against <u>Sigatoka</u> (<i>Cercospora musae</i>). Dosage: up to 5 lbs./acre in sufficient water to cover. Start application as soon as disease appears and repeat at 2 weeks intervals. Postharvest: against <u>crown rot</u> , surface molds (<i>Gleosporium</i> , <i>Thielaviopsis</i> , <i>Fusarium</i> species). Dosage: up to 4 lbs./100 gallons of water as a dip or spray. Use 5 gallons of dilute suspension per 100 trays of fruit as a spray to fruit crowns after their passage through hydrocooler or water rinse spray.	Grapes against <u>black rot</u> . Dosage: 2 lbs./100 gallons of water. Treatment to be effected before and after bloom and 8 days later.
Beans	against <u>downy mildew</u> . Dosage: 2 lbs./100 gallons of water. Begin application before disease appears. Repeat applications at 5 to 7 day intervals or as needed to maintain control. Do not apply to Succulent Beans within 4 days of harvest.	Onions against <u>downy mildew</u> and <u>purple blotch</u> . Start spraying as soon as disease appears. Treatment to be repeated weekly. Dosage: 2 lbs./100 gallons of water (up to 3 lbs. per acre).
Carrots	against <u>leafspot</u> . Dosage: 2 lbs./100 gallons of water. Sprays to be started when plants are 6 weeks old and to be continued every 7-10 days as needed.	Potatoes against <u>early and late blight</u> . Begin application before disease appears. Dosage: 2 lbs./100 gallons of water. First treatment 2 lbs. per acre. Second and third treatment 3 lbs. per acre. Fourth and following treatment 4 lbs. per acre. Applications to be repeated every 10 days.
Cranberries	against <u>fruit rot</u> . Dosage: 2 lbs./100 gallons of water. Immediately after bloom 2 sprays to be effected with an interval of 14 days. Do not apply later than 4 weeks after mid-bloom.	Spinach against <u>downy mildew</u> and <u>white rust</u> . Dosage: 2 lbs./100 gallons of water. Start spraying as soon as disease appears and continue every 10 days, but not later than 7 days before harvest. Remove excess residues by washing.
Cucumbers	against <u>downy mildew</u> and <u>anthracnose</u> . Dosage: 2 lbs./100 gallons of water. Begin application as soon as disease appears. Repeat applications at 7 to 10 day intervals as necessary for control.	Sugar beets against <u>cercospora</u> , up to 2 lbs. per acre. Dosage: 2 lbs./100 gallons of water. Start application end July/early August, to be repeated every 7-10 days for 3-5 sprays, according to local recommendations. Do not apply later than 3 weeks before harvest. Do not feed treated tops to livestock.
Endive	against <u>downy mildew</u> . Dosage: 2 lbs./100 gallons of water.	Tomatoes against <u>early and late blight</u> , <u>leafspot</u> and <u>gray mold</u> . Begin application before disease appears. Dosage: 2 lbs./100 gallons of water. Treatment to be repeated every 8-10 days.

Thielaviopsis, Fusarium species).
Dosage: up to 4 lbs./100 gallons of water as a dip or spray.
Use 5 gallons of dilute suspension per 100 trays of fruit as a spray to fruit crowns after their passage through hydrocooler or water rinse spray.

Beans

against downy mildew.
Dosage: 2 lbs./100 gallons of water.
Begin application before disease appears.
Repeat applications at 5 to 7 day intervals or as needed to maintain control.
Do not apply to Succulent Beans within 4 days of harvest.

Carrots

against leafspot.
Dosage: 2 lbs./100 gallons of water.
Sprays to be started when plants are 6 weeks old and to be continued every 7-10 days as needed.

Cranberries

against fruit rot.
Dosage: 2 lbs./100 gallons of water.
Immediately after bloom 2 sprays to be effected with an interval of 14 days. Do not apply later than 4 weeks after mid-bloom.

Cucumbers

against downy mildew and anthracnose.
Dosage: 2 lbs./100 gallons of water.
Begin application as soon as disease appears. Repeat applications at 7 to 10 day intervals as necessary for control.

Endive

against downy mildew.
Dosage: 2 lbs./100 gallons of water.

ment to be repeated weekly.
Dosage: 2 lbs./100 gallons of water (up to 3 lbs. per acre).

Potatoes

against early and late blight.
Begin application before disease appears.
Dosage: 2 lbs./100 gallons of water.
First treatment 2 lbs. per acre.
Second and third treatment 3 lbs. per acre.
Fourth and following treatment 4 lbs. per acre.
Applications to be repeated every 10 days.

Spinach

against downy mildew and white rust.
Dosage: 2 lbs./100 gallons of water.
Start spraying as soon as disease appears and continue every 10 days, but not later than 7 days before harvest.
Remove excess residues by washing.

Sugar beets

against cercospora, up to 2 lbs. per acre.
Dosage: 2 lbs./100 gallons of water.
Start application end July/early August, to be repeated every 7-10 days for 3-5 sprays, according to local recommendations.
Do not apply later than 3 weeks before harvest.
Do not feed treated tops to livestock.

Tomatoes

against early and late blight, leafspot and gray mold.
Begin application before disease appears.
Dosage: 2 lbs./100 gallons of water.
Treatment to be repeated every 8-10 days.

CAUTION: Keep Out of Reach of Children

May cause irritation of nose, throat, eyes and skin.
Do not breath dust or spray. In case of contact, flush affected area with water. For eyes get medical attention.

General instructions: The recommended amount of MV-4 should be washed through the filter-screen into the spray-tank with the required amount of water.

MV-4 can be applied in a mixed sprav with DDT, both Parathion and Malathion, Aldrin, Dieldrin and with most common insecticides in powder-form. It is not compatible with lead, lime or lime-sulphur.

Do not apply when weather conditions favor drift from areas treated. Do not apply where runoff is likely to occur. Do not contaminate water by cleaning of equipment, or disposal of wastes. Apply this product only as specified on this label.

Important: Keep in cool, dry place. Do not allow to become wet during storage. This will bring about a chemical change which will reduce the effectiveness of this material, as well as generate flammable vapors. Keep container closed when not in use.

Do not reuse empty container. Destroy it by burying with waste or burning. Stay away from smoke or fumes.

KEEP AWAY FROM FIRE AND SPARKS

Notice to buyers: All goods supplied by us are of outstanding quality and we believe them to be suitable; however, since the manufacturer cannot guarantee weather conditions, crop conditions, or other influencing factors, we cannot accept any responsibility for damage arising through the application or use of this product other than that limited to the terms of this label.

N.V. CHEFARO MAATSCHAPPIJ

Rotterdam - Holland

Postbox 6014 - Telex 22658 - Tel.: (010) 256860 - Cables: Chefaro Rotterdam

Net 50 lbs.



Distributed in U.S.A. by:

Henley & Co., Inc.

202 East 44th Street

NEW YORK N.Y. - 10017

Cables: Henleysale New York

Tel: YUkon 6-5544