




# FERTILIZER

Contains 0.3% DI-SYSTON

USDA Registration No. 9936-1



### STATEMENT OF INGREDIENTS FOR INSECT CONTROL

Di-syston 0.3%  
Inert ingredients 99.7%

### STATEMENT OF INGREDIENTS AS PLANT FOOD

Di-syston 0.3%  
Inert ingredients 99.7%

### DIRECTIONS FOR USE

Apply to soil at planting time. For best results, apply to soil at planting time. For best results, apply to soil at planting time.

### VEGETABLES

**BEANS**  
For Snap or Green Lima Beans. Dry Beans.

### IMPORTANT RECOMMENDED DI-SYSTON APPLICATIONS

**BRUSSELS SPROUTS, CAULIFLOWER**

**BROCCOLI, CABBAGE**

**POTATOES**

**SPINACH**

**PEAS** Apply to soil at planting time. For best results, apply to soil at planting time. For best results, apply to soil at planting time.

**SUGAR BEETS** Apply to soil at planting time. For best results, apply to soil at planting time. For best results, apply to soil at planting time. **IMPORTANT** See "RECOMMENDED DI-SYSTON APPLICATIONS" for more information.

### NON-BEARING FRUIT

**STRAWBERRIES PROPAGATING PLANTS ONLY**

### RECOMMENDED DI-SYSTON APPLICATIONS

**IMPORTANT** Apply to soil at planting time. For best results, apply to soil at planting time. For best results, apply to soil at planting time.

### CONTAINER DISPOSAL

**DO NOT RE-USE THE CONTAINER**



**DANGER POISON**

Keep out of reach of children



### SYMPTOMS OF POISONING

### TREATMENT

See a doctor immediately. If you have been poisoned, call your doctor or the nearest hospital.

and Thrips, apply at the rate of 3.35-6.70 lbs. of 1 to 2 lbs. actual insecticide per acre. **For Snap or Green Lima Beans**, apply specified dosage per acre in a band on each side of the seed furrow at planting time only. **For Dry Beans**, apply specified dosage per acre as a side-dressing on each side of the row after planting. Do not apply more than the specified amount. **IMPORTANT** statement under "RECOMMENDED DI-SYSTON APPLICATIONS" for additional information.

**BRUSSELS SPROUTS, CAULIFLOWER**

Apply Di-Syston at the rate of 3.35-6.70 lbs. of 1 to 2 lbs. actual insecticide per acre in a band on each side of the seed furrow at planting time only. Do not apply more than the specified amount. **IMPORTANT** statement under "RECOMMENDED DI-SYSTON APPLICATIONS" for additional information.

**BROCCOLI, CABBAGE**

Apply Di-Syston at the rate of 3.35-6.70 lbs. of 1 to 2 lbs. actual insecticide per acre in a band on each side of the seed furrow at planting time only. Do not apply more than the specified amount. **IMPORTANT** statement under "RECOMMENDED DI-SYSTON APPLICATIONS" for additional information.

**POTATOES**

Apply Di-Syston at the rate of 3.35-6.70 lbs. of 1 to 2 lbs. actual insecticide per acre in a band on each side of the seed furrow at planting time only. Do not apply more than the specified amount. **IMPORTANT** statement under "RECOMMENDED DI-SYSTON APPLICATIONS" for additional information.

**SPINACH**

Apply Di-Syston at the rate of 3.35-6.70 lbs. of 1 to 2 lbs. actual insecticide per acre in a band on each side of the seed furrow at planting time only. Do not apply more than the specified amount. **IMPORTANT** statement under "RECOMMENDED DI-SYSTON APPLICATIONS" for additional information.

**TOMATOES**

Apply Di-Syston at the rate of 3.35-6.70 lbs. of 1 to 2 lbs. actual insecticide per acre in a band on each side of the seed furrow at planting time only. Do not apply more than the specified amount. **IMPORTANT** statement under "RECOMMENDED DI-SYSTON APPLICATIONS" for additional information.

**IMPORTANT** statement under "RECOMMENDED DI-SYSTON APPLICATIONS" for additional information.

**VEGETABLES GROWN FOR SEED ONLY**

Apply Di-Syston at the rate of 3.35-6.70 lbs. of 1 to 2 lbs. actual insecticide per acre in a band on each side of the seed furrow at planting time only. Do not apply more than the specified amount. **IMPORTANT** statement under "RECOMMENDED DI-SYSTON APPLICATIONS" for additional information.

**FIELD CROPS**

**ALFALFA**

Apply Di-Syston at the rate of 3.35-6.70 lbs. of 1 to 2 lbs. actual insecticide per acre in a band on each side of the seed furrow at planting time only. Do not apply more than the specified amount. **IMPORTANT** statement under "RECOMMENDED DI-SYSTON APPLICATIONS" for additional information.

**BARLEY, OATS**

Apply Di-Syston at the rate of 3.35-6.70 lbs. of 1 to 2 lbs. actual insecticide per acre in a band on each side of the seed furrow at planting time only. Do not apply more than the specified amount. **IMPORTANT** statement under "RECOMMENDED DI-SYSTON APPLICATIONS" for additional information.

**CORN**

Apply Di-Syston at the rate of 3.35-6.70 lbs. of 1 to 2 lbs. actual insecticide per acre in a band on each side of the seed furrow at planting time only. Do not apply more than the specified amount. **IMPORTANT** statement under "RECOMMENDED DI-SYSTON APPLICATIONS" for additional information.

cool and wet or extremely dry weather, the insecticide may cause some delay in emergence, stunting of seedlings, or reduction of stand. Such damage may be more pronounced in light soil. Plant injury may also occur or may be enhanced when certain pre-emergent fertilizers are used in conjunction with DI-SYSTON.

**CONTAINER DISPOSAL**

**DO NOT RE-USE THE CONTAINER** for any other purpose. Do not use the container to hold water, milk, or other liquid. Do not use the container to hold other agricultural chemicals. Do not use the container to hold anything other than the product for which it was designed. Do not use the container to hold anything other than the product for which it was designed.



**DANGER POISON**

Keep out of reach of children



Keep out of reach of children. Do not use for any other purpose. Do not use the container to hold water, milk, or other liquid. Do not use the container to hold other agricultural chemicals. Do not use the container to hold anything other than the product for which it was designed.

**SYMPTOMS OF POISONING** - A sharp burning sensation in the mouth, throat, and stomach. Contact dermatitis of the skin. Irritation of the eyes.

**TREATMENT** - In case of skin contact, wash the affected area with plenty of water. If the liquid has been swallowed, do not induce vomiting. Give the patient plenty of water to drink. If the patient has inhaled the dust, remove the patient to fresh air and give plenty of water to drink.

**TO PHYSICIAN—ANTIDOTE** - Do not use for any other purpose. Do not use the container to hold water, milk, or other liquid. Do not use the container to hold other agricultural chemicals. Do not use the container to hold anything other than the product for which it was designed.

**CAUTION** - Do not use for any other purpose. Do not use the container to hold water, milk, or other liquid. Do not use the container to hold other agricultural chemicals. Do not use the container to hold anything other than the product for which it was designed.

**PATENT LICENSE**

Patent license information regarding the use of the product and the container. Do not use for any other purpose. Do not use the container to hold water, milk, or other liquid. Do not use the container to hold other agricultural chemicals. Do not use the container to hold anything other than the product for which it was designed.

**NOTICE** - Do not use for any other purpose. Do not use the container to hold water, milk, or other liquid. Do not use the container to hold other agricultural chemicals. Do not use the container to hold anything other than the product for which it was designed.

Manufactured by  
**W. R. GRACE & CO.**  
AGRICULTURAL PRODUCTS DIVISION  
Portland, Oregon  
Quincy, Wash. - Independence, Ore.

Net weight 50 lbs.