



FERTILIZER

Contains 0.5% DI-SYSTON

USDA Registration No. 9936-7

STATEMENT OF INGREDIENTS FOR INSECT CONTROL

Active ingredient: D-D-threo-TETRAHYDRO-5-HYDROXY-2,4,6-TRIMETHYL-4-PYRIMIDINE SULFONAMIDE. Other ingredients: Inert ingredients.

STATEMENT OF INGREDIENTS AS PLANT FOOD

Active ingredient: Di-Syston. Other ingredients: Inert ingredients.

DIRECTIONS FOR USE

Apply to soil at the rate of 100 lbs. per acre. For broadcast application, use 1/2 lb. per 100 sq. ft. For granular application, use 1/2 lb. per 100 sq. ft. Apply in early fall or early spring.

VEGETABLES

BEANS - Apply to soil at the rate of 100 lbs. per acre. For Snap or Green Lima Beans, apply 1/2 lb. per 100 sq. ft. For Dry Beans, apply 1/2 lb. per 100 sq. ft.

RECOMMENDED DI-SYSTON APPLICATIONS

BRUSSELS SPROUTS, CAULIFLOWER - Apply to soil at the rate of 100 lbs. per acre. For Brussels Sprouts, apply 1/2 lb. per 100 sq. ft. For Cauliflower, apply 1/2 lb. per 100 sq. ft.

PROCCOLI, CABBAGE - Apply to soil at the rate of 100 lbs. per acre. For Proccoli, apply 1/2 lb. per 100 sq. ft. For Cabbage, apply 1/2 lb. per 100 sq. ft.

POTATOES - Apply to soil at the rate of 100 lbs. per acre. For Potatoes, apply 1/2 lb. per 100 sq. ft.

SPINACH - Apply to soil at the rate of 100 lbs. per acre. For Spinach, apply 1/2 lb. per 100 sq. ft.

PEAS - Apply to soil at the rate of 100 lbs. per acre. For Peas, apply 1/2 lb. per 100 sq. ft.

SUGAR BEETS - Apply to soil at the rate of 100 lbs. per acre. For Sugar Beets, apply 1/2 lb. per 100 sq. ft.

NON-BEARING FRUIT

STRAWBERRIES, PROPAGATING PLANTS ONLY - Apply to soil at the rate of 100 lbs. per acre. For Strawberries, apply 1/2 lb. per 100 sq. ft.

RECOMMENDED DI-SYSTON APPLICATIONS

IMPORTANT - When the product is used on fruiting plants, it should be applied to the soil. Do not apply to the fruit. Do not use on plants that are sensitive to the active ingredient.

CONTAINER DISPOSAL

DO NOT RE-USE THE CONTAINER - Do not reuse the container for any other purpose. Do not use the container for food or drink. Do not use the container for any other purpose.



DANGER POISON

Keep out of reach of children



SYMPTOMS OF POISONING - Nausea, vomiting, diarrhea, dizziness, headache, weakness, and loss of consciousness. **TREATMENT** - Induce vomiting, give water, and seek medical attention.

acre. For Snap or Green Lima Beans, apply specified dosage per acre in a band on each side of the seed furrow at planting time only. For Dry Beans, apply specified dosage per acre as a side-dressing on each side of the row after plants become established but not within 60 days of harvest. Do not apply directly to seed. Do not apply more than once per season. **IMPORTANT** statement under "RECOMMENDED DI-SYSTON APPLICATIONS" for additional information before using in this crop.

BRUSSELS SPROUTS, CAULIFLOWER Treat with Amdro Flea Beetle, Leafhopper, Mite, and Thrip, or Braconid, Spint, and Aphid, Flea Beetle, and Leafhopper, or Dinitro, with any 2 of the above. Apply specified dosage per acre at planting time or as a side-dressing after planting. Do not apply directly to seed. Do not apply within 70 days of harvest. Higher rate of dosage for Cabbage. A minimum of 2 days between applications.

BROCCOLI, CABBAGE Treat with Amdro Flea Beetle, Leafhopper, Mite, and Thrip, or Braconid, Spint, and Aphid, Flea Beetle, and Leafhopper, or Dinitro, with any 2 of the above. Apply specified dosage per acre at planting time or as a side-dressing after planting. Do not apply directly to seed. Do not apply within 70 days of harvest. Higher rate of dosage for Cabbage.

POTATOES Treat with Amdro Flea Beetle, Leafhopper, Mite, and Thrip, or Braconid, Spint, and Aphid, Flea Beetle, and Leafhopper, or Dinitro, with any 2 of the above. Apply specified dosage per acre at planting time or as a side-dressing after planting. Do not apply directly to seed. Do not apply within 70 days of harvest. Higher rate of dosage for Cabbage.

SPINACH Treat with Amdro Flea Beetle, Leafhopper, Mite, and Thrip, or Braconid, Spint, and Aphid, Flea Beetle, and Leafhopper, or Dinitro, with any 2 of the above. Apply specified dosage per acre at planting time or as a side-dressing after planting. Do not apply directly to seed.

TOMATOES Treat with Amdro Flea Beetle, Leafhopper, Leaf Mite, and Thrip, or Braconid, Spint, and Aphid, Flea Beetle, and Leafhopper, or Dinitro, with any 2 of the above. Apply specified dosage per acre in a band on each side of the seed furrow at planting time or as a side-dressing after planting. Do not apply directly to seed. Do not apply within 70 days of harvest. Higher rate of dosage for Cabbage.

IMPORTANT statement under "RECOMMENDED DI-SYSTON APPLICATIONS" for additional information before using in this crop.

VEGETABLES GROWN FOR SEED ONLY Treat with Amdro Flea Beetle, Leafhopper, Mite, and Thrip, or Braconid, Spint, and Aphid, Flea Beetle, and Leafhopper, or Dinitro, with any 2 of the above. Apply specified dosage per acre at planting time or as a side-dressing after planting. Do not apply directly to seed. Do not apply within 70 days of harvest. Higher rate of dosage for Cabbage.

FIELD CROPS

ALFALFA Treat with Amdro Flea Beetle, Leafhopper, Mite, and Thrip, or Braconid, Spint, and Aphid, Flea Beetle, and Leafhopper, or Dinitro, with any 2 of the above. Apply specified dosage per acre at planting time or as a side-dressing after planting. Do not apply directly to seed.

BARLEY, OATS Treat with Amdro Flea Beetle, Leafhopper, Mite, and Thrip, or Braconid, Spint, and Aphid, Flea Beetle, and Leafhopper, or Dinitro, with any 2 of the above. Apply specified dosage per acre at planting time or as a side-dressing after planting. Do not apply directly to seed.

CORN Treat with Amdro Flea Beetle, Leafhopper, Mite, and Thrip, or Braconid, Spint, and Aphid, Flea Beetle, and Leafhopper, or Dinitro, with any 2 of the above. Apply specified dosage per acre at planting time or as a side-dressing after planting. Do not apply directly to seed.

be more pronounced in light sandy soils. Plant injury may also occur or may be enhanced when certain pre-emergent herbicides are used in conjunction with DI-SYSTON.

CONTAINER DISPOSAL

DO NOT RE-USE THE CONTAINER Completely empty the container and burn the unused chemical at least 10 inches deep in a hole 100 feet away from water supply. Burn the empty container completely. Burn the container at least 100 feet away from the hole. **Do not use** container or any part of it for food or drinking water.



DANGER POISON



Keep out of reach of children

Do not use if you are sick, pregnant, or nursing. Do not use if you are allergic to any of the ingredients. Do not use if you are taking any medicine. Do not use if you are taking any other medicine. Do not use if you are taking any other medicine. Do not use if you are taking any other medicine.

When using this product, wear protective clothing, including gloves, shoes, and socks. Do not eat, drink, or smoke while using this product. Do not use if you are sick, pregnant, or nursing. Do not use if you are allergic to any of the ingredients. Do not use if you are taking any medicine. Do not use if you are taking any other medicine.

SYMPTOMS OF POISONING Aversion to light, nausea, vomiting, diarrhea, and abdominal pain.

TREATMENT In case of poisoning, call a physician immediately. Do not induce vomiting unless directed by a physician. Do not give anything by mouth to someone who is unconscious or who is having seizures. Do not give anything by mouth to someone who is having seizures.

TO PHYSICIAN — ANTIDOTE Administer activated charcoal immediately. Do not give anything by mouth to someone who is unconscious or who is having seizures. Do not give anything by mouth to someone who is having seizures.

CAUTION Do not use if you are sick, pregnant, or nursing. Do not use if you are allergic to any of the ingredients. Do not use if you are taking any medicine. Do not use if you are taking any other medicine.

PATENT LICENSE

This product is a registered trademark of W. R. Grace & Co. Do not use if you are sick, pregnant, or nursing. Do not use if you are allergic to any of the ingredients. Do not use if you are taking any medicine. Do not use if you are taking any other medicine.

NOTICE Do not use if you are sick, pregnant, or nursing. Do not use if you are allergic to any of the ingredients. Do not use if you are taking any medicine. Do not use if you are taking any other medicine.

Manufactured by
W. R. GRACE & CO.
AGRICULTURAL PRODUCTS DIVISION
Portland, Oregon
Quincy, Wash. Independence, Ore.

Net weight 80 lbs.