



PUBLIC HEALTH SERVICE

**Dichlorvos Aircraft  
 Disinsection Cartridge**

USE AND DIRECTIONS FOR  
 For the control of mosquitoes (Sanitary Regulations), flies and other insects. Apply dichlorvos vapor as described in publication "The Dichlorvos Insecticide System", DHEW USPHS. Use cartridge only once.

GENERAL DIRECTIONS

Apply dichlorvos as a vapor by the cartridge (ADC-20100) containing 4.2 grams of active ingredients to the USPHS aircraft disinsection system. Operate system for not less than 30 minutes over 35 minutes while the aircraft is in flight, landing or taxiing. Stop the system in all emergency situations. Use a dosage of .20-.25 grams per liter of air.

Packed for U. S. PUBLIC HEALTH SERVICE  
 U. S. D. A. Registration  
 by Wix Corporation, Gastonia, N. C.

**WARNING!**

KEEP OUT OF REACH OF CHILDREN, domestic animals, and pets. Poisonous if swallowed, inhaled, or absorbed through skin. Do not swallow or get in eyes, or on skin, or on clothing. Wash hands after handling cartridge. Call a physician immediately in all cases of suspected poisoning. Wash thoroughly with soap and water in case of contact. Remove all spillage by washing with large amounts of water. If the chemical gets into the eyes, flush with water for at least 15 minutes and get medical attention.

**DANGER**

DO NOT STORE ABOVE 120 FAHRENHEIT (APPROXIMATELY 50 CENTIGRADE)

NET CONTENTS 4.2 GRAMS



ANTIDOTE—Atropine is the emergency antidote for Dichlorvos (DDVP).

ACTIVE INGREDIENTS  
 2,2-Dichlorovinyl dimethyl phosphate\* 30% by wt  
 INERT INGREDIENTS  
 Dichlorodifluoromethane 70% by wt

\*Dichlorvos (DDVP) TOTAL 100% by wt



0-1  
 ACCEPTED  
 23 ALB  
 PESTICIDE  
 PESTICIDE  
 REGISTERED  
 COMMENTS

**HEALTH SERVICE**  
**Dichlorvos Aircraft**  
**Disinsection Cartridge**

**DANGER**

DO NOT STORE  
 ABOVE 120  
 FAHRENHEIT  
 (APPROXIMATELY  
 50 CENTIGRADE)

CONTENTS  
 4.2 GRAMS



**POISON**

ANTIDOTE—Atropine is the emergency antidote for Dichlorvos (DDVP).

ACTIVE INGREDIENTS  
 2,2-Dichlorovinyl dimethyl phosphate\* 30% by wt  
 INERT INGREDIENTS  
 Dichlorodifluoromethane 70% by wt  
 \*Dichlorvos (DDVP) TOTAL 100% by wt

**USE AND DIRECTIONS FOR APPLICATION**  
 For the control of mosquitoes (WHO International Sanitary Regulations), flies and other small flying insects. Apply dichlorvos vapor ONLY with system described in publication "The Dichlorvos Aircraft Disinsection System", DHEW USPHS NCDC 1966. Use cartridge only once.

**GENERAL DIRECTIONS**  
 Apply dichlorvos as a vapor by attaching the cartridge (ADC-20100) containing 4.2 grams of active ingredients to the USPHS aircraft disinsection system. Operate system for not less than 30 minutes nor over 35 minutes while the aircraft is taking off, in flight, landing or taxiing. Stop the system in emergencies. Use a dosage of .20-.30 micrograms per liter of air.

**Packed for U. S. PUBLIC HEALTH SERVICE, Savannah, Ga.**  
**U. S. D. A. Registration No. 9990-1**  
**by Wix Corporation, Gastonia, North Carolina 28052**

