

DIRECTIONS FOR USE

OLD SCRATCH CATTLE OIL is a ready-to-use formulation for the control of HORN FLIES on beef cattle. A reduction in FACE FLY population will take place if beef cattle rub their heads and faces on the back rubber device. This treatment also aids in controlling LICE on beef cattle and swine.

Old Scratch Cattle Oil should be used in Old Scratch or similar tank type automatic back rubber devices that are constructed low enough to allow cattle to rub their heads.

Discontinue use 30 days before slaughter.

Do not make accessible to dairy animals.

Do not reuse empty drum. Return to drum reconditioner or destroy by perforating or crushing and burying in a safe place.

NOTICE: Seller makes no warranty, express or implied, concerning this material, except that it conforms to the chemical description on the label. The seller shall not be held responsible in any manner for any personal injury or property damage or other type of loss resulting from the handling, storage or use of this material. The buyer assumes all risk and liability therefrom and accepts and uses this material on the conditions stated.

Formula No. 1

Net Contents

30 Gallons


Old Scratch[®] CATTLE OIL

CONTAINS 5% TOXAPHENE

ACTIVE INGREDIENTS

Toxaphene	5.00%
Mineral Oil	95.00%
	100.00%

*Chlorinated Camphene containing 67 to 69% Chlorine



H & H FARM SERVICE

OLD SCRATCH DISTRIBUTOR

P.O. BOX 425

MILLEDGEVILLE, ILLINOIS

CAUTION - POISON

KEEP OUT OF REACH
OF CHILDREN

CAUTION: May be fatal if swallowed! May be absorbed through skin. Do not get in eyes, on skin or on clothing. Wash hands thoroughly with soap and warm water after handling. Do not contaminate food or feed products. Keep out of reach of children. Do not breathe spray mist. Do not use or store near heat or open flame.

ANTIDOTE-INTERNAL: Use emetic made by dissolving 15 grams (1/2 ounces) of Sodium Chloride (table salt) in glass of warm water and repeat until vomit is clear. If not vomited, give 15 grams of Sodium Chloride in 150 ml of water. If not vomited, give 15 grams of Sodium Chloride in 150 ml of water. If not vomited, give 15 grams of Sodium Chloride in 150 ml of water. If not vomited, give 15 grams of Sodium Chloride in 150 ml of water.

USDA REG. NO. 10004-1