

Dobercide 325

ACCEPTED
with COMMENTS
in EPA Letter Dated

June 1983

Under the authority of Insecticide
Fungicide and Rodenticide Act
as amended by the
FIFRA Act of 1972
10076-10

ACTIVE INGREDIENT:

Poly[oxyethylene (dimethyliminio) ethylene-
(dimethyliminio) ethylene dichloride] 25.0 %

INERT INGREDIENT 75.0 %

This product contains 2.19 lb. of active ingredient per gallon and weighs
8.79 lb. per gallon.

KEEP OUT OF REACH OF CHILDREN

CAUTION

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION: Harmful if swallowed. Avoid breathing vapors. Avoid contact with skin, eyes, or clothing.

FIRST AID: If swallowed, drink promptly a large quantity of milk, egg whites, gelatin solution, or, if these are not available, drink large quantities of water. Avoid alcohol. Call a physician immediately.

ENVIRONMENTAL HAZARDS: This pesticide is toxic to fish. Keep out of lakes, streams, or ponds. Permits may be required for discharges containing this pesticide into lakes, streams, ponds, or public water. For guidance, contact the regional office of the Environmental Protection Agency.

GENERAL CLASSIFICATION

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Dobercide 325 is used to control algae, bacteria, and fungi in recirculating commercial and industrial water cooling towers. Prior to its use, systems must be cleaned to remove algal growth, microbiological slime, and other deposits. An initial slug addition of 2.38 to 5.8 fluid ounces of Dobercide 325 per 1000 gallons of water to provide a concentration of 19.2 to 48 parts per million of Dobercide 325, based on the total weight of water in the system, is recommended. Repeat initial dosage until control is evident.

Subsequent slug additions of 0.58 to 5.8 fluid ounces of Dobercide 325 per 1000 gallons of water (4.8 to 48 parts per million of Dobercide 325) should be employed every 2 to 5 days, or as needed. The frequency of addition depends upon the relative amount of bleedoff and the severity of the microbiological problem. Slug additions should be made in the sump of water cooling towers.

Dobercide 325 is used to control bacteria in industrial air-washing systems that maintain effective mist eliminating components. Prior to its use, systems should be cleaned to remove bacterial slime and other deposits. An initial slug dose of 8.8 to 14.7 fluid ounces of Dobercide 325 per 1000 gallons of water is recommended. Repeat initial dosage until control is evident. Subsequent slug additions of 5.95 to 14.7 fluid ounces of Dobercide 325 per 1000 gallons of water should be employed each 1 to 5 days, or as needed. The frequency of addition depends upon the relative amount of bleedoff and severity of the bacterial problem. Slug additions may be made to the sump or to the water collection trays of the airwash system.

STORAGE & DISPOSAL: Keep container closed when not in use. Do not contaminate water, food, or feed by storage, disposal or cleaning of equipment. Rinseate that cannot be used or reprocessed should be disposed of in a landfill approved for pesticides or buried in a safe place away from water supplies. Open dumping is prohibited.

- METAL CONTAINERS:** Triple rinse and offer for recycling, reconditioning, or disposal in an approved landfill or bury in a safe place.
- PLASTIC CONTAINERS:** Do not reuse empty container. Triple rinse and incinerate or dispose of in an approved landfill or bury in safe place.

Manufactured by:

Dober Water Treatment, Inc.

14465 S. Waverly Avenue

Midlothian, IL 60445

EPA REG. NO. _____

EPA EST. NO. _____

NET CONTENTS _____

U.S. ENVIRONMENTAL PROTECTION AGENCY
REGISTRATION DIVISION
WASHINGTON, D.C. 20460

EPA REGISTRATION NUMBER

DATE OF ISSUANCE

FORM NO. 8570-6

NAME OF REGISTRANT

NOTICE OF PESTICIDE

Under the Federal Insecticide, Fungicide, and Rodenticide Act

NAME AND ADDRESS OF REGISTRANT (Include ZIP code)

<input type="checkbox"/>	<input type="checkbox"/>
<input type="checkbox"/>	<input type="checkbox"/>

NOTE: Changes in labeling formula differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above U.S. EPA registration number.

On the basis of information furnished by the registrant, the above named pesticide is hereby Registered/Reregistered under the Federal Insecticide, Fungicide, and Rodenticide Act.

A copy of the labeling accepted in connection with this Registration/Reregistration is returned herewith.

Registration is in no way to be construed as an indorsement or approval of this product by this Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.

ATTACHMENT IS APPLICABLE

SIGNATURE OF APPROVING OFFICIAL

DATE