

SPRAY-FOG

SYNERGIZED PYRETHRIN SPRAY FOR MILLS, FOOD PLANTS AND THE HOME

ACTIVE INGREDIENTS:

Pyrethrins	.100%	100%
*Piperonyl butoxide, technical	.200%	
N-octyl bicycloheptene dicarboximide	.333%	
Petroleum distillate	99.367%	

*Consists of 0.16% (butylcarbityl) (6-propylpiperonyl) ether and 0.04% related compounds.

This spray is especially designed for use in cafes, restaurants, food storage warehouses, flour mills, meat packing plants and the like. (Meat Inspection Division Memorandum No. 52 - Insect Sprays.) This spray is designed for use in either fogging equipment or hand sprayers.

CAUTION:

Causes irritation of skin and eyes. Avoid contact with skin, eyes, or clothing.

KEEP OUT OF REACH OF CHILDREN

See other precautions on the back panel.

USDA Registration No. 10088-20

ATHEA LABORATORIES, INC.
MILWAUKEE, WISCONSIN

Directions For Use

Flies, Mosquitoes, Gnats: Close room and discharge mist vigorously in all parts of room, using 1½ to 2 ounces for each 1000 cubic feet of space. Leave room closed for at least ten minutes. Sweep up and destroy fallen insects. Repeat as needed.

Fleas (Infesting Premises): Remove and destroy animals' bedding and spray sleeping quarters freely, forcing the spray into all cracks in the floor and about the room. Repeat when re-infested.

Roaches, Waterbugs: Apply in all cracks and crevices in woodwork, underneath sinks, dark places behind pipes, and in all places where roaches and waterbugs infest, contacting as many insects as possible. Repeat the operation as often as necessary.

Clothes Moths: Spray all articles to be protected, giving particular attention to folds and seams. Interior surfaces of storage containers such as closets, chests and trunks should be sprayed regularly. The treatment should be repeated every 30 days unless the articles are stored in moth-tight containers.

Carpet Beetles: Apply thoroughly to infested areas of carpet, localized areas of floors, baseboards and closet shelves, into cracks and under carpets where pests may be found. Repeat application as often as necessary to maintain effective control.

Grain Weevils, Cadelles, Flour Beetles, Grain Mites (Accessible Stages): Wet surface thoroughly and treat all infested surface areas, insect runways and hiding places at the rate of 1 gallon of this material to 1000 square feet of area, or to the point of run-off. Repeat when necessary. Do not spray on grain. WHEN USED IN AN ENCLOSED AREA DO NOT REMAIN IN AREA. VENTILATE AREA WHEN TREATMENT IS COMPLETED.

CAUTION

KEEP OUT OF REACH OF CHILDREN. Harmful if swallowed. Do not apply directly to food. Do not apply while food processing is underway. Thoroughly wash with potable water all treated chutes, processing surfaces, etc. that food will touch prior to re-use. In case of impracticality of washing, do not allow food to come in contact with treated surfaces for at least 48 hours. Remove bird cages and fish bowls during spray application. Do not pour, store or spill near heat or open flame. Keep container closed.

10088-20

7/15/1970

Page 1 of 2

2/2

ACCEPTED
JUL 15 1970
UNDER THE FEDERAL INSECTICIDES
FUNCTIONS AND ROENTGENIDE ACT
FOR ECONOMIC POISON REGISTER-
ED UNDER NO. 12088-20

12088-FA