

## DIRECTIONS FOR USE

**Selective Herbicide #4** is a foliage herbicide for use in controlling certain woody and herbaceous plants growing on drainage ditch banks, fence rows and right-of-ways.

**Selective Herbicide #4** will control: blackjack oak, wild blackberry, ground ivy, northern (white, pin, red, scrub) oaks, post oak, mesquite, maples, poison ivy. Not recommended for ash.

Certain broadleaf weeds are also controlled such as broomweed, cocklebur, croton weed, curly indigo, lambs quarters, Mexican weed, pokeberry, ragweed, sunflower.

**FOLIAGE TREATMENT:** Spray woody growth up to 6 or 8 feet tall after the foliage is fully developed. Use spray mixture consisting of 4-6 quarts Selective Herbicide #4 per 100 gallons of water. Cover all plant parts including leaves and stems with a drenching spray. For knapsack application use 1 pint Selective Herbicide #4 per 10 gallons water.

**STUMP TREATMENT:** Use 4-6 quarts Selective Herbicide #4 per 100 gallons of water; for knapsack application use 1 pint Selective Herbicide #4 in 10 gallons of water. New sprouts from recently cut stumps should not be sprayed until they have good growth. Applications can generally be effectively made up until 3 weeks before killing frost. Application after leaves have lost their normal green color may not give effective control. Less effective control may also result when application made during hot, dry weather. A single application is usually effective for any one year. If new growth develops, repeat application may be necessary in succeeding years.

**AIR APPLICATION:** Use 4 quarts Selective Herbicide #4 in 1 gallon of diesel oil and 2 to 4 gallons of water; or 4 quarts of Selective Herbicide #4 in 3 to 4 gallons of water; or 4 quarts of Selective Herbicide #4 in 3 gallons of diesel oil per acre. Make application when foliage is fully developed (May or June).

An additional spray may be necessary the second or third year. These applications will also control broadleaf weeds in the sprayed areas.

# SELECTIVE HERBICIDE # 4

CONTAINS 2# SILVEX ACID EQUIVALENT  
PER GALLON

### ACTIVE INGREDIENTS:

* Isooctyl ester of Silvex [2 (2,4,5-trichlorophenoxy) propionic acid]	33.19%
INERT INGREDIENTS	66.81%
TOTAL	100.00%

\*Equivalent to 23.43% (2 (2,4,5-trichlorophenoxy) Propionic Acid)

## CAUTION

Keep out of the reach of children.

May cause skin irritation.

Do not take internally, harmful if swallowed

Avoid contact with eyes, skin or clothing.

Avoid inhaling spray mist

Wear goggles while handling concentrate during mixing operations.

### NET CONTENTS

EPA Reg. No. 10088-43-AA-

EPA Est. No. 10088-WI-1

Sold by:

Do not reuse empty drum. Return to drum re-conditioner or destroy by perforating or crushing and burying in a safe place away from water suppliers.

### WEED CONTROL IN LAWNS AND GOLF COURSE:

Control of broadleaf, plantain, buckhorn, common chickweed, mouse-eared chickweed, dandelion, knotweed, white clover and black medic in established turf can be obtained using 3 quarts Selective Herbicide #4 in 25 to 40 gallons of water per acre. Apply in early spring or fall. Hot weather may cause damage to turf. Do not apply to new turf or to bent, Bermuda and St. Augustine grasses. This product can cause possible injury to Dichondra. For spot treatment with knapsack equipment use 1 cup Selective Herbicide #4 in 3 gallons of water.

### WARNING

This product is toxic to fish. Keep out of lakes, streams or ponds. Apply this product only as specified on this label.

This product will kill or seriously injure many desirable forms of vegetation. Avoid spray drift to susceptible plants, as this product may injure cotton, fruits, vegetables, grapes, ornamentals, etc. Do not use when there is hazard from drifting spray mists. (Coarse sprays are less likely to drift).

Vapors from this product may injure susceptible plants in the immediate vicinity.

Do not contaminate water in irrigation ditches or that used for domestic purposes. Excessive amounts of this product in the soil may temporarily inhibit seed germination and plant growth.

Do not store near fertilizers, seeds, insecticides or fungicides.

Spray equipment used in applying this product should be thoroughly cleaned before using for any other purpose. Use repeated flushing with soap and warm water or suitable chemical cleaner. It is best to use a separate sprayer for application of insecticides and fungicides.

### NOTICE

Seller makes no warranty, expressed or implied, concerning the use of this product other than indicated on the label. Buyer assumes all risk of use and/or handling of this material when such use and/or handling is contrary to label instructions.