

DIRECTIONS

IMPORTANT — Use a high pressure sprayer for surface sprays. Avoid contamination of feed and feeding troughs. **RESIDUE TOLERANCE:** Poultry-4 ppm in meat and by products Eggs-0 ppm.

DIRECT APPLICATION FOR NORTHERN FOWL MITE, POULTRY, LICE AND CHICKEN RED MITE*. Use 2 tbs. Insecticide Concentrate per gallon of water per 100–150 birds. Repeat application in 4–8 weeks or when necessary.

*As a supplement to premise treatment for chicken red mite.

ROOST PAINT FOR CHICKEN RED MITE AND POULTRY LICE. Use 2–7 ounces of insecticide concentrate per gallon of water. Brush on at a rate of 1 pint per 150 feet of roost.

PREMISE TREATMENTS FOR NORTHERN FOWL MITE, CHICKEN RED MITE, POULTRY LICE AND FLIES. Use 4 tbs. of insecticide concentrate per gallon of water. Apply liberally to litter, walls, ceilings, roost nests, and adjacent areas. Force spray into cracks and crevices.

CAUTION

This product is toxic to fish. Keep out of lakes, streams, or ponds. Do not contaminate water by cleaning of equipment, or disposal of wastes.

Apply this product only as specified on this label.

EMULSIFIABLE INSECTICIDE CONCENTRATE MB

A Concentrated Insecticide for use on Poultry
(Chickens, Ducks, Geese, and Turkeys) and
Poultry areas.

Active Ingredients:		
Malathion*	57%
Heavy Aromatic Naphtha	34%
Inert Ingredients	9%
	Total	100%

*O, O -Dimethyl dithiophosphate of diethyl mercaptosuccinate

CAUTION

Keep Out of Reach of Children

Harmful if swallowed. Avoid breathing of dust or spray mist. Avoid contact with skin. Wash thoroughly after using. Avoid contamination of Feed or Foodstuffs. Keep away from heat, sparks & open flames. Read entire label before using product.

EPA Reg. No. 10088-58-AA
EPA Est. No. 10088-WI-1

Sold by:

W. A. LAPOSA, INC.
1000 E. BIRD ST.
DALLAS, TEXAS

PREMISE TREATMENTS FOR POULTRY TICKS
Use 6–7 ounces per gallon of water. Apply liberally to walls, ceilings, and adjacent areas. Force spray into cracks and crevices.

TAIL DIPPING FOR NORTHERN FOWL MITE, CHICKEN BODY LICE, AND SHAFT LICE. Use 8½ ounces per 15 gallons of water per 400 birds. Hold bird by wings and dip 3 to 4 inches of tail into solution. Treat vent and surrounding areas. Repeat in 7–10 days if necessary.

“Do not reuse empty drum. Return to drum reconditioner or destroy by perforating or crushing and burying in a safe place away from water supplies.”

NET CONTENTS