

# PROKIL Cryolite 75 Dus

ACTIVE INGREDIENTS
SODIUM FLUOALUMINATE\*
INERT INGREDIENTS

72 0 28 0

100 0

\*Flourine expressed as elemental

40.6

#### CAUTION

KEEP OUT OF REACH OF CHILDREN

Avoid Inhalation of Dust

THIS PRODUCT HIGHLY TOXIC TO BEES exposed to treatment or residues on crops. Protective information may be obtained from your Cooperative Agricultural Extension Service.

### NET WEIGHT 50 LBS.

READ ENTIRE LABEL. USE STRICTLY IN ACCORDANCE WITH CAUTIONS, WARNINGS AND DIRECTIONS, AND WITH APPLICABLE STATE AND FEDERAL REGULATIONS. USES, TIMING AND DOSAGE MAY VARY AS A CONSEQUENCE OF LOCAL WEATHER OR CONDITIONS. WE RECOMMEND REFERENCE TO STATE AGRICULTURAL EXTENSION SERVICE CONCERNING SPECIFIC USES.

KEEP PESTICIDE IN ORIGINAL CONTAINER!

DO NOT REUSE CONTAINER. DESTROY WHEN EMPTY.

Use at rate of 17 to 40 pounds per acre depending on plant size and insect population. Repeat application as necessary. Remove residues remaining at harvest by washing, brushing or other effective means. Do not apply under conditions involving possible drift to food, forage or other planting that might be damaged or the crops thereof rendered unfit for sale, use or consumption.

6-72-113A IM

MARED BUILDING A

LEANS Merican bean decrees Bean Leaf. office -- Repeat at two week in

CABRACIE CAULIFLOWER, BROCCOLL. Flea Beetles, Cabbage Caterpillar NOT APPLY to Broccoli after the ed bile parts start to form. DO NO to Cabbage after the heads start to form. DO NOT APPLY to Cauliflothe seedling stage.

CARROTS: Vegetable Weevil.

CUCUMBERS: Striped Cucumber Beetle, Twelve-Spotted Ci Beetle, Flea Beetle — Dust both sides of foliage at first : Beetles.

EGGPLANT: Flea Beetle - Dust thoroughly.

LETTUCE: Climbing Cutworms, Alfalfa Looper, Corn Earworm — Approxims first appear. DO NOT APPLY after the seedling stage.

MELONS: Cucumber Beetle — Dust both sides of foliage. Repeat if n

FEPPERS. Pepper Weevil — Dust thoroughly at first set of fruit. Repeat days until about 6 applications have been made.

STRAWBERRIES: Leaf Beetle - Dust both sides of leaves.

TOMATOES: Flea Beetle — Apply when plants are first set out and renecessary. If applied after fruit starts to set, remove residue at ha washing, bushing or other effertive means.

#### NOTICE ON CONDITIONS OF SALE

The property of the product are based upon tests believed at the later of the product are based upon tests believed at the later of the product are based upon tests believed at the later of the product of the manufacturation of the product of the effects of such or the resultance of the later of the later of the later of the product of the later of the l

CALIFORNIA (Plant)
P. O. Box 406
340 East Main Street
Calipatria, Calif. 92233
Telephone: 348-5150

## The Dune Company

AGRICLUTURAL CHEMICALS

Route 1, B

Telephone: '627