

DIRECTIONS
For ground application, use from 5 to 15 gallons of water. For aerial application, use from 3 to 10 gallons of water.

COTTON
Do not feed treated cotton trash to dairy animals or animals being finished for slaughter. Drift may contaminate lakes or streams. The dosage depends on size of plants. Repeat application as necessary.

ORNAMENTALS
CANKERWORM AND FALL WEBWORM (on elm and Maple) apply 4 quarts per acre. **CATALPA WORM** (on Catalpa). **TENT CATERpillARS** (on Oak, Willow, Poplar) Use 2 quarts Toxaphene E-6 in 100 gallons of water. **MAY BEETLES** (on Nursery Stock of Elm, Maple, Oak). **BAGWORM** (on Evergreens): Use 1 1/2 to 2 quarts per 100 gallons of water. Make full coverage spray.

LIVESTOCK SPRAYING
DO NOT USE FOR DIPPING Do not use on animals within 30 days of slaughter.
CATTLE For control of Horn Flies and Ticks, dilute 3 qts. per 100 gallons of water. For lice use 1/2 qt. per 100 gallons of water. Agitate thoroughly and apply as spray only; use driving spray to wet skin. 2 qt. of water per adult animal. Repeat as needed. Do not use on dairy cattle or calves under 6 months of age.

SHEEP and GOATS For control of Fleecce Worms and Ticks, Lone Star and Winter Tick, dilute 1 pint in 15 gallons of water. Keep spray agitated, spray animals thoroughly. For Lone Star and Sheep Ticks use 1 pint in 10 gallons of water. Do not apply to dairy goats.

Avoid treating animals during cold weather, overhead or in the rain.

WARNING

May be fatal if swallowed! Do not breathe spray mist. Do not get in eyes, on skin or on clothing. Wash thoroughly after using. Not for use or storage in or around the house.

This product is toxic to fish, birds, and other wildlife. Birds feeding on treated areas may be killed. Keep out of any body of water. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment, or disposal of wastes. Apply this product only as specified on this label.

Avoid contamination of feed and foodstuffs. Do not use full strength. Do not store or pour near open flames. Keep out of reach of children. Do not apply undiluted to animals or to animal quarters. Misuse as to quantity, timing or method of application can cause damage or injury to animals, persons, property or crops or cause residues in excess of official tolerances. Do not reuse empty container. Destroy it by perforating or crushing. Bury or discard in a safe place.

NOTICE: Seller makes no warranty, expressed or implied, concerning the use of this product other than indicated on the label. Buyer assumes all risk of use and/or handling of this material when such use and/or handling is contrary to label instructions.

DO NOT USE, POUR, SPILL OR STORE NEAR HEAT OR OPEN FLAME.

DIRECTIONS
MIXING INSTRUCTIONS: Determine the amount of Toxaphene E-6 to be used in the spray by multiplying the number of acres to be treated by the quantity of Toxaphene E-6 recommended per acre. Add this amount of Toxaphene E-6 to one-half the water required for coverage. Mix well, then add remaining water. Continue mixing until spray mixture is uniform. **BY AIR:** Use 10 to 15 gallons of water per acre. **BY GROUND:** Use 20-50 gallons of water per acre.

ACCEPTED
MAY 1 1975
HARRIS SERUM & SUPPLY CO., INC.
P.O. BOX 410, MCCOOK, NEBR. 68001

HARRIS

TOXAPHENE E-6

6 LBS. PER GALLON

WARNING: Keep Out of Reach of Children

SEE SIDE PANEL FOR ADDITIONAL CAUTIONS.

BEFORE USE READ DIRECTIONS,

WARNING AND NOTICE STATEMENTS CAREFULLY

ACTIVE INGREDIENTS:

Toxaphene*	57.0%
Xylene	31.0%
INERT INGREDIENTS	<u>12.0%</u>
	100.0%

*Technical Chlorinated Camphene-Chlorine content 67.7 to 69.7%

Manufactured for:
HARRIS SERUM & SUPPLY CO., Inc
P.O. Box 410, McCook, Nebr. 68001

EPA REG. NO. 10107-11
EPA EST. NO. 10107-NB-1

NET CONTENTS---GALLON(S)

WARNING

Do not swallow! Do not breathe spray mist. Do not get in
 tin or on clothing. Wash thoroughly after using. Not for
 use in or around the house.

It is toxic to fish, birds, and other wildlife. Birds feeding
 areas may be killed. Keep out of any body of water. Do
 there runoff is likely to occur. Do not apply when weather
 favor drift from areas treated. Do not contaminate water
 of equipment, or disposal of wastes. Apply this product
 cified on this label.

mination of feed and foodstuffs. Do not use full strength.
 e or pour near open flames. Keep out of reach of children.
 y undiluted to animals or to animal quarters. Misuse as
 , timing or method of application can cause damage or
 animals, persons, property or crops or cause residues in
 flicial tolerances. Do not reuse empty container. Destroy
 ating or crushing. Bury or discard in a safe place.

Seller makes no warranty, expressed or implied,
 the use of this product other than indicated on the label.
 nes all risk of use and/or handling of this material when
 id/or handling is contrary to label instructions.

**DO NOT USE, POUR, SPILL OR STORE NEAR
 OR OPEN FLAME.**

DIRECTIONS

INSTRUCTIONS: Determine the amount of Toxaphene E-6
 i the sprayer by multiplying the number of acres to be
 re quantity of Toxaphene E-6 recommended per acre.
 unt of Toxaphene E-6 to one-half the water required for
 ix well, then add remaining water. Continue mixing
 mixture is uniform. **BY AIR:** Use 10 to 15 gallons of water
GROUND: Use 20-30 gallons of water per acre.

ACCEPTED
 MAY 14 1975
 U.S. Dept. of Agriculture
 Agricultural Research Service
 Registration Division
 EPA Reg. No. 10107-11

HARRIS

TOXAPHENE E-6

6 LBS. PER GALLON

WARNING: Keep Out of Reach of Children

SEE SIDE PANEL FOR ADDITIONAL CAUTIONS.

BEFORE USE READ DIRECTIONS.

WARNING AND NOTICE STATEMENTS CAREFULLY

ACTIVE INGREDIENTS:

Toxaphene* 57.0%

Xylene 31.0%

INERT INGREDIENTS:

12.9%

100.0%

*Technical Chlorinated Camphene-Chlorine content 67% to 69%

Manufactured for:
 HARRIS SERUM & SUPPLY CO., Inc.
 P. O. Box 410 McCook, Nebr. 69001

EPA REG. NO. 10107-11
 EPA EST. NO. 10107-N3-1

NET CONTENTS-----GALLON(S)

Do not apply to all species of cucurbits including melons as injury in
 the form of stunting may occur. Do not allow this product to drift
 into cucurbit as injury may occur.

INSECT	AMOUNT LBS. PER ACRE	TIME FOR TREATMENT
Beet Weevil	2-4 pts	From first square until maturity.
Cotton Fleahopper	1 qt	When cotton begins to square.
Cotton Leafworm	2-4 pts	As soon as young worms hatch.
Caterpillars	2-4 pts	Soon after budding cotton is up.
Field Armyworms	2-4 pts	When young worms appear.
Grasshoppers	2-4 pts	Usually early in growing season.
Lyral Bugs	2-4 pts	When cotton begins to square.
Thrips	1-1 1/2 pints	2 to 4 leaf stage of cotton.

This product is toxic to bees and should not be applied when bees
 are actively visiting the area.

ACCEPTED
 MAY 14 1975
 Under the Federal Insecticide,
 Fungicide, and Rodenticide Act,
 as amended, for the pesticide
 registered under
 EPA Reg. No. 10107-11

CROP	
Corn	
Sorghum	
Seed Alfalfa	Y V C C
Beans	A G L
Tomatoes	A C
Lettuce	C L
Small Grains, Wheat Oats Rice Barley Rye	G C A C S in

ARRIS

PHENE E-6

1. PER GALLON

Keep Out of Reach of Children

FOR ADDITIONAL CAUTIONS.

USE READ DIRECTIONS.

NOTICE STATEMENTS CAREFULLY

57.0%
31.0%
12.0%
100.0%

Chlorine content 67% to 69%

CO., Inc. EPA REG. NO. 10107-11
69001 EPA EST. NO. 10107-NB-1

CONTENTS-----GALLON(S)

Do not apply to all species of cucurbits including melons as injury in the form of stunting may occur. Do not allow this product to drift into cucurbit as injury may occur.

INSECT	AMOUNT E-6 PER ACRE	TIME FOR TREATMENT
Beet Weevil	2-3 pts	From first square until maturity.
Cotton Fleahopper	1 qt.	When cotton begins to square.
Cotton Leafworm	2-3 pts	As soon as young worms hatch.
Cutworms	2-3 pts	Soon after seedling cotton is up.
Ear Armyworms	2-3 pts	When young worms appear.
Grasshoppers	2-3 pts	Usually early in growing season.
Lygus Bugs	2-3 pts	When cotton begins to square.
Thrips	1-2 pts	2 to 4 leaf stage of cotton.

This product is toxic to bees and should not be applied when bees are actively visiting the area.

ACCEPTED
 MAY 14 1975
 Under the Federal Insecticide,
 Fungicide, and Rodenticide Act,
 as amended, for the Pesticide
 registered under
 EPA Reg. No. 10107-11

CROP	PEST	AMOUNT E-6	TIME TO SPRAY	LIMITATION
Corn Sorghum	Corn Earworms Chinch Bugs	2-3 pt.	When insects first appear. For corn earworms, apply to silks as soon as they appear. Make 3 to 4 applications of 2-3 day intervals. Re-apply immediately after rain.	Do not apply within 28 days of harvest. Do not apply more than once after heads begin to form. Do not ensile treated forage.
Seed Alfalfa	Yellow Striped Armyworms Webworms Climbing Cutworms Cloverleaf Weevil	2-4 to 4 pts.	When insects first appear.	DO NOT GRAZE DAIRY ANIMALS OR ANIMALS BEING FINISHED FOR SLAUGHTER.
Beans	Armyworms Grasshoppers Lygus Bugs	2-3 pt.	When insects first appear.	Do not apply to GREEN or SNAP BEANS after pods begin to form. Do not feed treated vines to dairy animals or animals being finished for slaughter.
Tomatoes	Armyworms Corn Earworms	2 qt.	When insects first appear.	Do not apply to TOMATOES within 3 days of harvest. Remove excess residues at harvest.
Lettuce	Cabbage looper	3 qts.	When insects first appear.	Do not apply to LEAF LETTUCE after the seedling stage. Do not apply to head lettuce after heads start to form.
Small Grains, Wheat Oats Rice Barley Rye	Grasshoppers Crunch Bugs Armyworms Climbing & Surface feed- ing cutworms	2-3 pt.	when grasshoppers hatch.	Do not feed treated forage to dairy or animals being finished for slaughter. Do not use on forage to be sold commercially or to be shipped interstate. Do not apply more than once after heads have begun to form. Do not apply to wheat, rye, or oats within one week of harvest. Do not apply to barley or rice within two weeks of harvest.