

-RECALL-
WARNING

THERMOPHILIC BACTERIA

卷之三

STORAGE AREA B - SAI

CORNBELT

TOXAPHENE E-6

ACTIVE INGREDIENTS
Sodium *
Copper Sulfate
INERT INGREDIENTS

6 LBS PER GALLON

Keep Out of Reach of Children
WARNING

WARNING

FIRST-AID TREATMENT

INTERNAL: If anyone is swallowed, do not induce vomiting—see "Physician's Note" never give anything by mouth to a unconscious person. Call a physician in all cases suspected of poison.

EXTERNAL: Remove contaminated clothing and wash skin thoroughly with soap and water. If in eyes, flush with clean water for at least 15 minutes.

PHYSICIAN'S NOTE: If swallowed, use gastric lavage. Avoid aspiration. Toxaphene produces central nervous system excitation, and responds to barbiturate acid formulae. Physician should consult with barbituates, preferably amobarbital, intravenously. If the barbiturate can be given a better convulsions start, sodium pentobarbital or the like should be used. Since Toxaphene has a fairly long duration of action, a long-acting barbiturate such as meprobamate may be used after initial control of convulsions. Refer to "Clinical Handbook on Economic Poisons," which was prepared by the U.S. Department of Health, Education and Welfare, Public Health Service, Communicable Disease Center, Atlanta, Ga.

SEE SIDE PANEL FOR ADDITIONAL CAUTIONS

NET CONTENTS 5 GALLONS

DIRECTIONS: For external application, use by water. For internal application, gallipot of 1 oz.

MAXIMUM POSSIBLE POSSIBILITIES
OF THE PROBLEMS

OPTION

BORNAMENT
CROWNED AND FEST