

Cameraman's Note

Poor Copy

INSTRUCTIONS FOR USE (CONTINUED)

General Information

Affacide may be effectively applied by aircraft. Aerial applications may be made on those crops where such application is practical or where application of water to the specific crop and use under the Directions for Use section for the date and timing of application. Mix the recommended amount of affacide in sufficient water to apply a minimum of 2 gallons per acre.

Aerial Application

Affacide is an emulsifiable solution which is to be diluted with water. When ground applications are to be made, dilute Affacide with water according to the Directions for Use section.

Affacide is intended for use against insect pests of agricultural crops and should not be used in barns or dwellings. Applications should be repeated as necessary to maintain control.

Observe the Caution statement and all precautions on the label to avoid any effects to application personnel or crops. Pilots and loaders should wash thoroughly after handling, before eating and at the end of each day's operation.

Affacide is not to be used under conditions where uniform coverage cannot be obtained. Heavy excessive spray drift may occur. Do not fly over power lines or dump contents while in flight.

Directions for Use

Large Insects

Airalfa and Clover (Pure Stands and Stands Containing Grass)

Airalfa Weevil, Larvae, Egyptian Airalfa Weevils, Pea Aphids, Yellow Clover Aphids, Spotted Airalfa Aphids, Leafhoppers, Plant Bug Nymphs, etc. Apply 2.5 quarts per acre in sufficient water to achieve thorough coverage. Apply when 50-60 percent of the growing tips show feeding damage. Repeat as infestations warrant. Do not treat during bloom to avoid injury to beneficial insects.

Airalfa Caterpillars, Armyworms, Fall Armyworms, Flea Beetles, Grasshoppers, Japanese Beetles, Ligus Bugs, Spottedbugs. Apply 2.5 quarts per acre in sufficient water to achieve thorough coverage when damage first appears. Repeat as infestations warrant. Do not treat during bloom to avoid injury to beneficial insects.

Do not feed and use cattle and sheep in any way 7 days following treatment. Do not feed green forage or hay from airalfa or clover to livestock for 7 days following treatment. Do not feed livestock directly from airalfa or clover following treatment.

Small Fruit Insects

For the control of the insects on the following crops, apply 2 1/2 quarts per acre in sufficient water to achieve thorough coverage when insects first appear. Repeat applications at 7 day intervals as infestations warrant.


Cranberries

Cranberry Fruitworms, Leafhoppers, Spottedbugs. Do not apply within 14 days of harvest.

Grapes

Fruit Flies, Grape Berry Moths, Grape Leaf Folders, Grape Leaf Skeletonizers, Japanese Beetles, Leafhoppers, Pacific Mites, Rose Chiggers, For Grape Berry Moths, apply a pre-bloom spray and repeat 10 days later, and then third and fourth applications at 10 day intervals thereafter. Do not apply within 14 days of harvest.

NOTE: The California Department of Agriculture has set a 5 day re-entry interval for Affacide on grapes. If you wish to drop between the time application and the time a worker may enter the field to engage in any activity requiring substantial contact with treated foliage, when a mixture of two or more organic phosphate pesticides are applied in combination, repeat application is required.



Affacide
contains
Diazinon and methoxychlor

For control of insects on
airalfa, clover, vegetables,
and small fruits

Active Ingredients:

- * Methoxychlor, technical..... 20.0 per cent
- * O,O-diethyl O-(2-isopropyl-6-methyl-4-pyrimidinyl) phosphorothioate..... 10.0 per cent
- Aromatic hydrocarbon derivative solvent..... 58.5 per cent

Inert Ingredients:..... 11.5 per cent

Total: 100.0 per cent

* Equivalent to 17.6 per cent 2,2-bis (p-methoxyphenyl)-1,1,1-trichloroethane and 2.4 per cent other isomers and reaction products.
** Diazinon Insecticide

Warning: Keep out of reach of children.

Avoid contact with skin, eyes, and clothing. Do not store near food or feed. May be fatal if swallowed. Do not breathe spray mist.

Fish and Wildlife Caution:

This product is toxic to fish and wildlife. Birds feeding on treated areas may be killed. Do not apply when weather conditions favor drift from areas treated. Do not apply where runoff is likely to occur. Do not contaminate any body of water by direct application, cleaning equipment, or disposal of wastes and containers. Shrimp and crab may be killed at application rates recommended on this label. Do not apply where these are important resources. Apply this product only as specified on this label.

This product is highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your Cooperative Agricultural Extension Service.

Container Disposal:

Do not reuse empty container. Destroy by perforating or crushing and burying in a safe place. Store at temperatures above 32 degrees F.

Broccoli

Aphids, Imported Cabbageworms, Diamondback Moths, Flea Beetles. Do not apply within 14 days of harvest.

Cabbage

Aphids, Imported Cabbageworms, Diamondback Moths, Flea Beetles. Do not apply within 7 days of harvest.

Cauliflower

Aphids, Imported Cabbageworms, Diamondback Moths, Flea Beetles. Do not apply within 7 days of harvest.

Carrots

Aphids, Rootmoppers. Do not apply within 10 days of harvest or 14 days if tops are used for food or feed.

Corn

Aphids, Corn Earworms, Diamondback Moths, Flea Beetles, European Cornborer, European Cabbageworms. Do not apply within 14 days of harvest.

Corn

Fall Armyworms, Flea Beetles, Grasshoppers, Japanese Beetles, Rose Chiggers. Do not apply within 14 days of harvest.

Cucumbers

Aphids, Cucumber Beetles, Serpentine Leaf Miners, Thrips. Do not apply within 7 days of harvest.

Kale

Aphids, Corn Earworms, Diamondback Moths, Flea Beetles, Harlequin Cabbageworm. Do not apply within 14 days of harvest.

Lettuce

Aphids, Diamondback Leaf Miners, Leafhoppers. Do not apply within 14 days of harvest.

Melons (Cantaloupes, Muskmelons, Watermelons)

Aphids, Cucumber Beetles, Leafhoppers, Nymphs, Worms, Mites, Serpentine Leaf Miners, Thrips. Do not apply within 13 days of harvest.

Peas

Aphids, Pea Weevils. Do not apply within 7 days of harvest or during.

Potatoes

Aphids, Banded Cucumber Beetles, Flea Beetles, Colorado Potato Beetles, Diamondback Leaf Miners, Flea Beetles, Leafhoppers, Southern Armyworms. Do not apply within 21 days of harvest.

Radishes

Aphids, Diamondback Leaf Miners, Flea Beetles. Do not apply within 10 days of harvest.

Squash (Summer and Winter)

and should not be used in barns or dwellings. Applications should be repeated as necessary to maintain control. Observe the Caution statement and all precautions on the label to avoid any effects to application personnel or crops. Flagmen and loaders should wash thoroughly after handling, before eating and at the end of each day's operation. Do not apply under conditions where uniform coverage cannot be obtained or where excessive spray drift may occur. Do not follow by lines or dump contents while in flight.

Directions for Use

Large Insects

Alfalfa and Clover (Pure Stands and Stands Containing Grass)

Alfalfa Weevil, Larvae, Egyptian Alfalfa Weevils, Pea Aphids, Yellow Clover Aphids, Spotted Alfalfa Aphids, Leafhoppers, Plant Bug Nymphs, Mites: Apply 2 1/2 qts. per acre in sufficient water to achieve thorough coverage. Apply when 30-50 percent of the growing tips show feeding damage. Repeat as infestations warrant. Do not treat during bloom to avoid injury to beneficial insects.

Alfalfa Caterpillars, Armyworms, Fall Armyworms, Flea Beetles, Grasshoppers, Japanese Beetles, Lygus Bugs, Spittlebugs: Apply 2 1/2-3 qts. per acre in sufficient water to achieve thorough coverage when insects first appear. Repeat as infestations warrant. Do not treat during bloom to avoid injury to beneficial insects.

Note: Dairy and beef cattle and sheep 1) may be grazed 7 days following treatment; 2) may be fed green forage or hay provided alfalfa or clover is not cut for 7 days following treatment. Do not spray livestock directly with either air or ground equipment.

Small Fruit Insects

For the control of the insects on the following crops, apply 2 1/2 qts. per 100 gals. of water per acre as a thorough cover spray when insects first appear. Repeat applications at 7-10 day intervals as infestations warrant.

Cranberries

Cranberry Fruitworms, Leafhoppers, Spittlebugs: Do not apply within 14 days of harvest.

Grapes

Flea Beetles, Grape Berry Moths, Grape Leaf Folders, Grape Leaf Skeletonizers, Japanese Beetles, Leafhoppers, Pacific Mites, Rose Chafers: For Grape Berry Moths, apply as a post bloom cover spray and repeat 10 days later; make third and fourth applications at 10 day intervals if necessary. Do not apply within 14 days of harvest.

Note: The California Department of Agriculture has set a 5 day re entry interval for Diazinon on grapes. Five days should lapse between the time of application and the time a worker may enter the field to engage in any activity requiring substantial contact with treated foliage. When a mixture of two or more organic phosphate pesticides are applied in combination, the interval should be prolonged by adding to the largest applicable interval an additional 50 per cent of that interval.

Strawberries

Aphids, Flea Beetles, Spittlebugs, Strawberry Leaf Beetles, Strawberry Leaf Rollers, Strawberry Weevils, Two Spotted Mites: Do not apply within 5 days of harvest.

Vegetable Insects

For the control of the insects on the following crops use 2 qts. per acre in 100 gals. of water per acre as a thorough cover spray when insects first appear. Repeat applications at 7-10 day intervals as infestations warrant.

DIRECTIONS CONTINUED ON RIGHT PANEL

For control of insects on alfalfa, clover, vegetables, and small fruits

Alfacide

contains
Diazinon and methoxychlor

Active Ingredients:

- * Methoxychlor, technical..... 20.0 per cent
- ** O,O-diethyl O-(2-isopropyl-6-methyl-4-pyrimidinyl) phosphorothioate..... 10.0 per cent
- Aromatic petroleum derivative solvent..... 58.5 per cent

Inert Ingredients:..... 11.5 per cent

Total: 100.0 per cent

* Equivalent to 17.6 per cent 2,2-bis (p-methoxyphenyl)-1,1,1-trichloroethane and 2.4 per cent other isomers and reaction products.
** Diazinon Insecticide

Warning: Keep out of reach of children.

Avoid contact with skin, eyes, and clothing. Do not store near food or feed. May be fatal if swallowed. Do not breathe spray mist.

Fish and Wildlife Caution:

This product is toxic to fish and wildlife. Birds feeding on treated areas may be killed. Do not apply when weather conditions favor drift from areas treated. Do not apply where runoff is likely to occur. Do not contaminate any body of water by direct application, cleaning equipment, or disposal of wastes and containers. Shrimp and crab may be killed at application rates recommended on this label. Do not apply where these are important resources. Apply this product only as specified on this label.

This product is highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your Cooperative Agricultural Extension Service.

Container Disposal:

Do not reuse empty container. Destroy by perforating or crushing and burying in a safe place. Store at temperatures above 32 degrees F.


IMPORTANT: Read the entire Directions For Use and the Conditions Of Sale and warranty before using this product.

NOTICE

Seller makes no warranty, expressed or implied, concerning the use of this product other than indicated on the label. Buyer assumes all risk of use and of handling of this material when purchased and/or handling in a contrary to label instruction.

EPA Est. No. 10107-NB-1
EPA Reg. No. 10107-14

Manufactured for:
HARRIS SERUM & SUPPLY CO., Inc.
P.O. Box 419, McCook, Neb., 68901



Aphids, Imported Catepillars, European Spruce Sawflies, etc. Do not apply within 7 days of harvest.

Carrots

Aphids, Leafhoppers: Do not apply within 10 days of harvest or 14 days if tubers are to be stored or feed.

Corn

European Cornborer, European Corn Rootworm, Fall Armyworm, etc. Do not apply within 14 days of harvest.

Corn

Fall Armyworms, Flea Beetles, Leafhoppers, etc. Do not apply within 14 days of harvest.

Cucumbers

Aphids, Cucumber Beetles, Serpentine Leaf Miners, etc. Do not apply within 7 days of harvest.

Kale

Aphids, Corn Earworms, Diamondback Moths, Flea Beetles, etc. Do not apply within 14 days of harvest.

Lettuce

Aphids, Lettuce Leaf Miners, Leafhoppers: Do not apply within 4 days of harvest.

Melons (Cantaloupes, Muskmelons, Watermelons)

Aphids, Cucumber Beetles, Leafhoppers, Melon Worms, Mites, Serpentine Leaf Miners, etc. Do not apply within 3 days of harvest.

Peas

Aphids, Pea Weevils: Do not apply within 7 days of harvest or grazing.

Potatoes

Aphids, Banded Cucumber Beetles, Blister Beetles, Colorado Potato Beetles, Dipterous Leaf Miners, Flea Beetles, Leafhoppers, Southern Armyworms: Do not apply within 35 days of harvest.

Radishes

Aphids, Dipterous Leaf Miners, Flea Beetles: Do not apply within 10 days of harvest.

Squash (Summer and Winter)

Aphids, Cucumber Beetles, Leafhoppers, Mites, Squash Vine Borers: Do not apply to winter squash within 3 days, or summer squash within 10 days of harvest.

Tomatoes

Aphids, Blister Beetles, Dipterous Leaf Miners, Fall Armyworms, Flea Beetles, Japanese Beetles, Tomato Fruitworms: Do not apply within 14 days of harvest.

Turnips

Aphids, Turnip Leaf Miners, Flea Beetles: Do not apply within 10 days of harvest or 14 days if tubers are to be stored or feed.