

**PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS & DOMESTIC ANIMALS**

WARNING

May be fatal if swallowed. May be absorbed through skin. Do not breathe spray mist. Do not get in eyes, on skin, or on clothing. Wash thoroughly after handling. Avoid contamination of food and feed. Do not use on humans or household pets. Do not permit children or pets to go on to the sprayed grass until spray has completely dried.

ENVIRONMENTAL HAZARDS

This product is toxic to fish, shrimp, crab, birds and other wildlife. Birds feeding on treated areas may be killed. Keep out of lakes, streams, ponds, tidal marshes, and estuaries. Shrimp and crab may be killed at application rates recommended on this label. Do not apply where these are important resources. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment, or disposal of wastes. Apply this product only as specified on this label.

This product is highly toxic to bees exposed to direct treatment or residues on crops.

Protective information may be obtained from your Cooperative Agricultural Extension Service.

PHYSICAL OR CHEMICAL HAZARDS

Do not use, pour, spill, or store near heat or open flame.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

STORAGE AND DISPOSAL

PROHIBITIONS: Do not contaminate water, food, or feed by storage or disposal. Open dumping is prohibited.

PESTICIDE DISPOSAL: Pesticide, spray mixture, or rinsate that cannot be used or chemically reprocessed should be disposed of in a landfill approved for pesticides or buried in a safe place away from water supplies.

CONTAINER DISPOSAL: Triple rinse (or equivalents) and: offer for recycling, reconditioning, or disposal in approved landfill, or bury in a safe place.

GENERAL: Consult federal, state or local disposal authorities for approved alternative procedures.

Do not reuse empty container.

CORN BELT

**DIAZINON[®]
AG 500**

ACTIVE INGREDIENTS:

*O, O-diethyl O-(2-isopropyl-6-methyl-4-pyrimidinyl) phosphorothioate 48.00%

Xylene Range Aromatic Solvent 35.37%

INERT INGREDIENTS: 16.63%

TOTAL 100%

*Diazinon is the Reg. T.M. of CIBA-GEIGY

**CONTAINS 4 LBS.
DIAZINON PER GALLON**

**Keep out of reach of children
WARNING**

If swallowed get medical attention immediately. Do not induce vomiting. If inhaled, remove to fresh air. If on skin, immediately, wash off with plenty of soap and water. In case of eye contact, wash immediately with plenty of water for 15 minutes and get medical attention. - 2-PAM or atropine may be antidotal.

SEE SIDE PANEL FOR ADDITIONAL CAUTIONS.

ACCEPTED

Manufactured by:
CORN BELT CHEMICAL CO.
JUN 8 1978
PO BOX 410
McCook, Nebr. 69001

Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the purpose of registration under EPA Reg. No. 10107-19

NET CONTENTS: 4 GALLON(S)

**GENERAL
INFORMATION**

Cornbelt Diazinon AG 500 is an emulsifiable solution which, when diluted with water according to the directions below, is intended for the control of insect pests of agricultural crops. Do not use in barns or dwellings. Do not use on livestock. In the directions for use given below, best control is obtained when application is made at first sign of insects. Application should be repeated as necessary to maintain control.

USE PRECAUTIONS

Observe number of days before harvest for last application noted in parenthesis by each food crop. Do not apply when honey bees are foraging in the garden.

**FIELD AND FORAGE
FOLIAR INSECTS**

Unless otherwise stated, foliar application of Cornbelt Diazinon AG 500 should always be applied to field & forage crops as a spray in a minimum of 2 gals. of water per acre for aerial applications and a minimum of ten gals. of water per acre for ground application. ALFALFA AND CLOVER - Spotted Alfalfa Aphids, Yellow Clover Aphids, Pea Aphids, Leafhoppers, Plant Bug Nymphs, Mites; Apply 1-1 pt. per acre. Grasshoppers: Apply 1 pt. per acre. Following the above applications, dairy and beef cattle and sheep (1) may be grazed immediately following application (2) may be fed green alfalfa or clover immediately following cutting, and (3) may be fed hay provided alfalfa or clover not cut for hay for 7 days following treatment. Do not spray livestock directly with either air or ground equipment. Do not treat during bloom to avoid injury to pollinating insects. CLOVER - Striped Ground Crickets: Broadcast 1-2 pts. per acre following seeding in the fall but prior to clover emergence.

**FIELD AND
FORAGE
SOIL INSECTS**

Unless otherwise stated, soil application of Cornbelt Diazinon AG 500 should always be applied to field and forage crops as a spray in a minimum of 5 gals. of water per acre for ground applications. ALFALFA AND CLOVER - Cutworms (Surface and Subterranean): Broadcast 2-4 qts. per acre just prior to planting. Work into the soil immediately 2-3 inches for Surface Cutworms, 3-6 inches for Subterranean Cutworms.

**READ CAREFULLY
AND FOLLOW ALL
LABEL DIRECTIONS WHEN
USING THIS PRODUCT**

Cornbelt Diazinon AG 500 is registered for use on the following crops:

- | | |
|------------------|---------------|
| Alfalfa | Olives |
| Almonds | Onions |
| Apricots | Oranges |
| Beets | Ornamentals |
| Bermudagrass | Parsnips |
| Blueberries | Pasture Grass |
| Broccoli | Peaches |
| Brussels Sprouts | Peanuts |
| Cabbage | Peas |
| Caneberries | Pecans |
| Carrots | Peppers |
| Cauliflower | Pineapples |
| Celery | Plums |
| Cherries | Pole Beans |
| Clover | Potatoes |
| Collards | Prunes |
| Cotton | Radishes |
| Cowpeas | Rangeland |
| Cranberries | Snap Beans |
| Cucumbers | Sorghum |
| Endive | Soybeans |
| Field Corn | Spinach |
| Figs | Squash |
| Filberts | Strawberries |
| Grapefruit | Sugar Beets |
| Grapes | Sugarcane |
| Hops | Sweet Corn |
| Kale | Swiss Chard |
| Kumquats | Tangelos |
| Lawns | Tangerines |
| Lemons | Tobacco |
| Lespedeza | Tomatoes |
| Lettuce | Turnips |
| Lima Beans | Turnip Tops |
| Limes | Walnuts |
| Melons | Watercress |
| Nectarines | (Hawaii only) |

**PLEASE SEE BOOKLET
ATTACHED TO THIS
CONTAINER FOR COMPLETE
LIST OF USE DIRECTIONS
FOR INSECT CONTROL
ON THE ABOVE CROPS**

NOTICE: Follow directions carefully. Timing and method of application, weather and crop conditions, mixtures with other chemicals not specifically recommended, and other influencing factors in the use of this product are beyond the control of the seller. Buyer assumes all risk of use, storage or handling of this material not in strict accordance with directions given hereon.

10107-19

6-8-78

10107-19