

10107-42

6-16-96

1/9



United States
Environmental Protection Agency
Washington, DC 20460

Registration
 Amendment
 Other

OPP Identifier Number
213912

Application for Pesticide - Section I

1. Company/Product Number 10107-42	2. EPA Product Manager Dennis Miller	3. Proposed Classification <input type="checkbox"/> None <input type="checkbox"/> Restricted
4. Company/Product (Name) Sevin 4L Insecticide	PM# 19	
5. Name and Address of Applicant (Include ZIP Code) Cornbelt Chemical Company P.O. Box 410 McCook, NE 69001 <input type="checkbox"/> Check if this is a new address	6. Expedited Review. In accordance with FIFRA Section 3(c)(3)(b)(i), my product is similar or identical in composition and labeling to: EPA Reg. No. _____ Product Name _____	

Section - II

<input type="checkbox"/> Amendment - Explain below.	<input type="checkbox"/> Final printed labels in response to Agency letter dated _____	NOTIFICATION JUN 16 1996
<input type="checkbox"/> Resubmission in response to Agency letter dated _____	<input type="checkbox"/> "Me Too" Application.	
<input checked="" type="checkbox"/> Notification - Explain below.	<input type="checkbox"/> Other - Explain below.	

Explanation: Use additional page(s) if necessary. (For section I and Section II.)

Notification to delete unsupported uses for the above product. The unsupported uses for this product include: sweet sorghum.

Section - III

1. Material This Product Will Be Packaged In:				2. Type of Container	
Child-Resistant Packaging <input type="checkbox"/> Yes <input type="checkbox"/> No	Unit Packaging <input type="checkbox"/> Yes <input type="checkbox"/> No	Water Soluble Packaging <input type="checkbox"/> Yes <input type="checkbox"/> No	If "Yes" Unit Packaging wgt. No. per container		<input type="checkbox"/> Metal <input type="checkbox"/> Plastic <input type="checkbox"/> Glass <input type="checkbox"/> Paper <input type="checkbox"/> Other (Specify) _____
3. Location of Net Contents Information <input type="checkbox"/> Label <input type="checkbox"/> Container		4. Size(s) Retail Container		5. Location of Label Directions <input type="checkbox"/>	
6. Manner in Which Label is Affixed to Product			<input type="checkbox"/> Lithograph <input type="checkbox"/> Paper glued <input type="checkbox"/> Stenciled <input type="checkbox"/> Other _____		

Section - IV

1. Contact Point (Complete items directly below for identification of individual to be contacted, if necessary, to process this application.)

Name Tom Kohmetscher	Title Registration Manager	Telephone No. (Include Area Code) 308-345-5057
-------------------------	-------------------------------	---

<p>Certification</p> <p>I certify that the statements I have made on this form and all attachments thereto are true, accurate and complete. I acknowledge that any knowingly false or misleading statement may be punishable by fine or imprisonment or both under applicable law.</p>		<p>6. Date Application Received (Stamped)</p> <p>SIG 8/14/96</p>
2. Signature 	3. Title Registration Manager	
4. Typed Name Tom Kohmetscher	5. Date 4-15-96	

SEVIN® 4L Insecticide

N - METHYL CARBAMATE

FOR PROFESSIONAL PEST CONTROL. FOR
COMMERCIAL USE ONLY.

ACTIVE INGREDIENT:	BY WEIGHT
Carbaryl (1-naphthyl N-methylcarbamate)	43.4% by wt.
INERT INGREDIENTS	56.6% by wt.
TOTAL 100.0%	

EPA Reg. No. 10107-42

EPA Est. No. 10107-NB-1

For MEDICAL and TRANSPORTATION Emergencies ONLY, Call
Chemtree 1-800-424-9300

SEVIN® is the registered trademark of Rhone-Poulenc Ag Company for Carbaryl insecticide.

PRECAUTIONARY STATEMENTS

CAUTION

STATEMENT OF PRACTICAL TREATMENT

GENERAL: Contact a physician immediately in all cases of suspected poisoning. Transport to a physician or hospital immediately and SHOW A COPY OF THIS LABEL TO THE PHYSICIAN. If poisoning is suspected in animals, contact a veterinarian.

IF SWALLOWED: Never give anything by mouth to an unconscious or convulsing person. If conscious and not convulsing, drink 1 to 2 glasses of water and induce vomiting by touching the back of the throat with finger.

IF IN EYES: Flush eyes with plenty of water. Get medical attention if irritation persists.

IF ON SKIN: Wash thoroughly with soap and water.

IF INHALED: Move from contaminated atmosphere and call a physician.

SEE ADDITIONAL PRECAUTIONARY STATEMENTS INSIDE BOOKLET.

Manufactured for:

Corbett Chemical Company, • P.O. Box 410 • McCook, Nebraska 69001

901208

Sevin® 4L Insecticide

N - METHYL CARBAMATE

For professional pest control. For commercial use only.

ACTIVE INGREDIENT:	BY WEIGHT
Carbaryl (1-naphthyl N-methylcarbamate)	43.4% by wt.
INERT INGREDIENTS	56.6% by wt.
TOTAL 100.0%	

For MEDICAL and TRANSPORTATION Emergencies ONLY,
Call Chemtree 1-800-424-9300

SEVIN is the registered trademark of Rhone-Poulenc Ag Company for Carbaryl insecticide.

PRECAUTIONARY STATEMENTS

CAUTION

STATEMENT OF PRACTICAL TREATMENT

GENERAL: Contact a physician immediately in all cases of suspected poisoning. Transport to a physician or hospital immediately and SHOW A COPY OF THIS LABEL TO THE PHYSICIAN. If poisoning is suspected in animals, contact a veterinarian.

IF SWALLOWED: Never give anything by mouth to an unconscious or convulsing person. If conscious and not convulsing, drink 1 to 2 glasses of water and induce vomiting by touching the back of the throat with finger.

IF IN EYES: Flush eyes with plenty of water. Get medical attention if irritation persists.

IF ON SKIN: Wash thoroughly with soap and water.

IF INHALED: Move from contaminated atmosphere and call a physician.

NOTE TO PHYSICIAN

Carbaryl is a carbamate insecticide, which is a cholinesterase inhibitor. Overexposure to this substance may cause toxic signs and symptoms due to stimulation of the cholinergic nervous system. These effects of overexposure are spontaneously and rapidly reversible. Gastric lavage may be used if this product had been swallowed. Carbaryl poisoning may occur rapidly after ingestion and prompt removal of stomach contents is indicated.

SEE ATTACHED BOOKLET FOR COMPLETE LABEL

EPA Reg. No. 10107-42 EPA Est. No. 10107-NB-1

NET CONTENTS: _____

Manufactured for:

CORBELT CHEMICAL COMPANY • P.O. Box 410
McCook, Nebraska 69001

901208

b/c

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS

PERSONAL PROTECTIVE EQUIPMENT

Applicators and other handlers must wear:

- Long-sleeved shirt and long pants
- Chemical-resistant gloves such as barrier laminate, butyl rubber, nitrile rubber, or viton.
- Shoes plus socks.
- Chemical resistant headgear for (overhead exposure).

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and Wash PPE separately from other laundry.

ENGINEERING CONTROLS STATEMENTS

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240 (d) (4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

USER SAFETY RECOMMENDATIONS

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.

May be harmful if swallowed. Avoid breathing spray mist. Do not take internally. Avoid contact with eyes, skin or clothing. Wear regular long-sleeved work clothing and head covering when making overhead applications. Change to clean clothing daily. Bathe and wash hair after each work day. Do not eat, drink, or use tobacco while working with this product or spray solutions. Wash hands and face before eating, drinking or using tobacco. Keep out of reach of children and domestic animals.

OVEREXPOSURE MAY CAUSE: Salivation, watery eyes, pinpoint eye pupils, blurred vision, muscle tremors, difficult breathing, excessive sweating, abdominal cramps, nausea, vomiting, diarrhea, weakness headache. **IN SEVERE CASES CONVULSION, UNCONSCIOUSNESS AND RESPIRATORY FAILURE MAY OCCUR. SIGNS AND SYMPTOMS OF OVEREXPOSURE OCCUR RAPIDLY FOLLOWING EXPOSURE TO THIS PRODUCT.**

ANTIDOTE STATEMENT

ATROPINE SULFATE IS HIGHLY EFFECTIVE AS AN ANTIDOTE. Narcotics and other sedatives should not be used. Further, drugs like 2-PAM (pyridine-2-aldoxime methiodide) are not recommended. See NOTE TO PHYSICIAN below.

NOTE TO PHYSICIAN

Carbaryl is a carbamate insecticide, which is a cholinesterase inhibitor. Overexposure to this substance may cause toxic signs and symptoms due to stimulation of the cholinergic nervous system. These effects of overexposure are spontaneously and rapidly reversible. Gastric lavage may be used if this product had been swallowed. Carbaryl poisoning may occur rapidly after ingestion and prompt removal of stomach contents is indicated.

Specific treatment consists of parenteral atropine sulfate. Caution should be maintained to prevent overatropinization. Mild cases may be given 1 to 2 mg. intramuscularly every 10 minutes until full atropinization has been achieved and repeated thereafter whenever symptoms reappear. Severe cases should be given 2 to 4 mg. intravenously every 10 minutes until fully atropinized, then intramuscularly every 30 to 60 minutes, as needed, to maintain the effect for at least 12 hours. Dosages for children should be appropriately reduced. Complete recovery from overexposure is to be expected within 24 hours.

Narcotics and other sedatives should not be used. Further, drugs like 2-PAM (pyridine-2-aldoxime methiodide) are not recommended. To aid in confirmation of a diagnosis, urine samples should be obtained within 24 hours of exposure and immediately frozen. Consultation of diagnosis therapy and analysis can be obtained at all hours by calling CHEMTREC 1-800-424-9300.

ENVIRONMENTAL HAZARDS

This product is extremely toxic to aquatic and estuarine invertebrates. Do not apply directly to water and wetlands, except under the forest canopy and use on rice. Discharge from rice fields may kill aquatic and estuarine invertebrates. Do not contaminate water by cleaning equipment or disposal of wastes.

BEE CAUTION: MAY KILL HONEYBEES IN SUBSTANTIAL NUMBERS.

This product is highly toxic to bees exposed to direct treatment or residues on blooming crops and weeds. Do not apply this product or allow it to drift to blooming crops or weeds if bees are visiting the treatment area. Contact your Cooperative Agricultural Extension Service or your local Majestic Green Products representative for further information.

DIRECTIONS FOR USE

3/9
IT IS A VIOLATION OF FEDERAL LAW TO USE THIS PRODUCT IN A MANNER INCONSISTENT WITH ITS LABELING. READ THIS LABEL BEFORE USE. STRICTLY OBSERVE LABEL DIRECTIONS AND CAUTIONS, AND APPLICABLE FEDERAL AND STATE REGULATIONS. DO NOT USE ON ANY CROP NOT LISTED ON THIS LABEL OR SUPPLEMENTAL LABELING AS ANY RESIDUES REMAINING MAY BE ILLEGAL AND/OR HARMFUL.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 12 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water is:

- Coveralls
- Chemical-resistant gloves such as barrier laminate, butyl rubber, nitrile rubber, or Viton.
- Shoes plus socks
- Chemical resistant headgear for overhead exposure

NON-AGRICULTURAL USE REQUIREMENTS

The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard for Agricultural Pesticides (40 CFR Part 173). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries, or greenhouses. Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. Do not enter treated areas without protective clothing until sprays have dried.

PREHARVEST INFORMATION AND LIMITATIONS

Tolerances established under the Federal Food, Drug and Cosmetic Act permit the sale of crops bearing probable carbaryl residues when this product is used in accordance with the label directions. Do not apply at greater rates or at more frequent intervals than are stated on the label.

PLANT RESPONSE PRECAUTIONS

To avoid possible injury to tender foliage, do not apply to wet foliage or during periods of high humidity. Do not use on Boston ivy, Virginia creeper and maidenhair fern as injury may result. Carbaryl may also injure Virginia and sand pines.

SPRAY PREPARATION

TO ASSURE A UNIFORM SUSPENSION, AGITATE, STIR OR RECIRCULATE ALL SPRAY MIXTURES OF THIS PRODUCT PRIOR TO AND DURING USE. Remove oil, rust, scale, pesticide residues and other foreign matter from mix tanks and entire spray system. Flush with clean water. Fill spray or mix tank with 1/2 to 3/4 the desired amount of water. Start mechanical or hydraulic agitation. Slowly add the required amount of SEVIN® 4L insecticide and then the remaining volume of water. Include rinse water from the container. Prepare only as much spray mixture as can be applied on the day of mixing. MAINTAIN CONTINUOUS AGITATION DURING MIXING AND APPLICATION TO ASSURE A UNIFORM SUSPENSION. DO NOT STORE SPRAY MIXTURE FOR PROLONGED PERIODS OR DEGRADATION OF CARBARYL MAY OCCUR.

Local water conditions may accelerate the degradation of spray mixtures containing carbaryl. See COMPATIBILITY STATEMENT below.

COMPATIBILITY

SEVIN® 4L Insecticide has been effectively used without plant injury with most common insecticides, miticides, fungicides, nutrients, adjuvants, and with petroleum oil as used on citrus. If compatibility with another product and the resulting crop response is unknown, it should be tested on a small scale. Curdling, precipitation, greasing, layer formation or increased viscosity are symptoms of incompatibility. Incompatibility will reduce insect control and may cause mechanical difficulties or plant injury. Observe all cautions and directions on labeling of all products used in mixtures. WHEN PREPARING COMBINATION SPRAYS, FIRST ADD SEVIN® 4L INSECTICIDE TO 1/2 THE DESIRED AMOUNT OF WATER, MIX THOROUGHLY, AND THEN ADD COMBINATION PRODUCTS TO THE MIXTURE. DO NOT APPLY TANK MIX COMBINATIONS UNLESS YOUR PREVIOUS EXPERIENCE INDICATES THE MIXTURE IS EFFECTIVE AND WILL NOT RESULT IN APPLICATION PROBLEMS, EXCESSIVE RESIDUES OR PLANT INJURY.

Carbaryl is unstable under highly alkaline conditions and mixtures with strong bases such as Bordeaux, lime-sulfur and casein-lime spreaders, will result in chemical degradation of the insecticide. Do not use this product in water with pH values above 8.0 unless a buffer is added. If necessary, water should be buffered to neutral (pH=7.0) before adding this product to the spray tank. Overhead irrigation with alkaline or muddy water after application will also accelerate chemical degradation and may result in reduced insect control.

APPLICATION

On all crops use sufficient gallonage to obtain thorough and uniform coverage. Observe crop label instructions for specific directions regarding spray volume where they occur. Calibrate spray equipment to deliver the required volume. Use 50 mesh slotted strainers in spray system and 25 mesh slotted strainers behind nozzles.

To clean spray system after use, drain and flush with water and detergent mixture. Refer to the Storage and Disposal directions for disposal instructions.

INSECT CONTROL

Apply when insects or their damage appears. To maintain control repeat at 7 to 14 day intervals or as necessary unless a shorter interval is specified below. Where a dosage range is indicated, use the lower rate on light to moderate infestations, young plants and early instars and the higher rate on heavy infestations, mature plants, advanced instars and adults. Thorough and uniform spray coverage is essential for effective control.

NOTE: All references to armyworm on the crops listed below refer to the species, *Pseudaletia unipuncta*, often called the "true armyworm". Except where indicated otherwise, this product is not registered for the control of other armyworm species. Regional differences have been noted in the susceptibility of certain strains of fall armyworm. Colorado potato beetle, spotted tentiform leafminer, and tobacco budworm (on cotton) to carbaryl. If local experience indicates inadequate control, use an alternative pesticide.

TREE FRUIT AND NUT CROPS

Use the recommended amount of Sevin® 4L Insecticide in 100 gallons of water by hand or power-operated sprayer to wet leaves and branches to the dripping point.

FORAGE, FIELD AND VEGETABLE CROPS

Apply in sufficient spray volume to obtain thorough coverage. Where maximum washoff resistance is needed, use up to a 1:1 (vol./vol.) dilution with water. Do not dilute greater than 1:11 where washoff resistance is desired. To prepare small volumes of spray, use 1 tablespoonful (1 fluid ounce) of SEVIN 4L per gallon of water where rates of 1 quart per acre or 1 quart per 100 gallons are indicated in the following tables.

CROP	INSECT	QTS. PRE-HARVEST SEVIN 4L INTERVAL ACRE	(DAYS)	SPECIFIC DIRECTIONS
All Forage, Field and Vegetable Crops in this section	Grasshoppers	1/2 to 1 1/2		See Specific Forage, Field, or Vegetable crop
Alfalfa Clovers	Blister Beetle, Mexican Bean Beetle	1/2 to 1		Observe plant response precautions
	Alfalfa Caterpillar, Bean Leaf Beetle, Cucumber Beetle, Green Cloverworm, Japanese Beetle, Leafhopper, Three-Cornered Alfalfa Hopper, Thrips, Velvetbean Caterpillar	1	3	
	Alfalfa Weevil Larvae, Armyworm, Cloverhead Weevil, Corn Earworm, Cutworm, European Alfalfa Weevil Looper, Essex Skipper, European Alfalfa Beetle, Fall Armyworm, Lygus Bug, Stink Webworm, Yellow-Striped Armyworm	1 to 1 1/2		For Alfalfa Weevil Larvae, if pretreatment damage is extensive, cut alfalfa and treat the stubble. Use higher rate in areas east of the Rocky Mountains. On dense growth, use 25 to 40 gallons per acre with ground equipment to ensure adequate coverage.
Asparagus	Asparagus Beetle	1 to 2	1	Treat ferns or brush growth. Do not treat more than once every 3 days.
	Apache Cicada, Asparagus Beetle	2 to 4	Post-Harvest Application	

	ACRE	(DAYS)	DIRECTIONS	
Beans (including black-eyed peas crowder or Southern peas, dry beans, green beans, lima beans, navy beans, and snap beans)	Blister Beetle, Mexican Bean Beetle	1/2 to 1	Observe plant response precautions.	
	Alfalfa Caterpillar, Bean Leaf Beetle, Cucumber Beetle, Flea Beetle, Green Cloverworm, Japanese Beetle, Leafhopper, Three-Cornered Alfalfa Hopper, Thrips, Velvetbean Caterpillar, Western Bean Cutworm	1		0
	Amyworm, Corn Earworm, Cutworm, European Corn Borer, Fall Armyworm, Stink Bug, Tarnished Plant Bug, Webworm	1 to 1/2		3
Cowpeas	Cowpea Curculio	2	CALIFORNIA ONLY	
	Corn Earworm, Lima Bean Pod Borer, Lygus Bug, Stink Bug	4		
Cabbage Broccoli Brussels Sprouts Cauliflower Kohlrabi	Flea Beetle, Harlequin Bug	1/2 to 2	3	
	Amyworm, Corn Earworm, Fall Armyworm, Imported Cabbageworm	1 to 2		
	Chinese Cabbage Collards	Flea Beetle, Harlequin Bug, Leafhopper		1/2 to 1
Hanover Salad Horseradish, Kale Mustard Greens Radishes Rutabagas Turnips	Aster Leafhopper	1 to 1 1/2	14	
	Amyworm, Corn Earworm, Fall Armyworm, Imported Cabbageworm, Stink bug, Tarnished Plant Bug	1 to 2	(Chinese cabbage, collards, Hanover salad, kale, mustard greens & turnip greens)	
Carrots Parsnips Parsley	Flea Beetle, Leafhopper	1/2 to 1	0	
	Aster Leafhopper, Lygus Bug, Spittlebug	1 to 1 1/2	3	
	Amyworm, Corn Earworm, Fall Armyworm, Stink Bug, Tarnished Plant Bug	1 to 2	14	
Corn (Field, Sweet, Popcorn)	Amyworm, Corn Earworm, Corn Rootworm (Adult), European Corn Borer, Fall Armyworm, Flea Beetle, Japanese Beetle, Sap Beetle, Southwestern Corn Borer, Leafhopper	1 to 2	0	
	Western Bean Cutworm	2	0	
Corn (cont)	Cutworms	2 to 6 1/2	0	
Cotton	Cotton Fleahopper, Cotton Leafworm, Flea Beetle, Striped Blister Beetle, Thrips	1/2 to 1	7	
	Boll Weevil, Bollworm, Cotton Leaf Perforator, Fall Armyworm, Leafroller, Leafhopper, Tarnished Plant Bug, Yellow-Striped Armyworm (Cotton Cutworm)	1 to 2	7	

4/9

CROP	INSECT	QTS. PRE-HARVEST SEVIN 4L INTERVAL /ACRE	INTERVAL (DAYS)	SPECIFIC DIRECTIONS
Cotton	Pink Bollworm	1 1/2 to 2 1/2	7	Aphid populations will be suppressed by repeated applications of this insecticide.
	Stink Bug, Saltmarsh Caterpillar	2		
Cucumber, Melons, Pumpkin, Squash	Pickleworm, Melonworm, Cucumber Beetle, Flea Beetle, Leafhopper, Squash Bug	1/2 to 1	0	Observe plant response precautions. Avoid excessive applications.
Dandelion, Endive (Escarole), Lettuce, Salsify	Flea Beetle, Harlequin Bug, Leafhopper	1/2 to 1	3	Observe plant response precautions. Treat on a 5 to 7 day schedule after heads begin to form.
	Aster Leafhopper, Lygus Bug, Spittlebug	1 to 1 1/2	14	
	Armyworm, Corn Earworm, Fall Armyworm, Imported Cabbageworm, Stink Bug, Tarnished Plant Bug	1 to 2	14	
Forage Grasses, Pasture	Armyworm, Black Grass Bug, Chinch Bug, Essex Skipper, Fall Armyworm, Range Caterpillar, Range Crane Fly, Striped Grass Looper, Thrips	1 to 1 1/2	0	To control thrips in grasses grown for seed use high spray pressure to improve penetration into boot.
	White Grub (June Beetle)	1 1/2 to 2		
Garden Beet, Spinach, Swiss Chard	Flea Beetle, Harlequin Bug, Leafhopper	1/2 to 1	3	Treat on a 5 to 7 day schedule.
	Aster Leafhopper, Armyworm, Corn Earworm, Fall Armyworm, Stink Bug, Tarnished Plant Bug	1 to 1 1/2	14	
Okra	Corn Earworm, Stink Bug	1 to 2	0	Treat on a 5 to 7 day schedule.
Peanuts	Blister Beetle, Mexican Bean Beetle	1/2 to 1	0	Observe plant response precautions.
Peanuts (cont)	Alfalfa Caterpillar, Bean Leaf Beetle, Cucumber Beetle, Green Cloverworm, Japanese Beetle, Leafhopper, Three-Cornered Alfalfa Hopper, Thrips, Velvetbean Caterpillar	1	0	
	Armyworm, Corn Earworm, Fall Armyworm, Stink Bug, Webworm	1 to 1 1/2		
	White-Fringed Beetle adults, Cutworms	2		
Peas	Colorado Potato Beetle, Leafhoppers	1	0	
	Armyworm	1 to 1 1/2		
	Alfalfa Caterpillar, Cutworm, Pea Leaf Weevil, Pea Weevil, Yellow-Striped Armyworm	1 1/2		
	Alfalfa Looper	2 1/2		
Potato, Tomato, Eggplant, Pepper	Colorado Potato Beetle, Flea Beetle, Leafhopper	1/2 to 1	0	
	European Corn Borer, Fall Armyworm, Lace Bug, Stink Bug, Tarnished Plant Bug, Tomato Fruitworm, Tomato Hornworm, Tomato Pinworm	1 to 2		
	Cutworm	2		

CROP	INSECT	QTS. PRE-HARVEST SEVIN 4L INTERVAL /ACRE	INTERVAL (DAYS)	SPECIFIC DIRECTIONS
Rice	Armyworm, Chinch Bug, Fall Armyworm, Stink Bug	1 to 1 1/2	14	DO NOT APPLY PROPANIL HERBICIDES WITHIN 15 DAYS BEFORE OR AFTER SEVIN 4L APPLICATION OR PLANT INJURY WILL RESULT. Do not use on rice fields in which crayfish and/or catfish farming is included in the cultural practice. MISSISSIPPI DELTA & TEXAS
	Armyworm, Leafhopper, Tadpole Shrimp	2	14	
Sorghums (milo, grain sorghum, sweet sorghum & hybrids)	Armyworm, Chinch Bug, Corn Earworm, Fall Armyworm, Stink Bug, Webworm	1 to 2	21 (grain)	Direct spray into forming heads for optimum insect control. Treat for sorghum midge when 25% to 20% of heads have emerged from boot and are in bloom. Repeat application 3 to 5 days later if adults are still active. A third application may be necessary in late planted sorghum or if midges are abundant.
	Sorghum Midge, Southwestern Corn Borer	1 1/2	0 (forage)	
	Cutworm	2		
Soybeans	Bean Leaf Beetle, Cucumber Beetle, Green Cloverworm, Mexican Bean Beetle, Velvetbean Caterpillar	1/2	0	DO NOT APPLY A COMBINATION OF SEVIN 4L and 2,4-D HERBICIDES TO SOYBEANS. For light to moderate populations.
	Corn Earworm	1 1/2 to 3/4		
	Blister Beetle, Grape Colaspis, Mexican Bean Beetle	1/2 to 1		
	Alfalfa Caterpillar, Bean Leaf Beetle, Cucumber Beetle, Green Cloverworm, Japanese Beetle, Leafhopper, Three-Cornered Alfalfa Hopper, Thrips, Velvetbean Caterpillar	1	0	
	Armyworm, Corn Earworm, Cutworm, Fall Armyworm, Stink Bug, Webworm	1 to 1 1/2		
	Painted Lady (Thistle Caterpillar), Saltmarsh Caterpillar, Yellow-Striped Armyworm	2 to 2 1/2		
Sugar Beets	Armyworm, Beet Leaf Beetle, Fall Armyworm, Flea Beetle, Leafhopper, Webworm	1 to 1 1/2	14	
	Cutworm	1 1/2		
Tobacco	Tobacco Flea Beetle	2 lbs./gal. or 1 qt. per 50 gals. or 6 gals. per 100 sq. yds.		IN PLANT BEDS: To prevent injury, avoid excessive applications.

CROP	INSECT	QTS. PRE-HARVEST SEVIN 4L INTERVAL /ACRE (DAYS)		SPECIFIC DIRECTIONS
Tobacco	Green June Beetle Grubs	1/2 qt. per 100 gals.	0	Apply only to areas that larvae have up-rooted by sprinkling as a branch with 50 to 100 gallons per 100 square yards.
	Budworm, Fall Armyworm, Flea Beetle, Hornworm, Japanese Beetle, June Beetle, Suckly	1 to 2		IN FIELDS: Use lower rate on young plants (up to knee height). Use at least 10 gallons of prepared spray per acre. Begin treatments when worms are small. If late season applications are necessary, allow three days before pruning or cutting.

SMALL FRUIT CROPS

Recommended dosages refer to quarts of SEVIN 4L per acre. The optimum spray gallonage will depend on plant size, density and stage of growth. Typical spray gallonages per acre range from 100-300 gallons for dilute sprays, 30-100 gallons for concentrate sprays and 10-25 gallons for aerial sprays. do not exceed maximum label rate per acre per application

CROP	INSECT	QTS. PRE-HARVEST SEVIN 4L INTERVAL /ACRE (DAYS)		SPECIFIC DIRECTIONS
Blackberries Raspberries Dewberries (Including Boysenberries and Loganberries)	European Raspberry Aphid, Japanese Beetle, Leafhopper, Leafroller, Rose Chafer, Snowy Tree Cricket	1 to 2	7	CALIFORNIA ONLY
	Omnivorous Leafroller, Raspberry Sawfly	2		
Blueberries	Blueberry Maggot, Cherry Fruitworm, Cranberry Fruitworm, European Fruit Lecanium, Japanese Beetle	1 1/2	0	Apply 3 weeks before harvest and repeat as necessary.
Cranberries	Cutworm, Cranberry Fireworm, Cranberry Fruitworm, Japanese Beetle, Leafhopper, Rose Chafer	1 1/2 to 3	1	Apply in late bloom and as needed at 7 to 10 day intervals.
	European Fruit Lecanium, Grape Leaf Folder, Grape Leafhopper, Western Grapeleaf Skeletonizer, Western Yellow-Striped Armyworm	1 to 2	7	Apply before first brood leafroller larvae emerge from rofts.
Grapes (cont)	Cutworm, Eight-Spotted Forester, Grape Berry Moth, Japanese Beetle, June Beetle, Orange Tortrix, Omnivorous Leafroller, RedBanded Leafroller, Saltmarsh Caterpillar	2	7	
Strawberries	Flea Beetle, Meadow Spittlebug, Omnivorous Leaf Tier (Strawberry Fruitworm), Strawberry Leafroller, Strawberry Weevil	1 to 2	1	SEVIN may injure Early Dawn and Sunrise varieties on the Delmarva Peninsula

TREE FRUIT AND NUT CROPS

CROP	INSECT	PRE-HARVEST TSP/ GALLON INTERVAL (DAYS)		SPECIFIC DIRECTIONS
Almond	Fruitree leafroller, Peach twig borer, San Jose scale	1	8	OBSERVE BEE CAUTION - Apply in "popcorn" or petal fall stages and again when the May brood of the peach twig borer begins to hatch or thereafter as needed. Time early and midseason applications to correspond to moth flight peaks. Make a late season application at initiation of hull split or up to 10% hull split.
	Navel Orangeworm			
Apples, Pears	Apple aphid, Apple maggot, Apple rust mite, Apple sucker, Bagworms, California pearslug, (pear sawfly), Eastern tent caterpillar, European apple sawfly, Eyespotted bud moth, Forbes scale, Fruitree leafroller, Gypsy moth, Green fruitworm, Japanese beetle, Lecanium scales, Lesser appleworm, Lygus bugs, Orange tortrix, Cystershell scale, Pear leaf blister mite, Pear psylla, Pear rust mite, Periodical cicada, Plum curculio, Redbanded leafroller, Rosy apple aphid, San Jose scale, Tarnished plant bug, Tentiform leafminers, Woolly apple aphid, Yellowheaded fireworm	3/4 to 1	1	To avoid undesired apple thinning, delay use until at least 30 days after full bloom. For psylla control, apply when eggs hatch or young nymphs are present. For optimum scale control, apply when crawlers are present.
Citrus Fruits (such as grapefruit, lemons, limes, oranges, tangelos, tangerines, citrus citron, kumquats, and hybrids)	Avocado leafroller, California orangedog, Citrus cutworm, Citrus root weevil, Fruitree leafroller, Orange tortrix, Western tussock moth, West Indian sugarcane borer (adults)	1	8	Do not apply less than 10 gal. of dilute spray mixture per mature tree. May be mixed with petroleum oils commonly used on citrus.
	Black scale, Brown soft scale, California red scale, Citricola scale, Citrusto 1 snow scale, Yellow scale	3/4		
Olives	Olive scale	3/4 to 1	0	For optimum scale control add 1 1/2 gallons of summer oil and apply mixture when crawlers are present. Do not exceed 2 applications per year
Peaches Apricots Nectarines	Apple pandemis, Codling moth, Cucumber beetle, European earwig, Fruit tree leafroller, Gypsy moth, Japanese beetle, June beetles, Lecanium scales, Lesser peachtree borer, Olive scale, Orange tortrix, Oriental fruit moth, Peach twig borer, Periodical cicada, Plum curculio, Redbanded leafroller, San Jose Scale, Tarnished plant bug, Tussock moths, Variegated leafroller	1	1 (peaches) 3 (apricots & nectarines)	For optimum scale control, apply when crawlers are present. For lesser peachtree borer control, spray limbs and tree trunks thoroughly, weekly during moth flight.
Pecans	Black margined aphid, Fall webworm, Hickory shuck worm, Lesser webworm, Pecan leaf phylloxera, Pecan stem psyllocera, Pecan nut casebearer, Pecan spittlebug, Pecan weevil, Twig girdler, Walnut caterpillar	1 to 2 1/2	0	

CROP	INSECT	PRE-HARVEST TSP/ GALLON INTERVAL (DAYS)		SPECIFIC DIRECTIONS
Almond	Peach twig borer, Fruit tree leafroller, San Jose scale	4	28	Apply in "popcorn" or petal fall stages and again when the May brood of the peach twig borer begins to hatch or thereafter as needed.

LAWNS AND RECREATIONAL AREAS

For control of certain turfgrass pests, apply the recommended rates in sufficient spray volume for thorough coverage and turf thatch penetration. Where a dosage range is indicated, use the higher rate on large pest populations, advanced instars and dense vegetation. Repeat treatment as necessary.

AMOUNT OF SEVIN 4L

FL. OZ/ QTS/

SITE	INSECT	1000 SQ.FT. ACRE	FL. OZ/ QTS/	SPECIFIC DIRECTIONS
Turf grasses	Ants, Armyworm, Centipedes, Chiggers, Cutworms, Earwigs, Essex skipper, European chafer, Fall armyworm, Fiery skipper, Grasshoppers, Green June beetle grub, June beetles, Leafhoppers, Lucerne moth, Millipedes, Mosquitoes (adults), Sowbugs, Spittle bugs, Springtails, Ticks, Yellowstriped army-worms	11/2 to 3	2 to 4	For Armyworm, Cutworm, and Fall Armyworm control: Do not irrigate treated areas following insecticide application. For Green June Beetle Grub control: Make applications when grubs are feeding near the soil surface. Water or irrigate turfgrasses soon after treatment. DO NOT ALLOW PUBLIC USE OF TREATED AREAS DURING APPLICATIONS OR UNTIL SPRAYS HAVE DRIED.
	Chinch bugs	4.4 to 6	6 to 8	Treat entire grass area rather than just damaged areas. Irrigation of turf grass area before treatment will aid in penetration of insecticide into turf grass.
	Sod Webworm (lawn moths)			Do not irrigate treated areas following insecticide application.
Turf grasses (cont.)	Bluegrass billbug, European crane fly, Fleas	6	8	For European Crane Fly Control: Treatments should be applied in early spring, April 1st to April 14th, 15th, or as recommended by local Agricultural Extension Service agents. Water or irrigate turf grasses thoroughly soon after treatment.
	White grubs (such as Japanese beetle Chafer beetle and <i>Phyllophaga</i> spp. larvae)			For White Grub control: Applications should be made when grubs are feeding near the soil surface, usually during late March through May, or July to early September, or as recommended by local Agricultural Extension Service agents. Water or irrigate turf grasses thoroughly soon after treatment.

IMPORTED FIREANT CONTROL

SITE	AMT/GAL OF WATER	SPECIFIC DIRECTIONS
Lawns, Cemeteries and Recreational Areas (including turf, golf courses, and parks) Pastures Rangeland Forested Lands and Waste-land	1-1/2 qts. or 1/2 Tblsp. per gal.	Apply a total of 2 gallons of the diluted solution over the surface of each mound or at least 1 qt. per 6 inches of mound diameter using a bucket, can, or other appropriate equipment. Thoroughly wet mound and surrounding area to a 4 ft. diameter (12 sq. ft.). Do not disturb mounds prior to treatment. Pour solution from a height of about three feet to give sufficient force to break mound apex and flow into ant tunnels. For best results apply in cool weather, 65° to 80° F or in early morning or late evening hours. Repeat application if mound activity resumes after 10 days. Treat new mounds as they appear. Pressurized sprays may disturb the ants and cause migration, reducing product effectiveness. DO NOT ALLOW PUBLIC USE OF TREATED AREAS DURING APPLICATIONS OR UNTIL SPRAYS HAVE DRIED.

ADULT MOSQUITO CONTROL

SITE	AMT/GAL PER 100 GAL.	SPECIFIC DIRECTIONS
Pastures, Rangelands, Parks, Recreational areas Logging Camps, Military Posts and Adjacent forested lands or Waste-lands	1/4 to 1 or 1/2 to 1 Tblsp. per gal.	DO NOT ALLOW PUBLIC USE OF TREATED AREAS DURING APPLICATIONS OR UNTIL SPRAYS HAVE DRIED. Caution: May kill shrimp and crabs. Do not use in areas where these are important resources. OBSERVE BEE CAUTION Treat shrubbery and areas where adult mosquitoes congregate. Treat when adult mosquitoes are active in early mornings or late evenings. Repeat at 7 to 10 day intervals. Use 1/4 to 1/2 quart per 100 gallons in mist-blowers, and 1 quart per acre in low pressure ground sprayers. For residual control in subtropical regions apply 4 gallons of prepared spray per 2000 square feet of surface area. Repeat in 3 to 6 months or when necessary.

CONTROL OF TICKS WHICH VECTOR LYME DISEASE

For control of juvenile and adult ticks which vector Lyme Disease, apply the recommended rate in sufficient volume for thorough coverage. To prepare small amounts, use 1 tablespoon (1/2 fluid ounce) of SEVIN® 4L insecticide per gallon of water.

SITE	PEST	QTS/ACRE	SPECIFIC DIRECTIONS
Lawns and Recreational turf grass (Including: Lawns & Perimeters, Golf Courses, Sports Fields, Cemeteries, Parks, and Pastures) Shrubs, Ornamentals, Wooded areas (Including: Military Posts Logging camps, and Campsites) Wastelands	<i>Ixoides</i> spp. Deer Tick, Bear tick, and Black legged tick <i>Amblyomma</i> spp. Lone Star tick	1 to 2	Use the high rate for heavy tick infestations. Use higher spray volumes for dense ground cover or heavy leaf litter and on lush turf with thatch buildup. Target applications for nymphal control in late spring or early summer. Control of adult ticks can be achieved with late summer and fall applications. Do not use spot treatments. Treat entire area and perimeter areas where exposure to ticks may occur. Ticks may be reintroduced from surrounding areas on host animals. Retreat as necessary to maintain adequate control levels. DO NOT ALLOW PUBLIC USE OF TREATED AREAS DURING APPLICATION OR UNTIL SPRAYS HAVE DRIED.

GENERAL INFORMATION

Staining may occur on certain surfaces such as stucco, brick, cinder block and wood. Therefore, applications of SEVIN® 4L Insecticide to surfaces where a noticeable residue or discoloration is objectionable should be avoided. Do not apply to carpets or draperies as staining may occur. Care should also be exercised to avoid spotting of wallpaper and fabrics.

Do not use this product in commercial food areas of food handling establishments, restaurants or other places where food is prepared or processed. Do not use in serving areas while food is exposed.

INDOORS

Ants, Crickets, Firebrats, and Silverfish: Mix 3 oz. this product per gallon of water and apply as fine, low pressure (20 psi) spot spray or as crack and crevice application to areas where these pests hide, such as baseboards, storage areas, closets, around water pipes, doors and windows, behind and under refrigerators, cabinets, sinks, stoves, dishwashers, hot water heaters, the underside of shelves, drawers and similar areas. For ants, apply to ant trails, around doors and windows and other places where ants enter premises.

Bees and Wasps: Mix 3 oz. this product per gallon of water and thoroughly spray nest and entrance and surrounding areas where insects alight. It is generally advisable to spray the nests in the evening when the insects are less active and have returned to the nest. For best results, check nest carefully one or two days after spraying to ensure complete kill, then remove and destroy nest to prevent emergence of newly hatched insects.

Brown dog Ticks and Fleas: Mix 3 oz. this product per gallon of water and thoroughly spray infested areas such as pet beds and resting quarters, nearby cracks and crevices, between and under cushions of upholstered furniture, along and behind baseboards, window and door frames and other areas where pests may be present.

Carpenter Ants: Mix 3 oz. this product per gallon of water and apply to ant trails, around doors and windows and other places where ants enter premises. Where possible, apply this product directly to ant nest of infested wood.

Centipedes, Earwigs, Millipedes, and Scorpions: Mix 3 oz. SEVIN® 4L Insecticide per gallon of water and apply around water pipes, doors and windows, and other places where these pests may enter premises. Spray baseboards, storage areas, garages, carports, basements and other areas where these pests are found.

Cockroaches: Efficacy varies with species sensitivity. This product is generally not highly effective in controlling German cockroaches. However, the following may be controlled with the rates as stated. American roach, Australian roach, Brown roach, Smoky brown roach, and others. Mix 3 oz. this product per gallon of water and apply as a fine, low pressure (psi) spot spray or as crack and crevice application to areas where these pests hide, such as baseboards, storage areas, closets, around water pipes, doors and windows, behind and under refrigerators, cabinets, sinks, stoves, dishwashers, hot water heaters, the underside of shelves, drawers and similar areas.

Spiders: Mix 3 oz. this product per gallon of water and apply to infested baseboards, window and door frames, corners, pipes, storage areas, attics and eaves. Make spot applications to other areas which these pests are ant.

OUTDOORS

Perimeter Treatment: Residual spray for control of ants, bees and wasps, brown dog ticks, carpenter ants, centipedes, cockroaches, crickets, earwigs, firebrats, and silverfish, fleas, millipedes, scorpions and spiders. Mix 16 oz. this product per 50 gallons of water and apply with power spray or other spray methods.

To help prevent infestations of buildings by the above pests, outside perimeter treatment should be in a band 6 to 10 feet wide and confined to shrub beds, foundation plantings and lawn or soil areas immediately adjacent to the structure. Direct application to structures should be made and restricted to cracks and crevices and other areas where insects tend to congregate.

STORAGE AND DISPOSAL**Storage**

Store unused SEVIN® 4L Insecticide in original container only, in cool, dry area out of reach of children and animals. Do not store in areas where temperatures frequently exceed 100° F. This product may be used following exposure to several freeze-thaw cycles.

If container is damaged, before cleaning up, put on long-sleeved shirt, full length trousers, head covering, and protective gloves.

Pesticide Disposal

Do not contaminate water, food, or feed by storage or disposal. Open dumping is prohibited. Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

Container Disposal

Triple rinse (or equivalent). Then offer for recycling or reconditioning, or dispose of in a sanitary landfill, or incineration, or if allowed by state and local authorities, by burning. If container is burned, stay out of smoke.

LIMITED WARRANTY AND DISCLAIMER

The manufacturer warrants (a) that this product conforms to the chemical description on the label; (b) that this product is reasonably fit for the purposes set forth in the directions for use when it is used in accordance with such directions; and (c) that the directions, warnings and other statements on this label are based upon responsible experts' evaluation of reasonable tests of effectiveness, of toxicity to laboratory animals and to plants, and of residues on food crops, and upon reports of field experience. Tests have not been made on all varieties or in all states or under all conditions. THE MANUFACTURER NEITHER MAKES NOR INTENDS, NOR DOES IT AUTHORIZE ANY AGENT OR REPRESENTATIVE TO MAKE ANY OTHER WARRANTIES, EXPRESS OR IMPLIED, AND IT EXPRESSLY EXCLUDES AND DISCLAIMS ALL IMPLIED WARRANTIES OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE.

THIS WARRANTY DOES NOT EXTEND TO, AND THE BUYER SHALL BE SOLELY RESPONSIBLE FOR, ANY AND ALL LOSS OR DAMAGE WHICH RESULTS FROM THE USE OF THIS PRODUCT IN ANY MANNER WHICH IS INCONSISTENT WITH THE LABEL DIRECTIONS, WARNINGS OR CAUTIONS.

BUYER'S EXCLUSIVE REMEDY AND MANUFACTURER'S OR SELLER'S EXCLUSIVE LIABILITY FOR ANY AND ALL CLAIMS, LOSSES, DAMAGES, OR INJURIES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, WHETHER OR NOT BASED IN CONTACT, NEGLIGENCE, STRICT LIABILITY IN TORT OR OTHERWISE, SHALL BE LIMITED, AT THE MANUFACTURER'S OPTION, TO REPLACEMENT OF, OR THE REPAYMENT OF THE PURCHASE PRICE FOR, THE QUANTITY OF PRODUCT WITH RESPECT TO WHICH DAMAGES ARE CLAIMED. IN NO EVENT SHALL MANUFACTURER OR SELLER BE LIABLE FOR SPECIAL, INDIRECT OR CONSEQUENTIAL DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT.

9/9