

10107-97

12-8-89

1/2

DEC 8 1989

Ms. Sharon K. Snyder, President
Cornbelt Chemical Company
P.O. Box 410,
McCook, Nebraska 69001

Dear Ms. Snyder:

Subject: Seed Shield™ Potato Seed Treater No. 7.5 With
Bark
EPA Registration No. 10107-97
Your Letter Dated November 24, 1989

The label submitted with the subject letter is acceptable under the Federal Insecticide, Fungicide and Rodenticide Act (FIFRA) as amended, provided that:

You have received notice from this Agency's Office of Compliance Monitoring that the EPA suspension of this product has been rescinded prior to any shipment under this label.

If this condition is not complied with, the registration will be subject to cancellation in accordance with FIFRA, Section 6(e). Your release for shipment of the product constitutes acceptance of this condition.

A stamped copy of the label is enclosed for your records.

Sincerely yours,

Joanne I. Miller
Acting Product Manager (23)
Fungicide-Herbicide Branch
Registration Division (H-7505C)

Enclosure

PRECAUTIONARY STATEMENTS
Hazards to Humans and Domestic Animals

DANGER

Causes irreversible eye damage. Harmful if swallowed or inhaled. May cause allergic skin reactions. Do not get in eyes. wear goggles or face shield when handling. Avoid contact with skin and clothing. Remove and separately launder clothing before reuse. Wash hands, arms, and face thoroughly with soap after handling and before eating or smoking. Routinely shower or bathe after work and wash with soap and hot water before reusing.

ENVIRONMENTAL HAZARDS

This pesticide is toxic to fish. Do not apply directly to water or wetlands (swamps, bogs, marshes, and potholes). Drift and runoff from treated areas may be hazardous to aquatic organisms in neighboring areas. Do not clean equipment or dispose of equipment washwaters in a manner that will contaminate water resources. In the event that treated seeds are spilled and cannot be salvaged, completely cover them with soil.

DIRECTIONS FOR USE

It is a Violation of Federal law to use this product in a manner inconsistent with its labeling.

A dust mask must be worn when mixing and loading this product. Mixers/loaders when mixing and loading this product must wear a long-sleeved shirt and long pants or a coverall that covers all parts of the body except the head, hands and feet. Wear chemical resistant gloves, goggles or a face shield and a chemically resistant apron.

AFTER WORK: Take off all clothes and shoes. Shower using soap and water. Do not wear contaminated clothing. Wash protective clothing and equipment with soap and water after each use. Personal clothing worn during use must be laundered separately from household articles. Clothing and protective equipment heavily contaminated or drenched with CAP-TAN must be destroyed according to state and local regulations.

STORAGE AND DISPOSAL

PROHIBITIONS: Do not contaminate water, food, or feed by storage or disposal or cleaning of equipment. Open dumping is prohibited. Do not reuse empty container.

STORAGE: Keep pesticide in original container. Do not put concentrate or dilute into food or drink containers. Store in cool, dry place. Protect from excessive heat. Do not store or transport near feed or toed.

PESTICIDE DISPOSAL: Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous waste representative at the nearest EPA Regional Office for guidance.

CONTAINER DISPOSAL: Completely empty liner into application equipment. Then dispose of empty liner and container in a sanitary landfill or by incineration if allowed by State and local authorities, by burning. If burned, stay out of smoke.

SEED SHIELD™ POTATO SEED TREATER NO. 7.5

WITH BARK
(FUNGICIDE)

ACTIVE INGREDIENTS:	BY WEIGHT
*Captan	7.30%
Related Derivatives	0.20%
INERT INGREDIENTS	92.50%
TOTAL	100.00%

*N-Trichloromethylthio-4-cyclohexene-1, 2-dicarboximide
SEED SHIELD™ registered trademark of Combelt Chemical Co.

KEEP OUT OF REACH OF CHILDREN DANGER - PELIGRO

PRECAUCIONAL USUARIO: Si usted no lee ingles, no use este producto hasta que la etiqueta le haya sido explicada ampliamente.

SEE SIDE PANEL FOR ADDITIONAL PRECAUTIONARY STATEMENTS

Corrosive, Causes Irreversible Eye Damage

STATEMENT OF PRACTICAL TREATMENT

FIRST AID: If a known exposure occurs or is suspected, immediately start life procedures given below and contact a POISON CENTER, PHYSICIAN, OR THE NEAREST HOSPITAL. Describe the type and extent of exposure, the victim's symptoms and follow the advice given.

If in eyes: Flush with plenty of water. Call a physician.

If swallowed: Drink promptly a large quantity of milk, egg whites, gelatin solutions or if these are not available, drink large quantities of water. Avoid alcohol.

If inhaled: Remove victim to fresh air. If not breathing, give artificial respiration, preferably mouth-to-mouth. Get medical attention.

If on skin: Wash with plenty of soap and water. Get medical attention if abnormal reaction occurs. If medical advice cannot be obtained, take person and a copy of the label to the nearest emergency treatment facility.

NOTE TO PHYSICIAN: Emergency information —

call CHEMTREC (800) 424-9300.

See side panel for additional precautionary statements

STORE IN A COOL, DRY PLACE.

EPA Reg. No. 10107-97

EPA Est. No. 10107-NB1

NET CONTENTS 50 POUNDS

MANUFACTURED FOR:

Combelt Chemical Co.

P. O. Box 400 McCook, Nebraska 69001

DIRECTIONS FOR USE (Continued)

Treated seed must not be used for or mixed with food or animal feed, processed. Treat only those seeds needed for immediate use, minimizing the interval between treatment and planting. Do not store excess treated seed beyond planting time. Exposed treated seeds may be hazardous to birds and other wildlife.

PRODUCT INFORMATION: POTATO SEED TREATER NO. 7.5 is an agricultural Fungicide in a dust form containing Douglas fir Bark as the primary inert carrier. It is useful for treating Potato Seed Pieces and Potato Whole Seed for diseases listed and aids in the natural healing process of suberization and wound periderm in cut seed potatoes.

POTATOES: (Seed Pieces) Fusarium Seed Rot, Verticillium Wilt and Damping-off; Dust at 10 oz. per bushel cut seed pieces (1-1/2 lb. to 100 lbs. cut seed pieces). Apply in convenient container or by duster attachment over belt. Cut pieces should be treated within 6 hours of cutting. If planting is to be delayed more than 1 or 2 days the treated pieces should be stored for 2 days in open crates before bagging.

NOTE TO USER: Federal Government and most states have seed laws and regulations requiring an appropriate caution statement be affixed to bags containing treated seed such as: "Treated Seed- Do Not Use for food or feed."

CONDITIONS OF SALE: 1. Combelt Chemical Company (Combelt) warrants that this material conforms to the chemical description on the label and is reasonably fit for use as directed hereon, Combelt neither makes, nor authorizes any agent or representative to make, any other warranty of FITNESS or of MERCHANTABILITY, guarantee or representation, express or implied, concerning this material
2. Critical and unforeseeable factors beyond Combelt's control prevent it from eliminating all risks in connection with the use of chemicals. Such risks include, but are not limited to, damage to plants and crops to which the material is applied, lack of complete control, and damages caused by drift to other plants or crops. Such risks occur even though the product is reasonably fit for the uses stated hereon and even though label directions are followed. Buyer and user acknowledge and assume all risks and liability (except those assumed by Combelt under 1 above) resulting from handling and use of this material.

DEC 8 1989

Under the Federal Insecticide, Fungicide, and Rodenticide Act as amended, for the pesticide registered under EPA Reg. No.

10107-97

2/2