

**PRECAUTIONARY STATEMENTS  
HAZARDS TO HUMANS  
( & DOMESTIC ANIMALS )**

**CAUTION**

May be harmful if swallowed. Do not breathe dust. Do not get in eyes, on skin or on clothing. May cause eye or skin irritation. During exposure in treatment and cleaning operations wear goggles, long pants, long-sleeved shirt, hats and boots. Wash hands, arms, and face thoroughly with soap after handling and before eating or smoking. Routinely shower or bathe after work and wash work clothes with soap and hot water before reusing.

**ENVIRONMENTAL HAZARDS**

This product is toxic to fish, birds, and other wildlife. Do not apply directly to water. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on this label. Exposed treated seeds may be hazardous to birds and other wildlife.

DEC 02 1991

**DIRECTIONS FOR USE**

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

Treated seed must not be used for or mixed with food or animal feed, or processed for oil. Treat only those seeds needed for immediate use, minimizing the interval between treatment and planting. Do not store excess treated seeds beyond planting time.

**STORAGE AND DISPOSAL**

**PROHIBITIONS:** Do not contaminate water, food or feed by storage, disposal or cleaning of equipment. Open dumping is prohibited.

**STORAGE:** Store in cool, dry place. Keep container closed when not in use. If this product becomes wet in storage, its fungicidal effectiveness may be reduced and it may also produce flammable vapors.

**PESTICIDE DISPOSAL:** Pesticide, spray mixture, or rinse water that cannot be used according to label instructions must be disposed of according to applicable Federal, state or local procedures.

**CONTAINER DISPOSAL:** Completely empty liner by shaking and tapping sides and bottom to loosen clinging particles. Empty residue into application equipment. Then dispose of liner in a sanitary landfill or by incineration if allowed by State and local authorities. Do not reuse empty container. Dispose of container in same manner.

**SEED SHIELD™  
POTATO SEED  
TREATER M - Z WITH  
DOUGLAS FIR BARK**

(Contains Dithane® M - 45)

ACTIVE INGREDIENTS:	BY WT.
A co-ordination product of zinc ion and manganese.....	6.40%
ethylene bisdithiocarbamate in which the ingredients are:	
Manganese.....	1.28%
Zinc.....	0.16%
Ethylene bisdithiocarbamate ion.....	4.96%
<b>INERT INGREDIENTS:</b> .....	<b>93.60%</b>
<b>TOTAL</b> .....	<b>100.00%</b>

\*Trademark of Rohm & Haas Co.

SEED SHIELD™ is the registered trademark of Cornbelt Chemical Co.

**KEEP OUT OF REACH OF CHILDREN  
CAUTION**

PRECAUCION AL USUARIO: Si usted no lee ingles, no use este producto hasta que la etiqueta le haya sido explicada ampliamente.

**STATEMENT OF PRACTICAL TREATMENT**

**If swallowed:** Drink one or two glasses of water and induce vomiting by touching back of the throat with finger. Repeat until vomit fluid is clear. Call a Physician immediately. Do not induce vomiting or give anything by mouth to an unconscious person.

**If inhaled:** Remove victim to fresh air. If not breathing, give artificial respiration, preferably mouth-to-mouth.

**If on skin:** Wash the affected area with soap and water.

**In case of eye contact:** Immediately flush eyes with fresh water for 15 minutes. Call a physician.

**Note to Physicians:** Emergency Information — call CHEMTREC (800) 424-9300.

**STORE IN A COOL, DRY PLACE.**

EPA Reg. No: 10107 - 100

EPA Est No: 10107 - NB1

**50 POUNDS NET**

Manufactured For:  
Cornbelt Chemical Co.  
P. O. Box 410  
McCook, Nebraska 69001

**DIRECTIONS (CON'T)**

**PRODUCT INFORMATION:** POTATO SEED TREATER M-Z WITH BARK is an agricultural Fungicide in a dust form containing Dithane®-45. IT IS USEFUL FOR TREATING Potato Seed Pieces and Potato Whole Seed for diseases listed.

**RECOMMENDATIONS**

POTATO SEED TREATER M-Z is a special fungicide dust formulation for application on whole or cut potato tubers to protect them from seed-piece decay caused by Fusarium sambucinum.

**DIRECTIONS FOR USE**

USE 1 LB. OF POTATO SEED TREATER M-Z per 100 lbs. of whole or cut potato seed pieces. apply the dust in a manner such that the entire surface of the seed piece becomes thoroughly covered with the fungicide dust. Plant seed as soon after treatment as possible. If planting is delayed more than 1-2 days, consult the seed piece supplier or local extension specialist for proper handling. Do not use treated seed pieces for food or feed purposes.

This product will not control bacterial diseases.

**CONDITIONS OF SALE:** 1. Cornbelt Chemical Company (Cornbelt) warrants that this material conforms to the chemical description on the label and is reasonably fit for use as directed hereon. Cornbelt neither makes, nor authorizes any agent or representative to make, any other warranty of FITNESS or of MERCHANTABILITY, guarantee or representation, express or implied, concerning this material.

2. Critical and unforeseeable factors beyond Cornbelt's control prevent it from eliminating all risks in connection with the use of chemicals. Such risks include, but are not limited to, damage to plants and crops to which the material is applied, lack of complete control, and damage caused by drift to other plants or crops. Such risks occur even though the product is reasonably fit for the uses stated hereon and even though label directions are followed. Buyer and user acknowledge and assume all risks and liability (except those assumed by Cornbelt under 1 above) resulting from handling, storage, and use of this material