

10F 1
Pm 25
10107-103

ACCEPTED

APR 2 0 1987

Under the Federal Insecticide,
Fungicide, and Rodenticide Act,
as amended, for the pesticide
registered under
EPA Reg. No. 10107-103

Precautionary Statements
Hazards To Humans
& Domestic Animals
CAUTION
 May cause irritation of eyes, nose, throat and skin. Avoid contact with skin, eyes or clothing. Wash thoroughly after handling.

ENVIRONMENTAL HAZARD
 Do not apply directly to any body of water. Do not contaminate water by cleaning of equipment or disposal of waste.

DIRECTIONS FOR USE
 It is a violation of federal law to use this product in a manner inconsistent with its labelling.

STORAGE AND DISPOSAL
PROHIBITIONS
 Do not contaminate water, food, or feed by storage, disposal or cleaning of equipment. Open dumping is prohibited. Avoid contamination of food or food-stuff.

PESTICIDE DISPOSAL
 Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL
 Completely empty bag by shaking and tapping sides and bottom to loosen clinging particles. Then dispose of bags in a sanitary landfill or by incineration if allowed by state and local authorities.

NOTICE OF WARRANTY
 Cornbelt warrants that this product conforms to the chemical description on the label thereof and is reasonably fit for the purposes stated on such label only when used in accordance with the directions under normal use conditions. It is impossible to eliminate all risks inherently associated with the use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or the manner of use or application, all of which are beyond the control of Cornbelt. In no case shall Cornbelt be liable for consequential, special or indirect damages resulting from the use or handling of this product. All purchasers shall be assumed to be buyers. CORNBELT MAKES NO WARRANTIES OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE NOR ANY OTHER EXPRESS OR IMPLIED WARRANTY EXCEPT AS SET FORTH ON THE LABEL.

HYKILL

BROMACIL 10G GRANULAR WEED KILLER

ACTIVE INGREDIENT:	Bromacil (5-bromo-3-sec-butyl-6-methyluracil)	30%
INERT INGREDIENTS		50%
	TOTAL	100%

Keep Out Of Reach of Children
CAUTION

STATEMENT OF PRACTICAL TREATMENT
 If in eyes: Flush with plenty of water. Get medical attention if irritation persists.
 If on skin: Wash with plenty of soap and water.

EPA Reg. No. 10107-
 EPA Est. No. 10107-NB1

NET CONTENTS, 50 LBS
 Mfg. For:
 Cornbelt Chemical Co.
 P.O. Box 410
 McCook, NE 68001

BEST AVAILABLE COPY

IMPORTANT

Loss of desirable trees or other plants may result from application. Do not apply to or near desirable trees or other plants or to areas where the chemical may extend, or in locations where the chemical may be carried by wind into contact with their roots. Do not use on tennis courts, tennis courts, or similar areas. Do not contaminate lawns, lawns, or other areas with fertilizers, insecticides, herbicides, or seeds. Thoroughly clean all traces of HYKILL BROMACIL 10G from application equipment immediately after use.

GENERAL INFORMATION
 HYKILL BROMACIL 10G is a granular product to be spread and applied as a dry material for control of weeds and brush. It is non-corrosive to equipment, non-flammable, and non-volatile. HYKILL BROMACIL 10G is an effective general herbicide for the control of many annual weeds at low rates and perennial weeds and brush at higher rates and is particularly useful for control of perennial grasses. It may be used on non-cropland for non-selective weed and brush control. Effects are slow to appear and may not become apparent until the chemical has been carried into the root zone of the weeds by moisture. The degree of control and duration of effect will vary with the amount of herbicide applied, soil texture, rainfall, and other conditions.

NON CROP USE WEED CONTROL
 HYKILL BROMACIL 10G should be applied as furnished with a seed spreader, a fertilizer spreader or with any equipment which will distribute the chemical uniformly over the area to be treated. For control of most weeds for an extended period of time on non-cropland areas such as Railroad, Highway and Pipeline Right-of-Ways, Petroleum Tank Farms, Lumberyards, Storage Areas and Industrial Plant Sites. Apply 24 to 48 lbs. per acre to control annuals such as cheat grass, downy brome, foxtail, lambsquarters, puncturing ragwort, kangaroo, turkey mullein and wild oats. When applied just before or after emergence of annuals, rates as low as 16 lbs. per acre will control many annual weeds and grasses in low rainfall areas and give short-term control in higher rainfall areas. Apply 16 to 36 lbs. per acre to control perennials such as aster, barnyard grass, bermudagrass, brome, sandstone, dog fennel, goldenrod, green, purple, nutgrass, fiddlefoot, quackgrass, smooth brome and wild carrot. In areas with low or seasonal rainfall, rates as low as 10 lbs. per acre control many perennial weeds and grasses. Apply 120 to 240 lbs. per acre to control bermudagrass, bouncingbet, bromefern, dallisgrass, dogbane, horseail, johnsongrass, nutsedge, sandgrass, and vasegrass. Where limited rainfall usually less than 4 inches occurs during the active growth period such as some areas of the West, HYKILL BROMACIL 10G usually will not provide sufficient control of hard-to-kill, deep rooted perennial weeds such as salt-tolerant grass.

NOTE: Use the higher levels of the dosage ranges on absorptive soils (those high in organic matter or carbon).
 Retreatment - Apply 16 to 48 lbs. per acre when annual weeds and grasses reappear on sites where weed growth has been controlled.
 For Small Areas - 1 cupful of HYKILL BROMACIL 10G per 250 sq. ft. is approximately 150 lbs. per acre.

BRUSH CONTROL
 To control undesirable woods, etc. on non-cropland areas such as Railroad Right-of-Ways, Storage Areas, Industrial Plant Sites and Drainage Ditches. Apply in spring or summer as a broadcast or spot treatment for use on drainage ditches.

Note
 For effective brush control and prevention of damage to desirable vegetation, do not apply to brush standing in water. Do not use water from treated ditches for irrigation. Do not use in irrigation ditches for on right-of-way or other sites where marketable timber or other desirable trees or shrubs are immediately adjacent to the treated area.

Broadest Treatment - Apply 56 to 96 lbs. per acre to control oak, pine, sweet gum, and willow. Apply 120 to 240 lbs. per acre to control brush such as American elm, cottonwood, hackberry, sumac, and winged elm. Use the higher levels of the dosage ranges on absorptive soils (those high in organic matter or carbon).
 Moisture is necessary to activate the chemical, best results are obtained if moisture is supplied by rainfall or irrigation within two weeks after application.

SELECTIVE USE IN CROPS

CITRUS (Oranges, Grapefruit, Lemons)
 Trees Established for Four Years or More
 Make a single application per year during the period from winter to early summer, use at the following rates:

Soil Texture	Lbs. HYKILL BROMACIL 10G Per Acre
Sand, loamy sand	32 to 40
Sandy loam	40 to 48
Silt loam, clay loam	48 to 64

Alternatively, make two applications of 24 to 32 lbs. HYKILL BROMACIL 10G per acre per year. In Florida, Texas and Louisiana, apply in spring and summer. In California and Arizona, apply in fall and spring.

Note: Partial control usually occurs with a single treatment. Repeat applications are required to control perennials. HYKILL BROMACIL 10G controls the following:

Bermudagrass	All areas U.S.
Torpedograss	Barrier Strip Treatment
Paraguaygrass	Florida
Johnsongrass	Texas
Nutsedge	Texas, California

Control of perennials may be improved by cultivation prior to treatment, other wise, avoid working the soil as long as weed control continues or else effectiveness of the treatment may be reduced.

Torpedograss Control - Barrier Strip Treatment
 For control of torpedograss adjacent to citrus groves to prevent spread of the weed into groves. Apply 240 lbs. per acre. Treat a border strip 10 to 20 ft. wide adjacent to the citrus grove, but no closer than 10 ft. to the trunk of citrus trees. Examine treated area every four months after application and spot treat invading or surviving torpedograss at 240 lbs. per acre (5 lbs. per 1000 sq. ft.). For best results apply in late winter or early spring after the torpedograss has broken dormancy and is actively growing. For later season application where growth is rank, mow or disk the area prior to treatment.