

NOTICE OF PESTICIDE: **REGISTRATION**
 REREGISTRATION
(Under the Federal Insecticide, Fungicide,
and Rodenticide Act, as amended)

conditional

NAME OF PESTICIDE PRODUCT

Herb Pro Atrazine 4L

NAME AND ADDRESS OF REGISTRANT (Include ZIP code)

F
Corneilt Chemical Company
P.O. Box 410
McCook, NE 69001

6-18-71

L
NOTE: Changes in labeling formula differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above U.S. EPA registration number.

On the basis of information furnished by the registrant, the above named pesticide is hereby Registered/Reregistered under the Federal Insecticide, Fungicide, and Rodenticide Act.

A copy of the labeling accepted in connection with this Registration/Reregistration is returned herewith.

Registration is in no way to be construed as an indorsement of approval of this product by this Agency. In order to protect health and the environment, the Administrator, on his own motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.

This product is conditionally registered in accordance with FIFRA sec. 3(c)(7)(A) provided that you:

1. Submit and/or cite all data required for registration/reregistration of your product under FIFRA sec. 3(c)(5) and sec. 4 when the Agency requires all registrants of similar products to submit such data.

2. Add the phrase, "EPA Registration No. 10107-RLG" and submit five (5) copies of your final printed labeling before you release the product for shipment.

3. Revise your labeling as follows:

1. Please check and correct labeling for typographical errors.
2. Under "General Information" page two (2), fourth paragraph; change sentence to read "This product is nonflammable."
3. Remove the following states under "Tank Mixtures for Chemical Fallow"; CO, MT, ND, SD, and WY.
4. Remove "Additional Tank Mixtures for Chemical Fallow".

BEST AVAILABLE COPY

ATTACHMENT IS APPROPRIATE

SIGNATURE OF APPROVING OFFICIAL

DATE

5. Under "Texas"; incorporate "For best results when weeds are emerged, add a nonionic surfactant at a concentration of 2qts./100 gals. to the spray and apply before weeds exceed 1.5 inches in height. Under precaution #3, change 30pts/A to 20 pts./A.
 6. Under "Conifers", drop the last four words of the second sentence and insert "...before weeds are 1.5 inches tall. Apply to established trees between fall and early spring while trees are dormant."
 7. Update your "Environmental Hazards" heading by replacing "Do not apply directly to water or wetlands" with "Do not apply to water, to areas where surface water is present or to intertidal areas below the mean high water mark.
- Replace "Do not contaminate water by cleaning of equipment or disposal of wastes" with "Do not contaminate water when disposing of equipment washwaters."
8. We are under the assumption that the following products are compatible:
 - a. Latron AB-44 and Unite
 - b. Cyclone and Gramoxone Extra
 - c. Sprayfuse 90 and X-77

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA sec. 6(e). Your release for shipment of the product constitutes acceptance of these conditions.

A stamped copy of the label is enclosed for your records.

Robert J. Taylor
 Product Manager (25)
 Fungicide-Herbicide Branch
 Registration Division (H7505C)

BEST AVAILABLE COPY

CONCURRENCES							
SYMBOL							
SURNAME						
DATE						

RESTRICTED USE PESTICIDE

(GROUND WATER CONCERN)

For retail sale to and use only by certified applicators or persons under their direct supervision and only for those uses covered by the certified applicator's certification.
This product is also a "Restricted Use" herbicide due to Atrazine ground water concerns. Users must read and follow all precautionary statements and instructions for use in order to minimize potential for Atrazine to reach ground water.

WEED PRO

Atrazine 4L

HERBICIDE

For season-long weed control in corn and sorghum. For weed control in certain other crops; in hempseed areas; and industrial sites.

ACTIVE INGREDIENTS

ATRAZINE (2-chloro-4-ethylamino-6-isopropylamino-2-triazine)

40.8%

Related Compounds

2.2%

INERT INGREDIENTS

57.0%

EPA Reg. No. 10107-

EPA Est. No. 10107-NB-1

Atrazine 4L contains 4 lbs. active ingredients per gallon.
Shake well before using. Use entire contents at one time.

KEEP OUT OF REACH OF CHILDREN

CAUTION

SEE ADDITIONAL PRECAUTIONARY STATEMENTS AND DIRECTIONS FOR USE INSIDE BOOKLET

Aerial application: Use aerial application only where broadcast applications are specified. Apply in a minimum of 1 qt. of water for each quart of the product applied per acre. For postemergence treatments on corn and sorghum, apply recommended rate in a minimum of 2 gals. of water/A. Avoid applications under conditions where uniform coverage cannot be obtained or where excessive spray drift may occur.

Avoid application to humans or animals: Flagmen and loaders should avoid inhalation of spray mist and prolonged contact with skin, and should wash thoroughly before eating and at the end of each day's operation.

Application in water or liquid fertilizer: Nitrogen solution or complete liquid fertilizer may replace all or part of the water as a carrier for preemergence, preplant incorporated, or preplant surface ground application on corn and sorghum. Check the compatibility of this product with liquid fertilizer and/or nitrogen solution as shown below before use. Do not apply in nitrogen solution or complete liquid fertilizer after corn or sorghum emerges or crop injury may occur.

Compatibility Test: Since liquid fertilizers can vary, even within the same analysis, always check compatibility with herbicide(s) each time before use. Be especially careful when using complete suspension or fluid fertilizers as serious compatibility problems are more likely to occur. Commercial application equipment may improve compatibility in some instances. The following test assumes a spray volume of 25 gals. per acre. For other spray volumes, make appropriate changes in the ingredients. Check compatibility using this procedure:

1. Add 1 pt. of fertilizer to each of 2 one-quart jars with tight lids.
2. To one of the jars, add 1/4 tsp. or 1.2 milliliters of a compatibility agent approved for this use, such as Compex® or Unite® (1/4 tsp. is equivalent to 2 pts. per 100 gals. spray). Shake or stir gently to mix.
3. To both jars add the appropriate amount of herbicide(s). If more than one herbicide is used, add them separately with dry herbicides first, flowable next, and emulsifiable concentrates last. After each addition, shake or stir gently to thoroughly mix. The appropriate amount of herbicides for this test follows:

Dry herbicides: For each pound to be applied per acre, add 1.5 level

DIRECTIONS FOR USE AND CONDITIONS OF SALE AND WARRANTY

IMPORTANT: Read the entire Directions for Use and the Conditions of Sale and Warranty before using this product. If terms are not acceptable, return the unopened product container at once.

Conditions of Sale and Warranty

The Directions for Use of this product reflect the opinion of experts based on field use and tests. The directions are believed to be reliable and should be followed carefully. However, it is impossible to eliminate all risks inherently associated with use of this product. Crop injury, ineffectiveness, or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or the manner of use or application all of which are beyond the control of CORNBELT CHEMICAL CO. or the Seller. All such risks shall be assumed by the Buyer.

CORNBELT CHEMICAL CO. warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes referred to in the Directions for Use subject to the inherent risks referred to above. CORNBELT CHEMICAL CO. makes no other express or implied warranty of Fitness or Merchantability or any other express or implied warranty. In no case shall CORNBELT CHEMICAL CO. or the Seller be liable for consequential, special, or indirect damages resulting from the use or handling of this product. CORNBELT CHEMICAL CO. and the Seller offer this product, and the Buyer and use accept it, subject to the foregoing Conditions of Sale and Warranty, which may be varied only by agreement in writing signed by a duly authorized representative of CORNBELT CHEMICAL CO.

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

FAILURE TO FOLLOW THE DIRECTIONS FOR USE AND PRECAUTIONS ON THIS LABEL MAY RESULT IN POOR WEED CONTROL, CROP INJURY, OR ILLEGAL RESIDUES.

Do not apply this product in such a manner as to directly or through drift expose workers or other persons, except those knowingly involved in the application. The area being treated must be vacated by unprotected persons.

Do not enter treated areas without protective clothing until sprays have dried. Because certain states may require more restrictive reentry intervals for various

1

teaspoons to each jar.

Liquid herbicides: For each pint to be applied per acre, add 0.5 teaspoon or 2.5 milliliters to each jar.

4. After adding all ingredients, put lids on and tighten, and invert each jar ten times to mix. Let the mixture stand 15 minutes and then look for separation, large flakes, precipitates, gels, heavy oily film on the jar, or other signs of incompatibility. Determine if the compatibility agent is needed in the spray mixture by comparing the two jars. If either mixture separates, but can be remixed readily, the mixture can be sprayed as long as good agitation is used. If the mixtures are incompatible, test the following methods of improving compatibility: (A) slurry the dry herbicide(s) in water before addition, or (B) add 1/2 of the compatibility agent to the fertilizer and the other 1/2 to the emulsifiable concentrate or flowable herbicide before addition to the mixture. If incompatibility is still observed, do not use the mixture.

Application in water plus emulsifiable oil or oil concentrate: Adding emulsifiable oil (petroleum-derived, petroleum-derived oil concentrate, or single or mixed crop-derived oil concentrate) to postemergence water-based sprays in corn and sorghum may improve weed control. However, under certain conditions, the use of oil or type of oil may seriously injure the crop. To minimize this possibility, observe the following directions:

Use one of the following properly emulsified:

1. A suitable oil concentrate containing at least 1% but not more than 20% suitable emulsifier or surfactant blend.

2. Petroleum-derived oil containing at least 1% suitable emulsifier.

Note: In the event of a compatibility problem when mixing oil with this product and water, a compatibility agent such as Compex or Latron AG-44 should be used. Any of the above oils contaminated with water or other materials can cause compatibility problems and/or crop injury.

Mixing procedures - all uses: (1) Be sure sprayer is clean and not contaminated with any other materials, or crop injury or sprayer clogging may result. (2) Fill tank 1/4 full with clean water, nitrogen solution, or complete liquid fertilizer. (3) Start agitation. (4) Be certain that the agitation system is working properly and creates a rippling or rolling action on the liquid surface. (5) Pour product

c. related with this product, consult your State Department of Agriculture.

Written or oral warnings must be given to workers who will be exposed to this product in the treated area or in an area about to be treated with this product. These warnings must be given which inform workers of areas or fields where specific protective clothing until sprays have dried. Workers must take actions to take in case of accidental exposure as described in the Statements on this label. When oral warnings are given in a language customarily understood by workers, written warnings must be given in English. Written warnings must include the following information:
STATEMENT: Area treated with Weed Pro Atrazine 4L on (date) enter without appropriate protective clothing until sprays have dried. If accidental exposure, flush eyes or skin with plenty of water. If irritation persists. Remove and wash contaminated clothing. Do not apply this product through any type of irrigation system.

General Information

This herbicide controls many annual broadleaf and grass weeds, including barnyardgrass, johnsongrass, sugarcane, and certain other crops specifically effective in noncrop areas and industrial sites for controlling many perennial broadleaf and grass weeds. This product may be applied to the soil surface.

Following many years of continuous use of this product, biotypes of some of the weeds listed on this label which cannot be effectively controlled by this and related products are known or suspected, and weeds controlled by this product along with resistant biotypes, we recommend combinations or in sequence with other registered herbicides. If only resistant biotypes are expected to be present, use the most effective herbicide. Consult with your state Agricultural Experiment Station for recommendations.

ACCEPTABLE USES

with COMMENTS

In EPA Letter dated [REDACTED] this product acts mainly through root absorption. It depends on moisture to move it into the root zone. If water is not available, cultivation or rotary hosing will generally result in better control.

This product is noncorrosive to equipment and metal and has low electrical conductivity.

Under the Federal Insecticide, Fungicide, and Rodenticide Act as amended, for the pesticide registered under EPA Reg. No.

10107-153
directly from container into tank. (6) Continue filling tank if necessary to maintain surface action. (7) Concentrate, or tank mix herbicide(s) after this product. (8) Finish filling tank. (9) Empty tank as completely as possible to prevent buildup of oil or emulsifiable concentrate residue. (10) If an oil or emulsifiable concentrate starts to build up in tank, drain it and clean with solvent. (11) Clean sprayer thoroughly immediately after use with water containing a detergent.

Atrazine 4L Applied Alone - Corn or Grain Sorghum

Precipitated Surface-Applied, Precipitant Incorporated Postemergence at 4 pts./A With Oil

Broadleaf and Grass Weed Controlled

barnyardgrass (watergrass)	cocklebur
grass, foxtail*	groundcherry
green foxtail	jmponeeed
large (hairy) crabgrass**	kochia
wild oats	lambsquarters
witchgrass (<i>Panicum capillare</i>)	annual morningglory
yellow foxtail	mustards

Postemergence with Emulsifiable Oil or Oil Concentrate

annual morningglory	lambsquarters	ragweed
cocklebur	mustards	smartweed
jmponeeed	pigweed	velvetleaf

* Certain states may have established rate limitations in certain areas. Consult your state lead pesticide control information. It is a violation of this label to deviate in rates, for example, maximum use rates have been established for specific pesticide management areas. Contact the Iowa Department of Agriculture and Land Stewardship, CORNBELT CHEMICAL CO. Representative for additional information.

** Indicates partial control only.

Broadleaf control (eastern CO, western KS, western NE, NM, OK Panhandle, west TX, and eastern WY): On sand, loamy sand, sandy loam, mild to strongly alkaline soil, and all recently tilled soil, apply no more than 2.3 pts./A, either preplant surface, preplant incorporated, or preemergence. On other soils in these areas, apply rate in Table 1 for broadleaf and grass control.

Certain states may have established rate limitations within specific geographical areas. Consult your state lead pesticide control agency for additional information. It is a violation of this label to deviate from state use regulations. In Iowa, for example, maximum use rates have been established within a specified pesticide management area. Contact the Iowa Extension Service, the Iowa Department of Agriculture and Land Stewardship or your CORNBELT CHEMICAL CO. representative for additional information.

Postemergence with emulsifiable oil or oil concentrate in water: Add the following volume of one of the type oils indicated for aerial or ground application unless the oil label specifies otherwise:

Type Oil	Ground Application	Aerial Application
Oil concentrate (Crop or Petroleum-derived)	1 qt/A	1/2-1 qt/A
Petroleum-derived oil	1 gal/A	2 qts/A

Note: Crop-derived or petroleum-derived oil concentrates sr. 14 contain at least 1% but not more than 20% suitable emulsifier or surfactant blend. Petroleum-derived oils should contain at least 1% suitable emulsifier.

Broadleaf and grass control: For postemergence control of those weeds listed under Preplant Incorporated and Preemergence, broadcast 4 pts./A plus emulsifiable oil or oil concentrate after weed emergence, but before weeds reach 1.5 inches in height and before corn exceeds 12 inches in height.

Broadleaf control: For postemergence control of those weeds listed under Postemergence with emulsifiable oil or oil concentrate in water, broadcast 2.4 pts./A plus emulsifiable oil or oil concentrate before pigweed, lambquarters reach 6 inches in height and before all other weeds reach 4 inches in height. A cultivation may be necessary if all weeds are not controlled or if weeds regrow.

Precautions: For applications with emulsifiable oil or oil concentrate in water: (1) Infested areas or any breeding stock may be severely injured by applications with emulsifiable oil or oil concentrate. (2) Adding other insecticides, herbicides, liquid fertilizers, or other materials is not recommended, because they may cause compatibility problems or crop injury. (3) Store and handle emulsifiable

oil and oil concentrate carefully. Oil contaminated with even a small amount of water may not emulsify properly when added to the tank. To avoid crop injury, (4) Do not apply when crop is under stress from prolonged cold, heat, weather, poor fertility, or other factors, or when crop is wet and succulent from recent rainfall. (5) Do not exceed 3 lbs. active ingredients (or 6 pts. of this product) per acre per calendar year. (6) Postemergence applications to corn must be made before corn exceeds 12 inches in height. (7) Do not make more than one application per season except as recommended for partial control of yellow nutsedge, quackgrass, and Canada thistle.

Problem weeds suppression

Yellow nutsedge and Canada thistle: This product will partially control (up to) yellow nutsedge (*Cyperus esculentus*) and Canada thistle (*Cirsium arvense*) when applied according to directions. For best results, apply each year until yellow nutsedge or Canada thistle is eliminated or reaches a level of infestation where neither weed is an economic problem. If yellow nutsedge or Canada thistle regrow following last application, cultivate once.

Four alternative methods of use for partial control (suppression) of yellow nutsedge and Canada thistle are listed below in order of preference. If other weed species, such as annual grasses, are also expected, use alternative 2 or 4.

1. Broadcast 3 pts./A plus 1 gallon of emulsifiable oil or 1 qt. of oil concentrate. Apply after crop and yellow nutsedge or Canada thistle emerge, but before yellow nutsedge reaches a height of 3 inches or Canada thistle reaches a height of 5 inches. Repeat application before corn exceeds 12 inches in height.

2. Broadcast 3 pts./A preplant incorporated for control of yellow nutsedge, or broadcast 3 pts./A during or shortly after planting, but prior to crop or weed emergence for control of Canada thistle. Follow with an application of 3 pts./A plus 1 gallon of emulsifiable oil or 1 qt. of oil concentrate after corn and weeds emerge, but before yellow nutsedge or Canada thistle reaches a height of 6 inches and before corn exceeds 12 inches in height.

3. Broadcast 6 pts./A plus 1 gallon of emulsifiable oil or 1 qt. of oil concentrate after crop emerges, but before corn exceeds 12 inches in height. Apply after yellow nutsedge and Canada thistle emerge, but before yellow nutsedge is 3 inches tall or Canada thistle is 6 inches tall.

4. Broadcast 6 pts./A before, during, or shortly after planting, but before crop or weed emergence. (yellow nutsedge control only).

9

It is a violation of this label to not use emulsifiable oil or oil concentrate when corn is wet or under stress, especially when using 6 pts./A of this product, as crop injury may occur. See "Precautions for applications with emulsifiable oil or oil concentrate in water" for additional directions.

Quackgrass suppression on land going into corn production

Split application: Broadcast 3 pts./A in early spring. Plow 1-3 weeks later. Apply a second application at 3 pts./A in spring before, during, or after planting, but before weeds are 1.5 inches high and before corn exceeds 12 inches in height. This split application will suppress quackgrass and control most annual broadleaf and grass weeds.

Quackgrass control on land going into corn production

Split application: Make a single broadcast application of 6 pts./A in spring before planting. Plow 1-3 weeks later. Follow-up in the fall by utilizing a suitable contact herbicide, such as Roundup.

Tank Mixtures for Corn

This product may be tank mixed with these herbicides for control of certain broadleaf and grass weeds in corn:

Dual® (isoproturon)	Lasso or Lasso EC + Roundup
Dual + Gramoxone Extra	Lasso or Lasso EC + Gramoxone Extra
Dual + Roundup (Glyphosate)	Gramoxone Extra
Dual + Princep®	Princep
Dual + Princep + Gramoxone Extra	Princep + Gramoxone Extra
Dual + Princep + Roundup	Princep + Roundup
Benton® or RawFlex® (propachlor)	Roundup
Lasso® or Lasso EC (isoproturon)	Susan™

Use tank mix directions appearing on the labels of the above herbicides when tank mixing with this product. Observe all precautions and limitations on labeling of product used in a particular tank mix.

Note: When the labels of the above herbicides refer to atrazine 80W, use

10

Table 2: Tank Mixtures with Princep on Corn

Soil Texture	Broadcast rate/A		
	1:1 Rate ^a	1:2 Rate ^b	1:3 Rate ^c
Sand, loamy sand, loam	This Princep 80W	This Princep 80W	1.32 pts.
2 pts.	1.25 lbs.	1.32 lbs.	1.67 lbs.
Loam, silt loam, silty clay loam, sandy clay loam, silty day loam, sandy clay, or silty clay with low organic matter	2.4 pts.	1.5 lbs.	1.6 pts.
2 lbs.			
Loam, silt loam, silty clay loam, sandy clay loam, silty day loam, sandy clay, or silty clay with medium to high organic matter, and clay including dark prairie soils of the Corn Belt	3 pts.	1.3 lbs.	1.92 pts.
			2.4 lbs.

^a For control of most weeds

^b For control of especially heavy infestations of crabgrass and fall panicum.

^c When using Princep Caliber 50 or Princep 4L, use equivalent rates. One lb. of Princep 80W equates 0.8 lbs. of Princep Caliber 50 or 1.6 pts. of Princep 4L.

Princep 80W, Princep 4L, or Princep Caliber 50 plus Roundup: Use as tank mixture for preemergence and postemergence control of certain broadleaf and grass weeds where corn will be planted directly into a cover crop, established seed, or in previous crop residues. Refer to Roundup label for all directions, weeds controlled, precautions, and limitations.

Princep 80W, Princep 4L, or Princep Caliber 50 plus Gramoxone Extra: Use as tank mixture with Princep and Gramoxone Extra to kill existing vegetation and for residual weed control where corn will be planted directly into a cover crop, established seed, or in previous crop residues. Add this product and Princep to water in spray tank, agitating until thoroughly mixed. Then add Gramoxone Extra and a nonionic surfactant, such as X-77. Continue agitation during application. Broadcast 2-4 pts. of this product plus 1.25-2.5 lbs. of Princep 80W (or 2-4 pts. of Princep 4L, or 1.1-2.2 lbs. of Princep Caliber 50) plus a suitable amount of Gramoxone Extra in 20-60 pts. of water per sprayed acre. Refer to Gramoxone Extra label for the appropriate rates to utilize in this tank mixture. Apply before, during, or after planting, but before corn emerges. Add 0.5 pt. of a nonionic surfactant, such as X-77, per 100 pts. of spray mixture. Use the higher rate of Gramoxone Extra specified on the label if existing vegetation is 4-6 inches tall. This mixture will not control weeds taller than 6 inches.

12

Refer to further limitations and precautions on labels for this product, Princep, and Gramoxone Extra.

Precautions for all applications to corn: (1) To avoid crop injury and illegal residues, do not apply more than 3 lbs./A active ingredient (6 pts./A of the product) per calendar year. (2) For best control of velvetleaf and cocklebur, the application rate cannot be less than 2 lbs./A active ingredients, either alone or in tank mix combinations. (3) Following harvest, plow (moldboard or disk-plow) and thoroughly till soil in fall or spring to minimize possible injury to spring-seeded rotational crops, regardless of rate used.

Note for all applications to corn: Do not graze or feed forage from treated areas for 21 days following application, or illegal residues may result.

Rotational crops: (1) Do not rotate to any crop except corn or sorghum until the following year, or injury may occur. (2) If applied after June 10, do not rotate with crops other than corn or sorghum the next year, or crop injury may occur. (3) In the High Plains and Intermountain areas of the West where rainfall is sparse and erratic or where irrigation is required, use only when corn or sorghum is to follow corn or sorghum or when a crop of untreated corn or sorghum is to precede other rotational crops. (4) In eastern parts of the Dakotas, KS, western MN, and NE, do not rotate to soybeans if the rate applied to corn or sorghum was more than 4 pts./A or equivalent band application rate, or soybean injury may occur. (5) Injury may occur to soybeans planted the year following application on soils having a calcareous surface layer. (6) Do not plant sugar beets, tobacco, vegetables (including dry beans), spring-seeded small grains, or small-seeded legumes and grasses the year following application, or injury may occur.

Sorghum and Sorghum-sudan Hybrids (Grain and Forage Types)

Preplant Surface-Applied (Broadleaf and grass control): Use on medium- and fine-textured soil with minimum-tillage or no-till systems only in CO, IL, IN, IA, KS, KY, MN, MO, MT, NE, ND, SD, WI, and WY. Apply 2/3 the recommended rate of WeedsPro Atrazine 4L shown in Table 3 for the soil texture as a split treatment 30-45 days prior to planting and the remainder at planting. Applications made less than 30 days prior to planting may be as either a split or single treatment. Under dry conditions, irrigation after application is recommended to move WeedsPro Atrazine 4L into the soil.

If weeds are present at time of treatment, apply in a tank mix combination with a contact herbicide (for example, Gramoxone Extra or Roundup). Observe 13

it is a violation of this label to not use emulsifiable oil or oil concentrate when corn is wet or under stress, especially when using 6 pts./A of this product, as crop injury may occur. See "Precautions for applications with emulsifiable oil or oil concentrate in water" for additional directions.

Quackgrass suppression on land going into corn production

Split application: Broadcast 3 pts./A in early spring. Plow 1-3 weeks later. Apply a second application at 3 pts./A in spring before, during, or after planting, but before weeds are 1.5 inches high and before corn exceeds 12 inches in height. This split application will suppress quackgrass and control most annual broadleaf and grass weeds.

Quackgrass control on land going into corn production

Split application: Make a single broadcast application of 6 pts./A in spring before planting. Plow 1-3 weeks later. Follow-up in the fall by utilizing a suitable contact herbicide, such as Roundup.

Tank Mixtures for Corn

This product may be tank mixed with these herbicides for control of certain broadleaf and grass weeds in corn:

Dual® (isoproturon)	Lasso or Lasso EC + Roundup
Dual + Gramoxone Extra	Lasso or Lasso EC + Gramoxone Extra
Dual + Roundup (Glyphosate)	Gramoxone Extra
Dual + Princep®	Princep
Dual + Princep + Gramoxone Extra	Princep + Gramoxone Extra
Dual + Princep + Roundup	Princep + Roundup
Benton® or RawFlex® (propachlor)	Roundup
Lasso® or Lasso EC (isoproturon)	Susan™

Use tank mix directions appearing on the labels of the above herbicides when tank mixing with this product. Observe all precautions and limitations on labeling of product used in a particular tank mix.

Note: When the labels of the above herbicides refer to atrazine 80W, use

11

equivalent rates of WeedsPro Atrazine 4L.

Principle 80W, Roundup

In addition to this product, Cenix or Grain Sengrae rated, or Preemergence corn, and carpetweed. Broadcast tank mix helps and weeds emerge, all weeds. Low the 12-hp panicum. Cultivate soil.

Preplant Surface-Applied minimum-tillage or no-till MT, NE, ND, SD, WI, an Atrazine 4L and Princep 30-45 days prior to planting. On coarse-textured soils. Refer to the Weedseed A of the corn label for addi-

If weeds are present at a contact herbicide (for direction for use, precal herbicide).

Note: To the extent possi- untreated soil to the surfa

Preplant Incorporated: during, or after final seed results, apply within two weeks.

Preemergence: Apply weeds emerge.

Refer to Corn sections of Princep 4L labels for furt

11

Preemergence broadleaf ghem (AZ and CA only): broadcast 1.6-2.4 pts./A before sorghum and weed with several regular irriga

Preplant Incorporated (Broadleaf and grass control): Broadcast in spring after tilling at rate shown in Table 3. Apply before, during, or after final seed or preparation. If soil is tilled or worked after application, avoid deep incorporation. For best results, apply within two weeks prior to planting.

Preemergence (Broadleaf and grass control): Apply during or shortly after planting, but prior to weed or crop emergence at rate shown in Table 3.

Table 3: Preplant Surface, Preplant Incorporated, and Preemergence Broadleaf and Grass Weed Control in Sorghum

Soil Texture	Organic Matter	Broadcast rate/A
COARSE	any level	DO NOT USE (except preemergence use on treated sorghum in AZ and CA)
Sand, loamy sand, sandy loam	less than 1%	3.2-4 pts. ^a
MEDIUM AND FINE	1-1.5%	
Loam, silt loam, silty clay loam, sandy clay loam, silty clay loam, sandy clay, silty clay, clay	more than 1.5%	4-4.75 pts. ^a

^a Do not apply preplant surface or preplant incorporated in AL, AR, FL, GA, LA, MS, NC, NM, OK, SC, TN, or TX. Do not apply preemergence in NM, OK, or TX, except in northeast OK and the TX Gulf Coast.

Certain states may have established rate limitations within specific geographical areas. Consult your state lead pesticide control agency for additional information. It is a violation of this label to deviate from state use regulations. In Iowa, for example, maximum use rates have been established within a specified pesticide management area. Contact the Iowa Extension Service, the Iowa Department of Agriculture and Land Stewardship, or your CORNBELT CHEMICAL CO. representative for additional information.

In case of planting failure, sorghum or corn may be replanted. Do not make a second broadcast application, or injury may occur. If originally applied in a band and sorghum or corn is replanted in unseeded row residue, this product may be applied in a band to the second planting.

14

Preemergence broadleaf control in water: Broadcast

Apply before pigweed and

all other weeds reach 4 inches

in the regions of AZ and CA

15

before it reaches boot stage. In all other areas, apply after sorghum reaches the 3-leaf stage but before it exceeds 12 inches in height. Add 1 qt./A of emulsifiable oil/A for ground application and 0.5 gal./A for aerial application, or add 1 qt./A of oil concentrate for ground application. A cultivation may be necessary if all weeds are not controlled or if weeds regrow.

For the list of weeds controlled, see **Weedpro Alatazine 4L Applied Alone - Corn or Grain Sorghum - Postemergence with Emulsifiable Oil or Oil Concentrate in Water**.

Precautions for applications with emulsifiable oil or oil concentrate in water: See "Precautions for applications with emulsifiable oil or oil concentrate in water" in Corn section.

Post-emergence broadleaf weed control with surfactant (CO, Western KS, NM, OK, TX, and desert regions of AZ and CA only): Broadcast 2.4 pts./A plus 0.75-1.5 pt. of surfactant after sorghum reaches 6 inches in height, but before weeds exceed 1.5 inches in height. Apply only on sandy loam and finer textured soil.

Precautions for all applications to sorghum: (1) Heavy rain immediately following application tends to cause excessive concentrations of herbicide in seed furrow, resulting in possible crop injury. Do not apply to furrow-planted sorghum until furrows are leveled (plowed in). Level deep planter marks or seed furrows before application. (2) Application to sorghum growing under stress caused by minor element deficiency or to sorghum growing on highly calcareous soil may result in crop injury. (3) Following harvest, plow (rototiller or disk-till) and thoroughly till soil in fall or spring to minimize possible injury to spring-seeded rotational crops, regardless of rate used. (4) Injury may occur if both this herbicide, preplant surface, preplant incorporated, or preemergence, and an atrazine systemic insecticide applied in-furrow are used. (5) Do not apply more than 3 lbs. active ingredient (6 pts. of this product) per acre per calendar year.

Note: Do not graze or feed forage from treated areas for 21 days following applications, or illegal residues may result.

Rotational crops: See "Rotational crops" in Corn section.

Tank Mixtures for Grain Sorghum

Dual 8E: Use as tank mixture with Dual 8E for control of those weeds listed on the Dual 8E label, as well as on this label. Use this tank mixture only on sorghum seed treated with Concep®. Refer to the Dual 8E label for all directions, precautions, and limitations.

15

Winter Weed Control in Texas

For postemergence control of winter weeds only, such as henbit, seedling dock, and annual thistle on fall bedded land in the Gulf Coast and Blacklands of Texas. Apply 0.8-1 qt./A postemergence to the weeds in November or December to land that will be planted to corn, grain sorghum, or forage sorghum the following spring. For best results, add a suitable surfactant, such as X-77, at the rate of 0.5% of the spray volume, an emulsifiable oil at the rate of 1.0% of the spray volume, or an oil concentrate at the rate of 1 qt. per acre.

Normal weed control programs may be used in the following corn, grain sorghum or forage sorghum crop.

Note: Do not plant any crops except corn, grain sorghum, or forage sorghum the spring following this treatment, or illegal residues may result.

Weedpro Alatazine 4L Alone - Chemical Fallow

Wheat-Sorghum-Fallow: To control annual broadleaf and grass weeds following wheat harvest and in the following sorghum crop when grown under minimum tillage, broadcast 6 pts./A to wheat stubble immediately following wheat harvest. If weeds are present at application, remove them with a sweep plow or other suitable implement after application, or use an approved contact herbicide before or after the application of Weedpro Alatazine 4L. Plant sorghum into wheat stubble the following spring with minimum soil disturbance. Use a surface planter or a planter leaving a shallow furrow. If weeds are present at planting, remove them with a sweep plow or other suitable implement before planting.

For the list of weeds controlled, see **Weedpro Alatazine 4L Applied Alone - Corn or Grain Sorghum - Preplant Surface-Applied, Preplant Incorporated, or Preemergence**.

Precautions: (1) Use only on silt loam or finer textured soil or crop injury may result. (2) Wheat-sorghum-fallow cropping sequence must be followed. (3) Do not apply following sorghum harvest.

Note: To avoid illegal residues, do not graze or feed forage from treated area to livestock. To avoid illegal residues and crop injury, do not plant any crop other than those on the label within 18 months following treatment.

Wheat-Corn-Fallow (KS, NE): This product controls cheatgrass (downy brome, chess), Kochia, mustard, pigweed, Russian thistle, wild lettuce, wild sunflower, and volunteer wheat during period after wheat harvest. Weed control may

17

be suppressed at the 20 ounces per acre rate and will require up to 26 ounces per acre for control. The addition of Ammonium Sulfate is recommended to increase the performance of Roundup RT and Weedpro Alatazine 4L plus a nonionic surfactant such as Sprayfuse 90 in a tank mix. For ground applications, apply in 3 to 10 gallons of water/A. For aerial applications, apply in 3 to 5 gallons of water/A. Refer to the Roundup RT label for further directions, precautions, and limitations.

Roundup RT (CO, KS, MT, NE, ND, SD, and WY): Follow directions for use, notes, and precautions in the "Wheat-Sorghum-Fallow" section above, substituting corn for references to sorghum.

Wheat-Fallow-Wheat (CO, KS, MT, NE, ND, SD, and WY): For preemergence control of cheatgrass (downy brome, chess), common lambsquarters, field pennycress, Kochia, mustard, Russian thistle, wild lettuce, and suppression of volunteer wheat during fallow period of a wheat-fallow-wheat rotation, broadcast 1-2 pts./A on all soils except those listed under "Precautions." For control of pigweed and wild sunflower, use the higher rate. Apply to stubble ground. Treat only once during same fallow period.

Tank Mixtures for Chemical Fallow

Wheat-Sorghum-Fallow or Wheat-Corn-Fallow (CO, KS, MT, ND, NE, SD, WY)

Cyclone/Gramoxone Extra: If weeds are present at application, a tank mix with Cyclone/Gramoxone Extra may be used. Broadcast 6 pts. of Weedpro Alatazine 4L plus a suitable amount of Cyclone/Gramoxone Extra in 20-60 gals. of water/A by ground equipment. Add 0.5-1 pt. of a nonionic surfactant, such as SPRAYFUSE 90, per 100 gals. of spray mixture. Add Weedpro Alatazine 4L to spray tank first and thoroughly mix with water. Then add Cyclone/Gramoxone Extra, followed by surfactant. Use the higher rate of Cyclone/Gramoxone Extra specified on the label if weeds are 4-6 inches tall. This mixture will not control weeds taller than 5 inches. Apply to stubble ground. Treat only once during same fallow period. Refer to Gramoxone Extra/Cyclone label for further directions, precautions, and limitations.

Wheat-Fallow-Wheat (CO, KS, MT, NE, ND, SD, and WY):

Cyclone/Gramoxone Extra: If weeds are present at application, a tank mix with Cyclone/Gramoxone Extra may be used. Broadcast 1-2 pts. of Weedpro Alatazine 4L plus a suitable amount of Cyclone/Gramoxone in 20-60 gals. of water/A by ground equipment. Add 0.5-1 pt. of a nonionic surfactant, such as SPRAYFUSE 90, per 100 gals. of spray mixture. Add Weedpro Alatazine 4L to spray tank first and thoroughly mix with water. Then add Cyclone/Gramoxone Extra, followed by surfactant. Use the higher rate of Cyclone/Gramoxone Extra specified on the label if weeds are 4-6 inches tall. This mixture will not control weeds taller than 5 inches. Apply to stubble ground. Treat only once during same fallow period. Refer to Cyclone/Gramoxone Extra label for further directions, precautions, and limitations.

If weeds are present at application and this product is used alone, use either an approved contact herbicide before or after treatment, or tillage after treatment.

18

be suppressed at the 20 ounces per acre rate and will require up to 26 ounces per acre for control. The addition of Ammonium Sulfate is recommended to increase the performance of Roundup RT and Weedpro Alatazine 4L plus a nonionic surfactant such as Sprayfuse 90 in a tank mix. For ground applications, apply in 3 to 10 gallons of water/A. For aerial applications, apply in 3 to 5 gallons of water/A. Refer to the Roundup RT label for further directions, precautions, and limitations.

Landmaster BW: (All states except AL, FL, HI, AK, SC, ME, NH, VT, RI, CA, GA, MD)

If weeds are present at application a tank mix with Landmaster BW may be used. Apply 40 ounces of Landmaster BW plus 2 pts. or less of Weedpro Alatazine 4L/A. Use 54 ounces Landmaster BW plus 4 pts. or less of Weedpro Alatazine 4L/A. Use 64 ounces of Landmaster BW plus 8 pts. or less of Weedpro Alatazine 4L. The addition of Ammonium Sulfate is recommended to increase the performance of Landmaster BW and Weedpro Alatazine 4L. In Oregon and Washington, do not exceed 2 pts. of Weedpro Alatazine 4L per acre. For ground applications, apply in 3 to 10 gallons of water per acre. For aerial applications, apply in 3 to 5 gallons of water per acre. Refer to the Landmaster BW label for further directions, precautions, and limitations.

Precautions: To avoid crop injury, (1) Do not use on sand soil. (2) Do not treat eroded hillsides, cutbank and rocky outcroppings, or exposed calcareous subsoil. (3) Do not treat soils of the Rosedale and Canyon Series in western NE and adjoining counties in CO and WY. (4) Do not treat soils with calcareous surface layers. (5) Avoid spray overlap.

Note: Do not graze treated areas within 6 months after application, or illegal residues may result.

Aerial application: In order to ensure that spray will be controllable within the target area when used according to label directions, make applications at a maximum height of 10 ft., using low drift nozzles at a maximum pressure of 40 psi, and restrict application to periods when wind speed does not exceed 10 mph. To ensure that spray will not adversely affect adjacent sensitive nontarget plants, apply Weedpro Alatazine 4L alone by aircraft at a minimum upwind distance of 400 ft. from sensitive plants.

Residues: To control certain annual weeds in established perennial grasses along road-

20

sides in CO, KS, MT, NE, ND, SD, and WY, including cheatgrass (downy brome, chess), common (annual) broomweed, little barley, medusahead, sagebrush, and tumble mustard, broadcast 2 pts./A in a minimum of 10 gals. of water by ground equipment in the fall before ground freezes, or after thawing in the spring, but before the established grasses green-up and before weeds emerge. Examples of desirable established grasses include big bluestem, bluegrass, bromegrass, buffalograss, crested wheatgrass, Indiangrass, little bluestem, side-oats grama, switchgrass, and Western wheatgrass. Apply only once per year. Temporary decoloration or other form of injury to the desirable perennial grasses may occur following application.

Note: To avoid illegal residues, (1) Do not cut or feed roadside grass hay. (2) Do not allow livestock to graze treated areas.

Sugarcane

For control of many broadleaf and grass weeds, including amaranthe, crabgrass, hawksbeard, Florida's paintbrush, foxtail, junciperus and wiregrass, broadcast 5-8 pts./A of Weedpro Alatazine 4L at time of planting or retanning, but before sugarcane emerges. Broadcast serially in a minimum of 5 gals. of spray per acre, or broadcast or band by ground equipment in a minimum of 20 gals. per acre, unless indicated otherwise. One additional application may be made over the sugarcane as it emerges, and two additional applications may be made later after emergence as directed sprays. Repeat treatments, where needed, may be applied broadcast, band, or interline as suggested with the final application being prior to close-in. Do not exceed the rate of herbicide suggested for any one crop of sugarcane.

Note: Where high rates of Weedpro Alatazine 4L are used alone, apply in a minimum of 1 qt. of water for each 1 qt. of Weedpro Alatazine 4L applied per acre.

Aerial application: In order to ensure that spray will be controllable within the target area when used according to label directions, make applications at a maximum height of 10 ft., using low drift nozzles at a maximum pressure of 40 psi, and restrict application to periods when wind speed does not exceed 10 mph. To ensure that spray will not adversely affect adjacent sensitive nontarget plants, apply Weedpro Alatazine 4L alone by aircraft at a minimum upwind distance of 400 ft. from sensitive plants.

Florida

For control of emerged palmetto weed: Apply 0.8-1.2 pts./A in at least 40 gals. of water as a directed spray by ground equipment prior to close-in. Add 4 qt. of surfactant for each 100 gals. of spray. Thoroughly cover weed foliage. For control of amaranth, large crabgrass, palmetto (fertilizer) weed, and spiny amaranth, see one of the following methods at planting or retanning:

21

1. Apply 8 pts./A preemergence. Follow with one or two applications, as needed, postemergence to sugarcane and weed, at 4 pts./A. Treat before weeds exceed 1.5 inches in height.

2. Apply 1-3 times, as needed, at 4 pts./A postemergence to sugarcane and weeds. Treat before weeds exceed 1.5 inches in height.

Louisiana

For control of annual weeds during the summer fallow period, apply 4 pts./A to weed-free beds immediately after bed formation. Follow normal weed control program after planting.

Precautions: To avoid crop injury, (1) Do not apply more than 10 pts./A to any one crop of sugarcane. (2) If making 4 pts./A application during summer fallow period, do not exceed 8 pts./A during the remainder of the growing season, or illegal residues may result.

Texas

Use Weedpro Alatazine 4L for control of barnyardgrass, pigweed, purslane and sunflower, in plant or ratoon sugarcane.

Apply 8 pts./A of Weedpro Alatazine 4L preemergence. Follow with one or two applications, as needed, at 8 pts./A postemergence to sugarcane and weeds.

Precautions: (1) Injury to sugarcane may occur when under moisture stress, when soil is of low adsorptive capacity, or when land is first cropped to sugarcane. (2) Do not apply after close-in. (3) Do not apply more than 30 pts./A to any one crop of sugarcane, or crop injury may result.

Turfgrasses-for-Bed-(Florida-only)

St. Augustinegrass, Centipedegrass, and Zoysia Grass
Broadcast 4-8 pts./A according to soil texture to control those weeds listed under **Weedpro Alatazine 4L Applied Alone - Corn or Grain Sorghum - Preplant Surface-Applied, Preplant Incorporated, or Preemergence**.

Soil or peat	8 pts.	Old beds; Within 2 days after tilling bed. New beds: 3-4 days after spraying or plowing	Old beds: Within 2 days after tilling bed. New beds: 7-10 days after spraying or plowing
Sandy soil	4 pts.	Old beds: Within 2 days after tilling bed. New beds: 7-10 days after spraying or plowing	Old beds: Within 2 days after tilling bed. New beds: 7-10 days after spraying or plowing

If weeds regrow, apply an additional 4 pts./A on moist or peat, or 2 pts./A on sandy soil.

Precautions: To avoid crop injury, (1) Do not apply within 30 days prior to cutting

22

Use tillage to zone planting. For JAS or Gramoxone Extra, 1 application.

Additional Tank-Mix Fertilizer (CO, HI)

If weeds are present Apply 32 ounces of J Use 44 ounces of F Use 52 ounces of T and Washington, do will be suppressed acer for control. T increase the perform applications, apply in 3 to 5 gallons of wa precautions, and lim formulatio

Landmaster II (CO, HI)

If weeds are present Apply 40 ounces of L Use 54 ounces Land 64 ounces of Landma to control grasses spe Landmaster II per ac Ammonium Sulfate is ter II and Weedpro A pts. of Weedpro Ara gallons of water/A. Refer to the Landma

Roundup RT (CO, HI)

If weeds are present a 12 ounces of Roundup ounces Roundup RT of Roundup RT plus nonionic surfactant in Washington, do not e

19

or till. (2) Do no additives. (3) Use on nematodes, and disease of growth may follow

Turfgrasses-for-Fa

'In states other than Turfgrasses-for-Sod

Bermudagrass, Cen

Apply Weedpro Alat weeds for control of a grass species, herb control annual bluergr of summer annual we Alatazine 4L Applied / apply Weedpro Alat a minimum of 15 gals.

Where annual bluer 1,000 sq. ft.). Use 2 qt other weeds named a striped turfgrass or Ormond.

For continued summe days after the previous more than two applica

Precautions: On new sowing of growth and injury. (1) Use only a nematodes, and disease of NC (except may be a species of nematode, (2) Do not apply over 1 lb. (3) Do not over mow, after treating bermudagrass until it system. Do not excess grass.

Note: Do not graze o

23

Macadamia Nut

For preemergence control of many broadleaf and grass weeds, including crepegrass, fescue, ryegrass, Florida's paintbrush, Spanish needles, and fireweed, broadcast 4-8 pts./A before harvest and before weeds emerge. Repeat as necessary. Do not spray when nuts are on ground during harvest period. Do not apply by air.

Quince

Use only on established plantings which are at least 18 months old. Apply as a directed spray at 4-8 pts./A of Weedpro Atrazine 4L in 20-50 gals. of spray mix preemergence or early postemergence to weeds. When applying postemergence, the use of a surfactant and greater spray volume (80-100 gals. of spray mix per acre) may enhance weed control. This product controls many annual broadleaf and grass weeds, including fireweed, purslane, scarlet pimpernel, Spanish needles, and sowthistle.

Notes: To avoid illegal residues, (1) Do not allow spray to contact foliage or fruit. (2) Do not apply more frequently than at 4 month intervals. (3) Do not apply more than 16 pts. of Weedpro Atrazine 4L per year.

Conifers

For control of annual broadleaf and grass weeds prior to transplanting, after transplanting or in established conifers (including Douglas fir, grand fir, noble fir, white fir, Austrian pine, bishop pine, Jeffrey pine, knobcone pine, loblolly pine, longleaf pine (Shore pine), Monterey pine, ponderosa pine, Scotch pine, slash pine, blue spruce, and Sitka spruce). Broadcast 4-8 pts. in a minimum of 5 gals. of water per acre by air or 10 gals. by ground while trees are dormant. For new transplants, apply during or soon after transplanting. For applications prior to transplanting, allow sufficient precipitation to activate Weedpro Atrazine 4L before transplanting. In areas where spring and summer rainfall is inadequate to activate Weedpro Atrazine 4L, apply during fall prior to spring transplanting.

For the list of weeds controlled, see Weedpro Atrazine 4L Applied Alone - Corn or Grain Sorghum - Preplant Surface-Applied, Preplant Incorporated, or Preemergence.

24

grazing may begin 7 months after a fall application or 3 months after a winter or spring application. (2) To prevent illegal residues, do not graze treated areas of the Great Basin, or areas east of the Rocky Mountains. (3) Temporary injury to trees may occur following use of Weedpro Atrazine 4L or coarse-textured soil. (4) To avoid crop injury, do not apply to seedbeds. (5) Also apply only once per year.

Aerial application: In order to assure that spray will be controllable within the target area when used according to label directions, make applications at a minimum height of 10 ft. above vegetation, using low drift nozzles at a maximum pressure of 40 psi, and restrict application to periods when wind speed does not exceed 10 mph. To assure that spray will not adversely affect adjacent sensitive nontarget plants, apply Weedpro Atrazine 4L by aircraft at a minimum upwind distance of 400 ft. from sensitive plants.

Hazardous Weed Control on Noncrop Land

This product provides long-term weed control on industrial sites, highway medians and shoulders, railroad rights-of-way, lumberyards, petroleum tank farms, and in noncrop areas on farms, such as around buildings, equipment and fuel storage areas, along fences, and lanes. Do not use near adjacent desirable plants or in greenhouses, or injury may occur.

Apply before or soon after weeds begin growth. Make postemergence applications when weeds are young and actively growing. Use the higher rates on fine clay and muck soils. Use sufficient water to assure thorough coverage. Use at least 1 gal. of water for each quart of product; use more if practical.

Annual broadleaf and grass weeds (including barnyardgrass, cheatgrass, crabgrass, lambquarters, foxtail, ragweed, puncturevine, and turkey mullein): Broadcast 4-8-10 qts./A.

For small areas, 0.28 pt. per 1,000 sq. ft. is equivalent to 10 qts./A.

Precautions: (1) Do not allow livestock to graze treated areas, or illegal residues may result. (2) Do not exceed 10 lbs. active ingredient (10 qts. of the product) per acre per calendar year.

Storage and Disposal

Storage

Ground water contamination may be reduced by diking and flooring of permanent liquid bulk storage sites with an impermeable material.

25

I - do Disposal

Do not contaminate water, food, or feed by storage or disposal, or cleaning or equipment. Open dumping is prohibited. Wastes resulting from the use of this product are acutely toxic. Improper disposal of unused pesticide, spray mixture, or rinsate is a violation of federal law. Pesticide, spray mixture, or rinsate that cannot be used according to label instructions must be disposed of according to federal, state or local procedures. For guidance in proper disposal methods, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office.

Container Disposal

Triple rinse (or equivalent) and offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by incineration, or by open burning, if allowed by state and local authorities. If burned, keep out of smoke.

For minor spills, leaks, etc., follow all precautions indicated on this label and clean up immediately. Take special care to avoid contamination of equipment and facilities during cleanup procedures and disposal of wastes. In the event of a major spill, fire or other emergency, call CHEMTRAC at 1-800-424-9300.

Caliber® trademark of CIBA-GEIGY

Compa® trademark of KALO Agricultural Chemicals, Inc.

Concap® trademark of CIBA-GEIGY

Cyclone® trademark of ICI AMERICAS for paraquat

Dual® trademark of CIBA-GEIGY for metolachlor U.S. Patent No. 3,937,730

Fallowmaster® trademark of MONSANTO Company

Gramoxone® trademark of ICI AMERICAS for paraquat

Landmaster IP® and **Landmaster BW®** trademark of MONSANTO Company

Lasso® trademark of MONSANTO Company for alachlor

Latron AG-44® trademark of ROHM & HAAS

Princip® trademark of CIBA-GEIGY for simazine

Ramrod® trademark of MONSANTO Company for propachlor

Roundup® and **Roundup RT®** trademark of MONSANTO Company

Sprayfuse 90® trademark of CORNBELT Chemical Co.

Sutan +® trademark of STAUFFER Chemical Company for butylate

Precautionary Hazards to Humans

Hand & eye irritant, v/c or spray mist. Auto.

Users are required to wear chemical resistant gloves in mixing, rubber or -synthetic.

STATEMENT OF USE
IF SWALLOWED: Drink one or two glasses of water. Do not induce vomiting.

IF INHALED: Remove person, preferably to fresh air.

IF ON SKIN: Wash with water.

IF IN EYES: Flush with water.

Note to Physician: Ingested, induce emesis. Activated charcoal. Environmental Health.

Atrazine can travel which may be used.

Users are advised not to drink water from the water table (ground permeable, i.e., we further information water).

This product may be including abandoned.

This pesticide is toxic to wetlands. Do not use in weather conditions. These areas may have hazards that contaminate water.

BEST AVAILABLE COPY

26