


"U.S.D.A. Registration No. 10108 3"



Poisoned Grain
for
Ground Squirrels
and **Mice**

DANGER

KEEP OUT OF REACH OF CHILDREN

WARNING

Convulsive Poison.

Do not contaminate feed or food stuffs. Keep away from children and domestic animals. See antidote on back panel.



POISON

Active Ingredients:	
Strychnine Alkaloid	0.30%
Inert Ingredients	99.70%
	100.00%

Net Weight 3 lbs.

Price \$2.50

RED STAR POISON GRAIN CO.
 Rt. 1 Box 70E
 WOODBURN, OREGON

ANTIDOTE

When some time has passed since the poison was taken, give a tablespoon of salt in a glass of warm water. Have victim lie down in a quiet, darkened room and keep warm. Call a physician immediately.

Directions - Ground Squirrels

When putting this poison out use teaspoonful to a tablespoonful of bait distributed near the entrance of burrows and along trails made by the squirrels. This bait has been found effective on the Oregon and Douglas ground squirrels. The best time for poisoning these squirrels is early spring around March or when they become active, as they are hungry and take the bait more readily. Treated seeds exposed on soil surface may be hazardous to birds and other wildlife.

Directions - House Mice

Place rounded teaspoon of bait at 8 to 10 foot intervals where mice feed, water or travel. Place in locations not accessible to children, pets and domestic animals, or use tamper-proof bait boxes. Collect and destroy all uneaten baits when through.

**Do not reuse container -
destroy when empty.**

ANTIDOTE

If less than ten minutes has passed since the poison was taken, give a tablespoon of salt in a glass of warm water. Have victim lie down in a quiet, darkened room and keep warm. Call a physician immediately.


Directions – Ground Squirrels

When putting this poison out use teaspoonful to a tablespoonful of bait distributed near the entrance of burrows and along trails made by the squirrels. This bait has been found effective on the Oregon and Douglas ground squirrels. The best time for poisoning these squirrels is early spring around March or when they become active, as they are hungry and take the bait more readily. Treated seeds exposed on soil surface may be hazardous to birds and other wildlife.

Directions – House Mice

Place rounded teaspoon of bait at 8 to 10 foot intervals where mice feed, water or travel. Place in locations not accessible to children, pets and domestic animals, or use tamper-proof bait boxes. Collect and destroy all uneaten baits when through.

**Do not reuse container –
destroy when empty.**



RED STAR POISON GRAIN CO.
Rt. 1 Box 70E
WOODBURN, OREGON

RED STAR POISON GRAIN CO.
Rt. 1 Box 70E
WOODBURN, OREGON