

NET WEIGHT 50 POUNDS (22.7 kg.)
EPA REG. No. 18114-2
CPA EST. No. 4841-LA-1

DIRECTIONS

It is a violation of Federal law to use this product in a manner inconsistent with its labeling

FOR USE ON NON-CROP LAND SUCH AS FARM ROAD AND PIPELINE RIGHT-OF-WAYS, INDUSTRIAL PLANTS, LUMBER YARDS, TANK FARMS, STORAGE AREAS AND PARKING LOTS for the control of most annual and perennial grasses and broadleaf weeds. Effects are slow to appear and will not become apparent until the chemical has been carried into the root zone of the weeds by moisture. Control will last for an extended period. The effectiveness and length of time control will last depends on type of soil, amount of rain and other conditions

HOW TO APPLY: Use a mechanical spreader or other device which will spread the required amount evenly over the area to be treated. It may be applied when top growth is dense, but better control will be obtained when weeds are short or when excess growth is mowed and removed before applying.

WHEN TO APPLY: Best results will be obtained if applied shortly before or at the time plant growth begins. Rain or thorough watering after application will cause the material to move to the root zone and thus speed up control. DO NOT use when ground is frozen

GENERAL WEED CONTROL


ANNUAL WEEDS AND GRASSES: For foxtail, crabgrass, wild oats, bromegrass, ryegrass, pigweed, ragweed, turkey mullein, and lambquarters use 60 to 120 pounds per acre

PERENNIAL WEEDS AND GRASSES: For quack grass, bluegrass, smooth brome, Bahig, sea, nuttop, purpletop, bromesedge, aster, plantain, dandelion, wild carrot, dog fennel, and goldenrod use 140 to 240 pounds per acre.

FOR HARD-TO-KILL WEEDS AND GRASSES such as Johnsongrass, Bermuda Grass, nutgrass, dallisgrass, Vasey grass, setgrass, bouncing bet, dogbane, bracken fern, and horsetail use 300 to 600 pounds per acre.

Use the higher level rates given on soils high in organic matter

Form ASL1330



Precautionary Statements Hazards to Humans and Domestic Animals Caution

Harmful if swallowed. May cause irritation of eyes, nose, throat and skin. Avoid breathing dust. Wash thoroughly after handling. Remove and wash contaminated clothing before reuse.

In case of contact with eyes, flush with plenty of water for 15 minutes. Get medical attention if irritation persists. In case of contact with skin, wash with soap and water.

ENVIRONMENTAL HAZARDS

Do not apply directly to water. Do not contaminate water by cleaning of equipment or disposal of wastes. Do not apply, allow to drift or drain or flush equipment onto or near croplands, lawns, trees or other desirable plants or to areas where their roots may extend. Application equipment should be thoroughly cleaned after use. Equipment which has been used to apply this product should not be used for applying insecticides, fungicides, fertilizers or other agricultural chemicals as plant injury may result.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

Pesticide Disposal: Pesticides that cannot be used according to label instructions must be disposed of according to applicable Federal, state, or local procedures.

Container Disposal: Destroy container so that it may not be used again for any purpose and dispose of it at some location where no attempt to salvage will be made.

ASPLUNDH WARRANTS THAT THE COMPOSITION OF THIS PRODUCT CONFORMS TO THE INGREDIENT STATEMENT AND IS SUITED TO THE PURPOSES DESCRIBED WHEN USED ACCORDING TO DIRECTIONS. THE USE OF THIS PRODUCT BEING BEYOND THE CONTROL OF THE MANUFACTURERS, NO GUARANTEE, EXPRESSED OR IMPLIED, IS MADE AS TO THE EFFECTS OR RESULTS TO BE OBTAINED IF NOT USED IN ACCORDANCE WITH DIRECTIONS OR ESTABLISHED SAFE PRACTICES. BUYER ACCEPTS ALL RISKS OF USE, STORAGE OR HANDLING OF THIS MATERIAL NOT IN STRICT ACCORDANCE WITH DIRECTIONS GIVEN HEREWITH.

Made in U.S.A.

Printed in U.S.A.

Expressly for
ASPLUNDH TREE EXPERT CO.
Blair Mill Road, Willow Grove, Pa. 19090

BEST AVAILABLE COPY