M

[7]

1



BO-RID CBB

LIQUID WEED KILLER

EPA Reg. No. 179-85

1 gallon contains 1.5 lbs. sodium chlorate 1 gallon contains 0.75 lbs. sodium metaborate anhydrous 1 gallon contains 0.05 lbs. bromacil

For use in killing weeds and grasses to keep areas free of vegetation and for treating soil prior to topping with asphalt or pavement. May be used in such locations as around warehouses, storage yards, camping places, recreational areas, fence lines and other similar uses where there is no danger to valuable plants, or where unproductive soil is not objectionable.

ACTIVE INGREDIENTS:

Sodium chlorate (NaClo ₃)	14.00%
Sodium metaborate anhydrous (Na ₂ B ₂ O ₄)	7.00%
Bromacil (5-bromo-3-sec-butyl-6-methyluraci	.46%
INERT INGREDIENTS (Water)	78.54%
Total	100.00%

*Equivalent to 1.15% boron expressed as elemental boron

CAUTION:

KEEP OUT OF REACH OF CHILDREN

CAUTION: Harmful if swallowed! Avoid getting in eyes; avoid breathing spray mist. Wash clothing and equipment after application. Do not leave containers where children and animals may take the chemical internally. Avoid contamination of, and storage near, feed and food products. Do not apply on or near turf, valuable trees or other plants, or in areas into which their roots may extend, or on slopes where "run-off" will move the chemical into the root areas of useful plants. Avoid contamination of irrigation or domestic waters. The chemical should be used only where unproductive soil is not objectional. Avoid drift to desirable plants.

Do not store below 32º F.

When this product has been supplied in drums. DO NOT reuse empty drum. Return to drum reconditioner or destroy by perforating or crushing and burying in a safe place away from water supplies.

KEEP LID TIGHTLY CLOSED Do NOT Store With Foodstuffs

NET CONTENTS _____ Gallons

DIRECTIONS:

GENERAL SPRAYING: Spray to wet plants thoroughly, using power or knapsack sprayer and apply chemical uniformly over infested area. Applications should be made while weeds are small. Avoid applications in hot, dry weather, as the killing action of BO-RID CBB is through both roots and tops of plants, and therefore moisture is necessary after application to take the chemical into the soil to kill deep-rooted perennials.

Fall applications may be made in areas of 20 inches or less rainfall in areas of heavy winter rainfall (20 inches or more) apply in the spring as the chemical is apt to be lost by leaching. Retreatment may be necessary under some conditions to keep areas freed of plant growth. By preventing weed growth early in the year, fire hazard from dead and dry vegetation later on is avoided.

Annual Weeds & Grasses—To control annuals use 2 or 3 pints of solution per 100 sq. ft, and spray to wet plants thoroughly.

Perennial Weeds & Grasses — Use 5 to 6 pints per 100 sq. ft, and apply in sufficient water for uniform coverage. For control of weeds with creeping rootstalks, treat soil 6 ft. beyond visible vegetation to get creeping roots.

Local conditions may affect the use of this chemical. Consult State agricultural extension or experiment station specialists for specific recommendations as to dosage and time of application and for the control of other weeds.

Follow directions carefully. Timing and method of application, weather and crop conditions, mixtures with other chemicals not specifically recommended, and other influencing factors in the use of this product are beyond the control of the seller.

NOTICE TO BUYER —Seller makes no warranty, expressed or implied, concerning the use of this product other than indicated on the label, Buyer assumes all risk of use and/or handling of this material when such use and/or handling is contrary to label instructions.

Manufactured for THE R. H. BOGLE COMPANY. ALEXANDRIX, VA.

JACKSONVILLE, FLA.

MEMPHIS, TENN.

ALVA, OKLA