DIRECTIONS FOR INSIDE USE: For flying insects, fog in processing plants, darrent

For flying insects, fog inside homes, food processing plants, dairies, institutions, warehouses and farm buildings. For best results close all openings and turn off all ventilating equipment before fogging CAUTION. Extinguish all sources of open flame prior to fogging. Fog until dense white smoke fills up entire volume of space. Allow fog to settle completely before reopening doors and win dows. Repeat as necessary.

DIRECTIONS FOR OUTSIDE USE:

Fog around structures such as motels, drive ins, roadside stands, industrial plants, dairy barns, stables, and in such open areas as lawns and gardens, playgrounds, around swimming pools for the control of flies, mos quitoes and other small flying insects. Keep the unit at least 10 feet from plants and shrubbery so as to avoid injury to foliage. Let fog drift in desired direction. Repeat as necessary.

OTHER USES:

Certain household insects such as cock roaches, ants, silverfish and spiders which are accessible and in exposed locations can be controlled by completely fogging area and directing fog into cracks and crevices. Allow fog to settle completely before opening doors or windows. Repeat as necessary



MULTI-PURPOSE ISTALLIFO STATE For control of Flies, Mosquitoes and other small flying insects.

AN INSECTICIDE FOR USE IN HIGH TEMPERATURE FOGGING UNITS SUCH AS THE TIFA, SWING FOG, AND DYNA FOG

CAUTION KEEP OUT OF REACH OF CHILDREN See side panel for additional cautions

| ACTIVE INGREDIENTS: | | BY WEIGHT | |
|------------------------------|---|-------------|----------|
| Petroleum Distillate . | | | . 98 2% |
| Technical Piperonyl Butoxide | | | 1.5% |
| Fyrethrins | | | 0 3% |
| Equivalent to 1 | Butylcarbityl 6 Propylpiperchyl ether and 0.3 | related cor | npour ds |
| | USDA REGISTRATION No. 10130-4 | | |



CAUTION

Repeated or prolonged contact with skin may cause toxic symptoms. Avoid inhalation or skin contact. In case of spillage on skin, wash with soap and water. Avoid contamination of foods. Do not use inside dairies or food processing plants, when plants are in operation. All utensils and surfaces which contact food should be covered. Cover fish ponds while fogging. Harmful if swallowed, keep out of ach of children. Occupants should vacate before indoor treatments and should not reoccupy the premises until it has been ventilated. All food processing surfaces should be cleaned tefore using

INSIST ON "ROSE" DEPENDABILITY

When you use genuine ROSE pest control products you are assured of highest quality and top effectiveness when applied according to directions. Rose Exterminator Co has specialized in scientific pest control products and professional technical service since 1860 and has built a reputation for dependability. Check Rose Exterminator Co. also for bird management, industrial weed control, odor removal and licensed furnigation.

NET GALLONS

