

**DIRECTIONS**


1. Remove gross filth and heavy soil deposits from surfaces prior to application.
2. Spray surface until completely wet from a distance of 6-10'
3. Wipe off excess moisture.
4. For Mildew control, clean surfaces and repeat application seven (7) days as needed.

If used on food contact surfaces, all such surfaces must be rinsed with potable water before reuse.

Empty containers must be rinsed thoroughly with water before discarding.

Non-flammable

NET CONTENTS: GALLON U.S. MEASURE



**BACTERIOSTATIC**

**PREVENTS MOLD AND MILDEW, REDUCES THE HAZARD OF CROSS INFECTION FROM ENVIRONMENTAL SURFACES**

ACTIVE INGREDIENTS:

Cetyl Pyridinium Bromide .....	.06250%
Citral .....	.01250%
Disodium salt of ethylene diamine tetraacetic acid .....	.00625%
INERT INGREDIENTS: .....	99.91875%

EPA Reg. No. 10141-5  
EPA Est. No.

**CAUTION** KEEP OUT OF REACH OF CHILDREN  
Avoid Contamination of Food.

**HOLLISTON LABORATORIES, INC.**

440 TOTTEN POND ROAD - WALTHAM, MASS. 02154

®Registered trademark of Holliston Laboratories, Inc.,  
for the patented disinfectant composition.  
U.S. Patent 3,595,975

**BAC-GARD®**

**BAC-GARD** is broad spectrum microbicidal-microbistatic spray which is effective in presence of soap films.

**BAC-GARD** may be used in meat and poultry processing establishments, super markets, food stores, restaurants, public transportation, hotels, motels, hospitals, nursing homes, vegetable and fruit ripening storage areas, schools, and institutions.

**BAC-GARD** - for use on metal, painted wood, plastic, rubber, tile, porcelain, and all washable surfaces.

**BAC-GARD** is a surface disinfectant.

**BAC-GARD** prevents mold and mildew.

**BAC-GARD** reduces the hazard of cross-infection from environmental surfaces.