

Authorized for Use Under U.S.D.A. Consumer and Marketing Service Inspection Program for Poultry, Meat, Rabbit and Egg Production Plants.

U.S.D.A. Registration No. 10147-10

ACTIVE INGREDIENTS

Trichloro-s-triazinetriene* 15%

INERT INGREDIENTS 85%

TOTAL 100%

*Also known as Trichloroisocyanuric acid.

WARNING

Harmful or fatal if swallowed.

ANTIDOTE: External — flood skin or eyes with plenty of water for 15 minutes. If irritation to skin persists, get medical attention. For eyes — call physician immediately. Internal — drink large quantities of water or milk. Follow with milk of magnesia, vegetable oil, or beaten eggs. Call physician immediately.

Do not get on skin or in eyes.

Causes severe skin burns or damage to the eyes.

AVOID CONTAMINATION OF FOOD.

USE AS DIRECTED.

KEEP DRY.

KEEP CONTAINER CLOSED WHEN NOT IN USE.

DO NOT REUSE CONTAINER.

DESTROY WHEN EMPTY.

**BIRKO
CHEMICAL**

DENVER, COLORADO

SYNCHLOROZENE

FOR

HIDE-BRINE CURING

WARNING: KEEP OUT OF REACH OF CHILDREN

NOT FOR USE OR STORAGE AROUND THE HOME ENVIRONMENT

SEE ADDITIONAL PRECAUTIONS ON LEFT SIDE OF LABEL

POUNDS

•
•
•
•
•
•
•
•
•
•

DI
IN
Ad
gall
SYN
ZEN

**BERKO
CHEMICAL**

EVANSTON, ILL. U.S.A.

SYNCHLOROZENE

FOR

HIDE-BRINE CURING

WARNING: KEEP OUT OF REACH OF CHILDREN


NOT FOR USE OR STORAGE AROUND THE HOME ENVIRONMENT

SEE ADDITIONAL PRECAUTIONS ON LEFT SIDE OF LABEL

POUNDS

ADVANTAGES

- POSITIVE ODOR CONTROL.
- REDUCES NUMBER OF SINKING HIDES.
- LONG LASTING RESIDUAL AVAILABLE CHLORINE.
Dissolves slowly — Remains over 24 hours.
- CONCENTRATED — One ounce SYNCHLOROZENE per gallon yields 1055 ppm available chlorine.



DIRECTIONS

IN RACEWAYS FOR HIDE — BRINE CURING.
Add one pound SYNCHLOROZENE per 5000 gallons of water daily or about one pound SYNCHLOROZENE per ten pounds of SYNTALAZENE added to the raceway.

Authorized for Use Under U.S.D.A. Consumer
and Marketing Service Inspection Program for
Poultry, Meat, Rabbit and Egg Production Plants.

U.S.D.A. Registration No. 10147-10

ACTIVE INGREDIENTS

Trichloro-s-triazinetrione* 15%

INERT INGREDIENTS 85%

TOTAL 100%

*Also known as Trichloroisocyanuric acid.

WARNING

Harmful or fatal if swallowed.

ANTIDOTE: External - flood skin or eyes with plenty of water for 15 minutes. If irritation to skin persists, get medical attention. For eyes - call physician immediately. Internal - drink large quantities of water or milk. Follow with milk of magnesia, vegetable oil, or beaten eggs. Call physician immediately.

Do not get on skin or in eyes.

Causes severe skin burns or damage to the eyes.

AVOID CONTAMINATION
OF FOOD.

USE AS DIRECTED.

KEEP DRY.

KEEP CONTAINER CLOSED
WHEN NOT IN USE.

DO NOT REUSE CONTAINER.

DESTROY WHEN EMPTY.



SYNGHI

HIDE-B

WARNING: KEEP

NOT FOR USE OR STN

SEE ADDITIONA

BIRKO CHEMICAL CORPORATION

P. O. BOX 131



SYNCHLOROZENE

BRINE CURING


SAFETY
OF CHILDREN
ENVIRONMENT
REVERSE OF LABEL

50 POUNDS

1500 S. W. 15TH AVENUE, DENVER, COLORADO
80202

ADVANTAGES

- POSITIVE ODOR CONTROL.
- REDUCES NUMBER OF SINKING HIDES.
- LONG LASTING RESIDUAL AVAILABLE CHLORINE.
Dissolves slowly — Remains over 24 hours.
- CONCENTRATED — One ounce SYNCHLOROZENE per gallon yields 1055 ppm available chlorine.



DIRECTIONS

IN RACEWAYS FOR HIDE — BRINE CURING.
Add one pound SYNCHLOROZENE per 5000 gallons of water daily or about one pound SYNCHLOROZENE per ten pounds of SYNTALAZENE added to the raceway.