


MILL SPRAY #4001

CAUTION

This product is toxic to fish and birds. Keep out of lakes, streams or ponds. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on this label.

Remove pets, birds, and cover fish aquariums before spraying.

Do not reuse empty drum. Return to drum reconditioner or destroy by perforating or crushing and burying in a safe place away from water supplies.



**ONLY TRAINED PERSONNEL PERMITTED
TO USE THIS PRODUCT**

Do not breathe spray mist. Wear a pesticide respirator of a type jointly approved by the Mining Enforcement and Safety Administration, and by the National Institute of Occupational Safety and Health under the provisions of 30 CFR Part 11. Wash hands, arms and face with soap and water after handling and before eating or smoking. Wash all contaminated clothing with soap and hot water before reuse.

A Ready To Use Insecticide For Control of Insect Pests of Grain and Food Products

For Use In Flour Mills, Elevators, Granaries, Food Processing Plants and Warehouses, Bakeries, Beverage Plants, Restaurants and Supermarkets

This oil base spray is suitable for application using conventional thermal, mechanical or compressed air fogging equipment.

The Insecticidal Components (Piperonyl Butoxide and Pyrethrins) are Classified as Food Additives and When This Product is Used as Directed, Allowable Tolerance Levels will not be Exceeded.

ACTIVE INGREDIENTS		100
Pyrethrins	0.2	
*Piperonyl Butoxide Technical	2.0	
Petroleum Distillate	97.8	

*Equivalent to 1.8-(4-butylcarbityl)-6-propylpiperonyl ether and to 0.2% of related compounds

**EPA REG. NO. 10148-14
EPA EST. NO. 10148-NY-1**

CAUTION

Keep Out Of Reach Of Children

Harmful if swallowed.

See rear panel for additional precautions

Net Contents

U.S. Gallons

Any infested stored food products should be fumigated or treated by other effective methods using an approved product intended for this purpose.

DIRECTIONS FOR USE

The use of this product in food processing plants should be confined to time periods when the plant is not in operation. Foods should be removed or covered before treatment. All food processing surfaces should be covered during treatment or thoroughly cleaned before using.

Clean out any waste grain, dust, dirt or debris and destroy before spraying.

Remove pets, birds and fish aquariums before spraying.

SURFACE SPRAYING This spray provides rapid initial kill of exposed or accessible stages of the following insects: granary weevil, rice weevil, confused flour beetle, rust red flour beetle, saw toothed grain beetle, spider beetle, cheese mites, and grain mites. Apply as a coarse droplet spray using approximately 1 gallon to 750 square feet of surface. Spray walls, floors, and other surfaces of bins and storage and handling areas thoroughly, being especially careful to treat cracks, crevices and similar hiding places. Treat unloading, handling and processing areas, and inside conveying, processing and handling equipment. Spray thoroughly around the base of heavy machinery and equipment. Spray insects directly when possible.

For in transit protection, clean and spray truck beds, boxcars, and ship's holds before loading. Thoroughly treat walls and floors and spray each layer of cartons or containers lightly as stacked.

To kill exposed stages of Indian meal moth and Angoumois grain moth in stored grain, apply to the surface of binned grain at a rate of 1 gallon per 1000 square feet. Apply at monthly intervals with the first appearance of moths and continuing until early fall. In case of severe infestation, break up webbing with rake before spraying and make a second application in two weeks.

As an aid in the control of mosquitoes, gnats, wasps and skippers, treat walls, ceilings, moldings, screens, door and window frames, beams, light cords, and similar resting places.

SPACE SPRAYING To control flies, mosquitoes, gnats, and small flying moths, close windows, doors, and other ventilation. Using a good sprayer adjusted to deliver a fine mist, direct spray toward the upper portions of the enclosure. Distribute spray throughout entire area using 2 ounces per 1000 cubic feet of space.

Sweep up and destroy fallen insects. Repeat treatment as necessary.

Keep area closed for at least 10 minutes after spraying. Vacate premises and ventilate before reoccupying.

Manufactured By:

Nationwide Chemical Co., Inc.
56 North First Street
Brooklyn, New York 11211