

CAUTION

Avoid contamination of feed and food-stuffs.

The use of this product in food processing plants should be confined to time periods when the plant is not in operation. Food should be removed or covered during treatment. All food processing surfaces should be covered during treatment or thoroughly cleaned before using.

Remove pets, birds, and cover fish aquaria before spraying.

This product is toxic to fish.

Keep out of lakes, streams or ponds.

Do not contaminate water by cleaning of equipment or disposal of wastes. Apply only as specified on the label.

Do not reuse empty drum. Return to drum reconditioner or destroy by perforating or crushing and burying in a safe place away from water supplies.

INDUSTRIAL SPRAY #4000

**Effective, Fast Acting Insecticide.
For Use In Institutions, Restaurants
and Food Processing Plants.**

Kills on contact exposed stages or accessible stages of:

Roaches	Gnats	Rice Weevil
Silverfish	Wasps	Grain Mites
Crickets	Hornets	Cheese Mites
Spiders	Sand Flies	Carpet Beetles
Ants	Mosquitoes	Angoumois Grain Moth
Fleas	Bedbugs	Confused Flour Beetle
Flies	Meal Worms	Saw Toothed Grain Beetle
	Meal Moths	Mediterranean Flour Moth

ACTIVE INGREDIENTS	100
Pyrethrins	0.10
*Piperonyl Butoxide, Technical	1.00
Petroleum Distillate	98.90

*Equivalent to 0.8% (butylcarbityl) (6-propylpiperonyl) ether and 0.2% of related compounds.

CAUTION

Keep Out Of Reach Of Children


Harmful If Swallowed

See rear panel for additional precautions

Net Contents

Gallons

Manufactured By:
Nationwide Chemical Co., Inc.
56 North First Street
Brooklyn, New York 11211



EPA REG. NO. 10148-15
EPA EST. NO. 10148-NY-1

DIRECTIONS FOR USE

SPACE SPRAY: For rapid control of flies, fruit flies, mosquitoes, gnats, small flying moths, sand flies, wasps, and hornets. Close doors and windows and shut off ventilating systems. Apply using a good machine adjusted to deliver a fine mist. Direct spray toward the upper portions of the room to fill the entire area with mist using 1 to 2 ounces per 1000 cubic feet of space. Vacate the treated area and ventilate before reoccupying. Keep the area closed for ten minutes after treating. Repeat application when reinfestation occurs. Sweep up and destroy fallen insects.

SURFACE APPLICATION:

To kill accessible, exposed stages of roaches, silverfish, flies, and spiders. Apply as a coarse wetting spray directed into locations where these insects may hide behind baseboards, moldings, cabinets, shelves, sinks, door and window frames, around plumbing drains and in dark corners and into cracks and crevices. Hit flying insects with the spray whenever possible. Repeat treatment as necessary.

Against infestation, treat shelves, cabinet and terrace areas, spray ant runs, baseboards, thresholds and other places of entry, and treat nests liberally. Repeat treatment as necessary.

In controlling flea and Brown Dog ticks in houses, direct the application toward the lower wall portions, baseboards, rugs, carpets and floors of infested areas. Animal quarters, kennels and bedding should also be sprayed. Repeat treatment as needed.

To control bedbugs effectively, apply liberally to top, ends, sides and bottom of mattresses, spraying thoroughly all tufts, seams and folds. Spray springs and frame, directing the liquid into all cracks and lock joints. Treat walls, baseboards, moldings, floors, closets and bedroom furniture, picture frames, with particular attention to protected bedbug hiding and breeding places. Repeat treatment as needed.

For most effective control of clothes moths and carpet beetles, remove infested articles from storage, brush thoroughly and, if possible, air several hours in direct sunlight. Apply spray liberally in empty chests, closets, bureaus and storages, directing the liquid into cracks, joints and crevices. After airing, infested articles may be treated by holding the spray nozzle at least 3 feet away from the fabric and applying the liquid lightly to the seams to avoid the possibility of staining. Treat both sides of rugs, carpets and pads, and all exposed and accessible surfaces of overstuffed furniture. Do not overspray if products are made of rubber or are rubber coated, to prevent expansion or swelling. Repeat treatments monthly unless the articles are stored in moth-tight containers.

To control infestations of granary weevil, rice weevil, saw-toothed grain beetle, confused flour beetle, rusty grain beetle, red flour beetle, meal worms, grain mites, Angoumois grain moth, Indian meal moth, and Mediterranean flour moth in food storage areas: If possible, clean out and destroy any infested food products; otherwise, fumigation or other effective treatment should be made using an approved product intended for this purpose. Apply this spray thoroughly to all surfaces of shelves, cabinets, bins, pantries, and similar food storage areas with particular attention to cracks, crevices, niches, and other hiding places. Repeat treatment as needed.

Any infested stored food products should be fumigated or treated by other effective methods using an approved product intended for this purpose.