


GLOBE

T - F

TURF FUNGICIDE (Contains Dyrene)*

DIRECTIONS FOR USE:

T-F Turf Fungicide gives effective control of *Rhizoctonia* brown patch, *Sclerotinia* dollar spot, copper spot, rust, snow mold, *Typhula*, and *Helminthosporium* diseases, such as melting-out of Kentucky bluegrass, red leaf spot of bentgrasses, leaf blotch of Bermuda Grass, *Helminthosporium* blight of fescues, brown blight of rye grasses, zonate eye spot, and *Helminthosporium* leaf spot of bluegrasses, bentgrasses, and fescues.



HOW TO USE:

In Garden Hose Sprayers: Put T-F Turf Fungicide into sprayer jar, fill with water and stir until a uniform mixture is obtained. Apply spray mixture uniformly to lawn area. For disease prevention use 4 ounces approximately 1 cup T-F Turf Fungicide and spray to cover 1,000 square feet. For spot treatment of diseased areas, use 6 ounces approximately 1 1/2 cups T-F Turf Fungicide and thoroughly wet these areas.

In Compressed Air Sprayers: For use in hand or knapsack-type sprayers, add 4 ounces approximately 1 cup of T-F Turf Fungicide to 2 1/2 gallons of water and apply to 1,000 square feet 50 ft. x 20 ft. of lawn or other turf area. For spot treatment of heavily diseased areas add 6 ounces approximately 1 1/2 cups to 2 1/2 gallons of water and thoroughly wet these areas.

ACTIVE INGREDIENT:

2, 4-Dichloro-6-(o-chloroanilino)-s-triazine 50%

INERT INGREDIENTS: 50%
100%

U.S. Patent No. 2,720,480 & 3,074,946
Canadian Patent No. 546,027

KEEP OUT OF REACH OF CHILDREN

CAUTION: SEE SIDE PANELS FOR WARNING STATEMENTS

STOP -- READ THE LABEL BEFORE USE

AN EXCLUSIVE PRODUCT OF

GLOBE CHEMICALS, INC.

45 Burbank Dr.
Toledo, O.

P.O. Box 713
Camp Hill, Pa.

WHEN TO APPLY:

For Foliar Turfgrass Diseases: Begin applications in the spring before diseases appear and repeat at 7 to 14 day intervals. During extended periods of wet weather, application every 7 days may be required. Continue treatments throughout the spring and summer months. If necessary, water the lawn before applying T-F Turf Fungicide.

For Control of Snow Mold: Apply after the first killing frost in the fall. A second application should be made when the lawn or other turfgrass area is free of snow in mid-winter. Apply a third application after the final melting of snow in late winter or early spring.

CAUTION:

Harmful if swallowed or inhaled. Avoid contact with eyes, skin, or clothing. Avoid breathing of dust or spray mist. Wash thoroughly with soap and warm water after handling. Wash contaminated clothing with soap and hot water before reuse.

Do not contaminate feed, foodstuffs, water supplies, and fish ponds. Keep out of reach of children.

This product is toxic to fish. Keep out of any body of water.

Do not graze treated areas. Do not feed droppings to livestock.

Never reuse container. Dispose of container by burying or burning. Keep out of smoke.

© 1971 Globe Chemicals, Inc. All rights reserved. For other information, contact Globe Chemicals, Inc.

NET CONTENTS _____ lbs.

USDA Reg. No. 10155-9