


10159-2

8-27-1971

4/3



Best Available Copy

(Front Panel)

VPG

5% SEVIN* DUST

ACTIVE INGREDIENTS:

Carbaryl (1-naphthyl N-methylcarbamate) 5.00%

INERT INGREDIENTS: 95.00%

Total 100.00%

CAUTION (18 Point)

KEEP OUT OF REACH OF CHILDREN (12 Point)

SEE SIDE PANEL FOR ADDITIONAL CAUTIONS (12 Point)

*Registered trademark of Union Carbide Corporation for the active ingredient 1-naphthyl N-methylcarbamate.

NET WEIGHT 4 POUNDS

(Back Panel)

VPG

5% SEVIN DUST

1. WHAT IS THIS PACKAGE FOR?

This package of VPG 5% SEVIN DUST provides an economical and easy method to control many insect pests as listed in the directions for use.


2. HOW DO YOU USE IT?

ON BIRDS -- to control Northern fowl mite, chicken mite, lice, fleas, use 1 lb. 5% Dust per 100 birds.

ON LITTER -- to control Northern fowl mite, chicken mite, lice, fleas, bedbugs, use 1 lb. 5% Dust per 40 sq. ft.

IN DUST BATH BOXES -- to control Northern fowl mite, chicken mite, lice and fleas, use 2½ lbs. 5% Dust per box for each 50 birds.

ON DOGS AND CATS -- for ticks and fleas dust liberally over the entire animal, avoiding the eyes, and rub thoroughly into the skin. Comb out dead ticks a few hours after treatment. Repeat at weekly intervals if needed. Avoid treatment of kittens less than 4 weeks of age. In and around Dog and Cat Quarters: For ticks and fleas dust liberally in and around sleeping quarters, kennels, and dog houses by means of a shaker can, hand or dust gun.



BEANS (green beans, lima beans, snap beans, cowpeas, black-eyed peas) -- to control Mexican bean beetle, bean leaf beetle, cucumber beetles, flea beetles, Japanese beetle, leafhoppers, velvetbean caterpillar, western bean cutworm, armyworms, corn earworm, lima bean pod borer, stink bugs and tarnished plant bugs.

SWEET CORN -- to control corn earworm, corn rootworm adults, European corn borer, fall armyworm, flea beetles, Japanese beetles, sap beetles and leaf hoppers. For larvae in whorl (budworm damage) and foliage feeders, apply to entire plant. Repeat as necessary.

For insects attacking silks and ears, apply at 2- to 3- day intervals starting when first silks appear, and continuing until silks begin to dry. Three or more applications may be required depending on severity of infestation. Timing and good coverage are essential for effective control.

CUCUMBER, MELONS, PUMPKIN, SQUASH -- to control pickleworm, melonworm, cucumber beetles, flea beetles, leafhoppers and squash bugs. Avoid excessive applications. Some leaf injury may occur if treatments are made when tender foliage is wet or in the presence of prolonged high humidity. SEVIN injures watermelons in Florida.

POTATO, TOMATO, EGGPLANT, PEPPER, PEAS -- to control Colorado Potato beetle, flea beetles, leafhoppers, European corn borer, fall armyworms, lace bugs, tomato fruit worm, tomato hornworm, tarnished plant bugs, stink bugs.

OKRA -- to control corn earworm and stink bugs.


CABBAGE, BROCCOLI, BRUSSELS SPROUTS, CAULIFLOWER, KOHLRABE -- to control flea beetles, harlequin bug, armyworms, cabbage caterpillars, corn earworm. To control small cabbage loopers, use a 5- to 7- day schedule.

STRAWBERRIES -- to control meadow spittlebug, strawberry leaf roller and strawberry weevil.

BLACKBERRIES, BOYSENBERRIES, DEWBERRIES, LOGANBERRIES AND RASPBERRIES -- to control Japanese beetles, leaf rollers, leafhoppers and raspberry aphids.

APPLY 5% SEVIN DUST AS OFTEN AS NECESSARY TO CONTROL GRASSHOPPERS ON THE ABOVE CROPS.

5% SEVIN DUST can be safely used on a wide range of ornamental plants, flowers, shrubs and small trees, such as rose, carnation, gladiolus, chrysanthemum, lilac, arborvitae, juniper, Hydrangea, azalea, elm, maple, oak, dogwood, birch and pines. SEVIN injures Boston Ivy.



(Back Panel)

Page 3.

Apply to the upper and lower leaf surface of all foliage for complete coverage to control such pests as: Apple aphid, June beetles, leaf-hoppers, leaf roller, boxwood leaf miner, mimosa webworm, orange tortrix, willow leaf beetles, Japanese beetle, psyllids, rose aphids, boxwlder bug, scale insects, oak leaf miners, thrips (exposed), Plant Bugs, lace bugs, rose-slug, mealy bugs, elm leaf aphid, flea beetles, gypsy moth, Bagworm, puss caterpillars, birch leaf miner, blister beetles, thorn bugs, elm leaf beetle, periodical cicada.

SEVIN does not control spider mites. Where infestations are encountered, use a recommended miticide.

Apply 5% SEVIN DUST when insects or their damage appear. Make applications in the morning or evening when it is cool. Treat upper and lower surface of all foliage for complete coverage. Repeat as necessary, generally at 7-day intervals. For best results, apply with a dust gun. Do not apply in excess.

CAUTIONS: Harmful if inhaled or swallowed. Do not breathe dust. Do not take internally. Avoid skin contact. Wash thoroughly after using. Avoid contamination fo feed or foodstuffs. Note for physician: Carbaryl is a moderate, reversible, Cholinesterase inhibitor. Atropine is antidotal.

For protection of honeybees avoid use, if possible, during periods when honeybees are visiting the crop. When necessary to use during such periods, warn beekeepers well in advance to locate hives at a safe distance until one week after application. Injury to tender foliage may occur if applications are made when plants are wet or in the presence of high humidity of several days duration following treatment.

Do not apply within 3 days -- to beans, corn forage or fodder, cucumbers, eggplants, melons, okra, garden peas, peppers, potatoes, pumpkins, summer squash, broccoli, bunssels sprouts, cabbage, cauliflower, and kohlrabi.

Allow 1 day between application and harvest of strawberries. Allow 7 days between dusting and harvest of blackberries, boysenberries, dewberries, loganberries and raspberries.

Last application either on birds or in premises; 7 days before slaughter. Do not treat nest litter. Avoid direct contamination of eggs. Do not feed treated forage to livestock.

NON-WARRANTY: Follow directions carefully. Timing and method of application, weather and crop conditions, mixtures with other chemicals not specifically recommended, and other influenceing factors in the use of this product are beyond the control of the seller. Buyer assumes all risks of use, storage or handling of this material not in strict accordance with directions given herewith.

USDA Reg. No. 10159-2

VOLUNTARY PURCHASING GROUPS, INC.
BONHAM, TEXAS

