

Feb 2 1972

1457-2



5% SEVIN* DUST

ACTIVE INGREDIENTS:

Carbaryl (1-naphthyl N-methylcarbamate) 5.00%

INERT INGREDIENTS: 95.00%

Total 100.00%

CAUTION

KEEP OUT OF REACH OF CHILDREN

SEE SIDE PANEL FOR ADDITIONAL CAUTIONS

*Registered trademark of Union Carbide Corporation for the active ingredient 1-naphthyl N-methylcarbamate.

NET WEIGHT 4 POUNDS

Continued from back

CABBAGE, BROCCOLI, BRUSSELS SPROUTS, CAULIFLOWER, KOHLRABI—to control flea beetles, harlequin bug, armyworms, cabbage caterpillars, corn earworm. To control small cabbage loopers, use a 5- to 7-day schedule.

STRAWBERRIES—to control meadow spittlebug, strawberry leaf roller and strawberry weevil.

BLACKBERRIES, BOYSENBERRIES, DEWBERRIES, LOGANBERRIES AND RASPBERRIES—to control Japanese beetles, leaf rollers, leafhoppers and raspberry aphids.

APPLY 5% SEVIN DUST AS OFTEN AS NECESSARY TO CONTROL GRASSHOPPERS ON THE ABOVE CROPS.

5% SEVIN DUST can be safely used on a wide range of ornamental plants, flowers, shrubs and small trees, such as rose, carnation, gladiolus, chrysanthemum, lilac, arborvitae, juniper, Hydrangea, azalea, elm, maple, oak, dogwood, birch and pines. SEVIN injures Boston Ivy.

Apply to the upper and lower leaf surface of all foliage for complete coverage to control such pests as:

Apple aphid, June Beetles, leafhoppers, leaf roller, boxwood leaf miner, mimosa webworm, orange tortrix, willow leaf beetles, Japanese beetle, psyllids, rose aphids, boxelder bug, scale insects, oak leaf miners, thrips (exposes), Plant Bugs, lace bugs, rose slug, mealy bugs, elm leaf aphid, flea beetles, gypsy moth, Bagworm, puss caterpillars, birch leaf miner, blister beetles, thorn bugs, elm leaf beetle, periodical cicada.

SEVIN does not control spider mites. Where infestations are encountered, use a recommended miticide.

Apply 5% SEVIN DUST when insects or their damage appear. Make applications in the morning or evening when it is cool. Treat upper and lower surface of all foliage for complete coverage. Repeat as necessary, generally at 7-day intervals. For best results, apply with a dust gun. Do not apply in excess.

Continued on other side

Continued from other side

CAUTIONS: Harmful if inhaled or swallowed. Do not breathe dust. Do not take internally. Avoid skin contact. Wash thoroughly after using. Avoid contamination of feed or foodstuffs. Note for physician: Carbaryl is a moderate, reversible, Cholinesterase inhibitor. Atropine is antidotal.

This product is highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your Cooperative Agricultural Extension Service.

Do not apply within 3 days—to beans, corn foliage or fodder, cucumbers, eggplants, melons, okra, garden peas, peppers, potatoes, pumpkins, summer squash, broccoli, brussels sprouts, cabbage, cauliflower and kohlrabi.

Allow 1 day between application and harvest of strawberries. Allow 7 days between dusting and harvest of blackberries, boysenberries, dewberries, loganberries and raspberries.

Last application either on birds or in premises, 7 days before slaughter. Do not treat nest litter. Avoid direct contamination of eggs. Do not feed treated forage to livestock.

Do not reuse empty container. Destroy it by burying with waste or burning. Stay away from smoke or fumes.

NON-WARRANTY: Follow directions carefully. Timing and method of application, weather and crop conditions, mixtures with other chemicals not specifically recommended, and other influencing factors in the use of this product are beyond the control of the seller. Buyer assumes all risks of use, storage or handling of this material not in strict accordance with directions given herewith.



1. WHAT IS THIS PACKAGE FOR?
This package of VPG 5% SEVIN control many insect pests as listed.

2. HOW DO YOU USE IT?
ON BIRDS—to control Northern 1 per 100 birds.

ON LITTER—to control Northern 1 lb. 5% DUST per 40 sq. ft.

IN DUST BATH BOXES—to control use 2 1/2 lbs. 5% DUST per box.

ON DOGS AND CATS—for tick avoiding the eyes, and rub thorax after treatment. Repeat at weekly puppies under 4 weeks old. In air dust liberally in and around sleep a shaker can, hand or dust gun.

BEANS (green beans, lima beans) control Mexican bean beetle, Japanese beetle, leafhoppers, worms, corn earworm, lima bean

SWEET CORN—to control corn ear fall armyworm, flea beetles, Japanese larvae in whorl (budworm damage as necessary).

For insects attacking silks and ears silks appear, and continuing until required depending on severity of for effective control. Application pollen shed period will seriously

CUCUMBER, MELONS, PUMPKIN cucumber beetles, flea beetles, leafhoppers. Some leaf injury may occur in the presence of prolonged high

POTATO, TOMATO, EGGPLANT beetle, flea beetles, leafhoppers, tomato fruit worm, tomato hornworm

OKRA—to control corn earworm. EPA Reg. No. 10159-2

VOLUNTARY BUY

Feb 2 1972

1457-2

Continued from other side

CAUTIONS: Harmful if inhaled or swallowed. Do not breathe dust. Do not take internally. Avoid skin contact. Wash thoroughly after using. Avoid contamination of feed or foodstuffs. Note for physician: Carbaryl is a moderate, reversible, Cholinesterase inhibitor. Atropine is antidotal.

This product is highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your Cooperative Agricultural Extension Service.

Do not apply within 3 days—to beans, corn foliage or fodder, cucumbers, eggplants, melons, okra, garden peas, peppers, potatoes, pumpkins, summer squash, broccoli, brussels sprouts, cabbage, cauliflower and kohlrabi.

Allow 1 day between application and harvest of strawberries. Allow 7 days between dusting and harvest of blackberries, boysenberries, dewberries, loganberries and raspberries.

Last application either on birds or in premises: 7 days before slaughter. Do not treat nest litter. Avoid direct contamination of eggs. Do not feed treated forage to livestock.

Do not reuse empty container. Destroy it by burying with waste or burning. Stay away from smoke or fumes.

NON-WARRANTY: Follow directions carefully. Timing and method of application, weather and crop conditions, mixtures with other chemicals not specifically recommended, and other influencing factors in the use of this product are beyond the control of the seller. Buyer assumes all risks of use, storage or handling of this material not in strict accordance with directions given herewith.

VPG 5% SEVIN DUST

1. WHAT IS THIS PACKAGE FOR?

This package of VPG 5% SEVIN DUST provides an economical and easy method to control many insect pests as listed in the directions for use.

2. HOW DO YOU USE IT?

ON BIRDS—to control Northern fowl mite, chicken mite, lice, fleas, use 1 lb. 5% DUST per 100 birds.

ON LITTER—to control Northern fowl mite, chicken mite, lice, fleas, bedbugs, use 1 lb. 5% DUST per 40 sq. ft.

IN DUST BATH BOXES—to control Northern fowl mite, chicken mite, lice and fleas, use 2 1/2 lbs. 5% DUST per box for each 50 birds.

ON DOGS AND CATS—for ticks and fleas dust liberally over the entire animal, avoiding the eyes, and rub thoroughly into the skin. Comb out dead ticks a few hours after treatment. Repeat at weekly intervals if needed. Avoid treatment of kittens and puppies under 4 weeks old. In and around Dog and Cat Quarters: For ticks and fleas dust liberally in and around sleeping quarters, kennels, and dog houses by means of a shaker can, hand or dust gun.

BEANS (green beans, lima beans, snap beans, cowpeas, black-eyed peas)—to control Mexican bean beetle, bean leaf beetle, cucumber beetles, flea beetles, Japanese beetle, leafhoppers, velvetbean caterpillar, western bean cutworm, armyworms, corn earworm, lima bean pod borer, stink bugs and tarnished plant bugs.

SWEET CORN—to control corn earworm, corn rootworm adults, European corn borer, fall armyworm, flea beetles, Japanese beetles, saw beetles and leaf hoppers. For larvae in whorl (budworm damage) and foliage feeders, apply to entire plant. Repeat as necessary.

For insects attacking silks and ears, apply at 2 to 3-day intervals starting when first silks appear, and continuing until silks begin to dry. Three or more applications may be required depending on severity of infestation. Timing and good coverage are essential for effective control. Application of this product to tassel region of corn during the pollen shed period will seriously reduce bee populations.

CUCUMBER, MELONS, PUMPKIN, SQUASH—to control pickleworm, melonworm, cucumber beetles, flea beetles, leafhoppers and squash bugs. Avoid excessive applications. Some leaf injury may occur if treatments are made when tender foliage is wet or in the presence of prolonged high humidity. SEVIN injures watermelons in Florida.

POTATO, TOMATO, EGGPLANT, PEPPER, PEAS—to control Colorado Potato beetle, flea beetles, leafhoppers, European corn borer, fall armyworms, lace bugs, tomato fruit worm, tomato hornworm, tarnished plant bugs, stink bugs.

OKRA—to control corn earworm and stink bugs.

EPA Reg. No. 10159-2

Instruction Cont'd on Side

VOLUNTARY PURCHASING GROUPS INC.