

ACCEPTED  
 3-9-72  
 UNDER THE FEDERAL INSECTICIDE  
 FUNGICIDE AND RODENTICIDE ACT  
 FOR ECONOMIC POISON RESISTANCE  
 ED UNDER NO. 10129-3

**Hi-Yield.**

**3 A.G.**

**MSMA+  
 SURFACTANT**

**FOR SELECTIVE POST-EMERGENT WEED CONTROL**

<b>Active Ingredient:</b>		
<b>Monosodium Acid Methanearsonate</b>	.....	<b>47.6%</b>
<b>Inert Ingredients:</b>	.....	<b>52.4%</b>
		<hr/>
	<b>Total</b>	<b>100%</b>

Total arsenic (as elemental) all in water soluble form 22.1%  
 Product contains 6 pounds of Monosodium Acid Methanearsonate  
 per gallon.  
 Use when temperature is at least 70° F.

**CAUTION** KEEP OUT OF REACH OF CHILDREN.  
 See Side Panel for Additional Cautions.

**VOLUNTARY PURCHASING GROUPS, INC.  
 BONHAM, TEXAS**

**Net Contents  
 1 Gallon**

**GENERAL INFORMATION:**

HI-YIELD 3 A. G. is a new highly effective, selective type post-emergence weed killer formulated particularly for grassy weeds. It contains surfactant — all you need is water. Best results are obtained on young, actively growing weeds when temperature is at least 70° F.

**GENERAL WEED AND GRASS CONTROL:**

HI-YIELD 3 A. G. is found very effective for controlling Johnson Grass, Goose Grass and Yellow Foxtail along fence lines, implement storage yards, and similar non-crop areas when cultivation is not practical.

**RECOMMENDED RATES FOR GENERAL WEED CONTROL:**

Mix 3 to 6 quarts of HI-YIELD 3 A. G. in 100 gallons of water. Apply 50 gallons of spray solution per acre, thoroughly wetting foliage and stems of weeds to be killed. Spot treatments to control re-growth may be necessary.

**COTTON:**

HI-YIELD 3 A. G. is very useful as a directed application for the post-emergence control of Johnson Grass, Goose Grass, Water Grass, Cocklebur, Nutgrass, Dallis Grass, and similar weeds in cotton. It should be applied at the rate of 1/4 gallon per acre.

Mix 3 quarts of HI-YIELD 3 A. G. in 90 gallons of water. Do not add any surfactant to the spray solution. Apply spray solution at 40 gallons per acre. Weed control is most effective at temperatures above 70° F. For band application in 40 inch rows, apply 1 gallon of spray solution per inch of band width per acre. Direct the spray solution to the base of the cotton plant. Nozzles should be placed so as to avoid spraying the cotton foliage. A second application may be required during

humid growing conditions to control perennials. Apply the second applications 1 to 3 weeks later.

Slight burning and a reddish discoloration of the cotton leaf may occasionally occur following the recommended treatment, but the cotton plant will develop normally. Clean application equipment after each use by flushing with water.

**WASTE DISPOSAL:** Rinse spray equipment and containers and dispose of liquid wastes in pit in non-crop lands located away from water supplies.

**DO NOT APPLY AFTER FIRST BLOOM. DO NOT GRAZE OR FEED FORAGE FROM TREATED AREAS TO LIVESTOCK. DO NOT RE-USE CONTAINER FOR ANY PURPOSE, DESTROY WHEN EMPTY.**

**CAUTION:** Harmful if swallowed. Avoid contact with skin. Wash hands after using. Avoid breathing spray mist. Avoid contamination of feed and foodstuffs. Avoid storage near food or feed products. Keep away from children and domestic animals. Keep children and animals off treated areas until washed into soil.

Do not contaminate water used by wildlife or aquatic life or water used for domestic or irrigation purposes.

**ANTIDOTE:** If taken internally, induce vomiting and call a physician at once.

**NON-WARRANTY:** Follow directions carefully. Timing and method of application, weather and crop conditions, mixtures with other chemicals not specifically recommended, and other influencing factors in the use of this product are beyond the control of the seller. Buyer assumes all risks of use, storage or handling of this material not in strict accordance with directions given herewith.

EPA Reg. No. 10159-3

**DO NOT USE OR STORE IN OR AROUND THE HOME  
DO NOT ALLOW DOMESTIC ANIMALS TO GRAZE TREATED AREA**

ACCEPTED

3-9-72

UNDER THE FEDERAL INSECTICIDE  
ACT OF 1947  
EPA REG. NO. 10159-3

# S H E F F

## NO-GRO LIQUID CONCENTRATE FOR PROFESSIONAL USE ONLY

For controlling grass and weeds in sand traps on golf courses

**DANGER** POISON: KEEP OUT OF REACH OF CHILDREN. KEEP CHILDREN AND ANIMALS OFF TREATED AREA. DO NOT USE OR STORE IN OR AROUND THE HOME. SEE OTHER PRECAUTIONS ON SIDE PANEL.

THIS PRODUCT IS TOXIC TO WILDLIFE. APPLY THIS PRODUCT ONLY AS SPECIFIED ON THE LABEL. DO NOT CONTAMINATE WATER BY CLEANING OF EQUIPMENT, OR DISPOSAL OF WASTES.

Active Ingredient  
 Arsenic Trioxide (30.7%) per gallon  
 Expressed as metallic all in water soluble form: 23.3%

40.4%  
59.6%

**CAUTION: POISON.** May be fatal if swallowed. Avoid contact with skin and eyes. Wash thoroughly with soap and water. Do not use in any connection with feed, food or drinking water. If irritation or burns persist, flush with vinegar — see a physician immediately.

**WARNING:** Spraying operator should wear protective gloves and a respirator mask. After spraying, operator should take a shower, washing thoroughly with soap and water, and change clean clothing. Never wear clothing that has been used in spraying operation before laundering. If spilled on skin or eyes, flush off immediately with water. Keep children and pets away from treated areas. Rinse container several times with water. Do not use in any connection with feed, food or drinking water. If irritation or burns persist, flush with vinegar — see a physician immediately.



**DANGER**  
 Contains ARSENIC  
 Contains CAUSTIC  
 POISON  
 See ANTIDOTE

**ANTIDOTE:** External: First Aid treatment. If skin irritation or burns should appear, flush areas with vinegar, dress with a wet dressing of milk of magnesia. If healing does not begin within 24 hours, see a Physician. If large areas of the skin should be irritated or blistered, see a physician immediately. Treat as for caustic burns.

Internal antidote. Call Physician at once. Give a tablespoon of salt in a glass of warm water until vomit is clear. Then give 2 tablespoons of epsom salts or milk of magnesia in water, followed by plenty of milk and water. Have patient lie down and quiet him.

PLEASE READ THIS ENTIRE LABEL BEFORE USING

**ACCEPTED**  
 10/19/68  
 UNDER THE FEDERAL INSECTICIDE FUNGICIDE AND RODENTICIDE ACT FOR ECONOMIC PEST CONTROL REGISTERED UNDER FIFED 25.2 SUBJECT TO ATTACHED COMMENTS.

**DIRECTIONS** for using NO-GRO in sand traps.  
 Treat sand traps at the rate of one gallon of NO-GRO in 15 gallons of water per 1,000 square feet. Sand should be dry when treated. Do not treat if water is standing in trap or if sand is wet. Careful application will give control of grass and weeds for months. Retreat when regrowth appears.  
**Caution:** this is a soil sterilizing solution and must be applied only IN the sand trap. Do not apply to turf. Do not apply to areas underlaid by roots of desirable trees and shrubs.

NET CONTENTS                      gallons.    USDA reg. no. 10193-1

DO NOT REUSE CONTAINER.                      DESTROY WHEN EMPTY.

**SHEFF CHEMICAL & SUPPLY COMPANY**

PHONE 746-5555    2827 RIVERVIEW BLVD. W.  
BRADENTON, FLORIDA 33505