

P R O K I L

Methyl Parathion-4

An emulsible solution containing 4-lbs. methyl parathion per gallon, for use in the control of certain insects as specified.

COMPOSITION

ACTIVE INGREDIENTS:

*O, O-dimethyl O-p-nitrophenyl thiophosphate	45.8%
Aromatic petroleum derivatives	49.1%
INERT INGREDIENTS	5.1%
Total	100.0%

*Methyl Parathion



DANGER POISON



KEEP OUT OF REACH OF CHILDREN

ATROPINE IS ANTIDOTAL

Keep an emergency supply on hand
CALL A PHYSICIAN AT ONCE

SEE ANTIDOTE STATEMENT AND OTHER WARNINGS
ON THE BACK - SIDE PANEL

NET CONTENTS _____ U.S. GALLONS

— DIRECTIONS FOR USE —

FROKIL METHYL PARATHION-4 is an emulsible solution containing 4 pound methyl parathion per gallon, for use in the control of certain insects and mites as specified. This product is for use only by thoroughly trained operators familiar with the handling of hazardous insecticides and is to be applied only by power sprayers or by aircraft. When applied use sufficient water for good coverage: By Aircraft 3-10 gallons per acre; and by ground equipment 10 or more gallons per acre.

READ ENTIRE LABEL. Use strictly in accordance with dangers, cautions, warnings, and directions, and with applicable State and Federal regulations. Timing and dosage vary greatly between districts, as a consequence of local weather or conditions. Always follow the recommendations of State Agricultural Extension Service concerning specific usage.

ALFALFA: Spotted alfalfa aphid—Apply 3/4 pint per acre in sufficient water for good coverage. DO NOT apply within 15 days of cutting or pasturing.

SEED ALFALFA: Lygus bugs — Apply 1 1/2 to 2 1/2 pints per acre in sufficient water to cover. DO NOT apply within 20 days of harvest.

BEETS: Armyworms (up to 3rd instar), Leafhoppers, Red spider mites — Apply 1 to 2 pints per acre in sufficient water to cover. DO NOT apply within 15 days of harvest or within 21 days if tops are to be used for feed or food.

BROCCOLI, BRUSSEL SPROUTS, CABBAGE. Aphids, Armyworms (up to 3rd instar), Fleabeetles, Loopers (immature), Thrips, Red spider mites—Apply 2 to 3 pints per acre to cover. Loopers (immature)—Apply 1 3/5 to 2 pints per acre in sufficient water to cover. DO NOT apply within 21 days of harvest.

SUGAR BEETS: Armyworms (up to 3rd instar), Fleabeetles, Red spider mites, Leafhoppers—Apply 1/2 pint per acre in sufficient water to cover. DO NOT apply within 20 days of harvest.

COTTON: Stink bugs, Lygus bugs, Red spider mites—Apply 1 to 2 pints per acre in sufficient water to cover. DO NOT apply within 15 days of harvest. Workers entering treated fields within 24 hours after application should wear protective clothing.

CARROTS: Aphids—Apply 1 to 2 pints per acre in sufficient water to cover. DO NOT apply within 15 days of harvest.

CELERY, PEAS, PEPPERS, LETTUCE, SPINACH: Aphids, Armyworms (up to 3rd instar), Fleabeetles, Leafhoppers, Red Spider Mites — Apply 1 to 2 pints per acre in sufficient water to cover. DO NOT APPLY to Celery, Peas, and Peppers within 15 days of harvest. DO NOT APPLY to Lettuce and Spinach within 21 days of harvest.

FROKIL METHYL PARATHION-4 IS COMPATIBLE WITH THE FOLLOWING: Pyrethrins, Oils, Dinitros, DDT, DDD, Methoxychlor, Perthane, BHC, Aldrin, Dieldrin, Endrin, Chlordane, Heptachlor, Toxaphene, Tedion, Thiodan, Kelthane, Chlorobenzilate, Malathion, Guthion, Phosdrin, Dibrom, Bidrin, Phosphamidon, Vapona, Cygon, Dylor, Trithion, Sevin; Diazinon, Delnav, Parathion.

USDA Reg. No. 10163-2

FIRST AID TREATMENT — ANTIDOTE

Do not wait for a doctor. Give 2 Atropine tablets at once if patient has blurred vision or pin-point pupils.

INTERNAL: If swallowed, give tablespoon of salt in warm water and repeat until vomit fluid is clear.

EXTERNAL: If spilled on body, immediately remove contaminated clothing and wash skin thoroughly with soap and water.

INHALATION: Remove patient from contaminated area and keep under continuous observation until doctor arrives.

DANGER

POISONOUS IF SWALLOWED, INHALED OR ABSORBED THROUGH THE SKIN. Repeated exposures may be increasingly hazardous. RAPIDLY ABSORBED THROUGH THE SKIN. DO NOT BREATHE VAPORS. DO NOT GET INTO EYES, OR SKIN OR ON CLOTHING. If spilled on skin, immediately wash thoroughly with soap and water. Wear protective clothing, rubber gloves and goggles. When filling spray tanks and during mixing and application, wear a mask respirator passed by the U. S. Department of Agriculture for methyl parathion protection. Airplane pilots should wear full-face cannister-type masks.

DO NOT STORE NEAR NOR CONTAMINATE FEED OR FOODSTUFFS. Wash hands, arms and face thoroughly with soap and water before eating or smoking. Bathe promptly with soap and water after spraying. Wash clothing with soap and hot water daily, before reuse.

CAUTION: Workers who enter treated fields within 24 hours after treatment must wear protective clothing. Keep all persons out of the operating areas or vicinity where there may be danger of drift. Vacated areas should not be re-entered until insecticidal residues have dissipated.

THIS PRODUCT IS TOXIC TO WILDLIFE. Birds and other wildlife in treated areas may be killed. Keep out of any body of water. This product is highly toxic to bees exposed to treatment or residues on crops. Protective information may be obtained from your Cooperative Agricultural Extension Service. Do not apply when weather conditions favor drift from areas treated. Apply this product only as specified on this label.

DESTROY THIS CONTAINER WHEN EMPTY: Wash out thoroughly with water. Punch drum full of holes or crush so that it cannot be used for any purpose. Dispose of it where it cannot be salvaged for any use.

NOTE: Many pesticidal chemicals are poisonous and may leave a toxic residue on the plants to which they are applied. The U.S. Food and Drug Administration has established minimum amounts of such pesticidal chemicals that may remain on raw agricultural products at harvest, and it is the user's responsibility to see that there is no residue on such crops at harvest in excess of these amounts.

The "Directions For Use" at left, are based on the best available information, and if followed carefully should not leave excessive residues at harvest.

CALIFORNIA (Plant)
340 East Main St.
P. O. Box 406
Calipatria, Calif. 92233
Telephone: 348-5150

The Dune Company

AGRICULTURAL CHEMICALS

ARIZONA

Rt. 1, Box 33

Somerton, Arizona 85350

Telephone: 627-2479

10163-2

10-23-69

101-2

ACCEPTED

OCT 23 1969

UN. GENERAL INSECTICIDE
FUNGICIDE AND RODENTICIDE ACT
FOR ECONOMIC POISON REGISTERED
UNDER NO. 10163-2 SUBJECT
TO ATTACHED COMMENTS.

File Symbol 10163-RN

2012