

SUGGESTED USE

READ ENTIRE LABEL. USE STRICTLY IN ACCORDANCE WITH WARNINGS AND DIRECTIONS, AND WITH APPLICABLE STATE AND FEDERAL REGULATIONS. TIMING AND DOSAGE VARY GREATLY BETWEEN DISTRICTS, AS A CONSEQUENCE OF LOCAL WEATHER AND OTHER CONDITIONS. ALWAYS FOLLOW THE RECOMMENDATIONS OF STATE AGRICULTURAL EXPERIMENT STATION CONCERNING SPECIFIC USAGE.

DIRECTIONS

DO NOT RE-USE EMPTY DRUM. RETURN TO DRUM RECONDITIONER OR DESTROY BY PERFORATING OR CRUSHING AND BURYING IN A SAFE PLACE.

Prokil Perthane 4 E C is for use only by thoroughly trained operators familiar with the handling of hazardous insecticides and is to be applied only by powered ground sprayers in 10 or more gallons water per acre or by aircraft in 3-10 gallons water per acre.

U.S.D.A. Reg. No. 10163 S

PROKIL PERTHANE⁺ 4

ACTIVE INGREDIENTS	BY WT.
Diethyl Diphenyl Dichloroethane	46.30%
Related Reactive Compounds	2.50%
Aromatic Petroleum Derivatives	47.20%
INERT INGREDIENTS	4.00%
TOTAL	100.00%

Perthane is a trademark Reg. U.S. Patent Office by Rohm & Haas Company.

Contains 4 lbs. Perthane per gallon.

WARNING

KEEP OUT OF REACH OF CHILDREN

See side panel for additional precautions.

NET CONTENTS

U. S. GALLONS

CALIFORNIA (Plant)
340 East Main St.,
P. O. Box 406
Telephone: 348-5150
Calipatria, Calif. 92233

The Dune Company

AGRICULTURAL CHEMICALS

ARIZONA
Rt. 1, Box 33
Somerton, Ariz. 85050
Telephone: 627-2479

BROCCOLI, E
ER, KOHLRA
Cabbage W
not apply wi
Sprouts, Cab
7 days of ho

LETTUCE: C
2 quarts per
vest.

Harmful if s
or eyes.

Avoid breath
with skin.


V/as. iha

Do not conta

AVOIL CON

Do not s

or feed crop



OKIL PERTHANE* 4

	BY WT.
Dichloroethane	46.30%
Compounds	2.50%
in Derivatives	47.20%
.....	4.00%
.....	100.00%

mark Reg. U.S. Patent
Haas Company.

lbs. Perthane per gallon.

WARNING
OF REACH OF CHILDREN
Additional precautions.

5 U. S. GALLONS

Haas Company
AGRICULTURAL CHEMICALS

ARIZONA
Rt. 1, Box 33
Somerton, Ariz. 85350
Telephone: 627-2479

BROCCOLI, BRUSSELS SPROUTS, CABBAGE, CAULIFLOWER, KOHLRABI, SPINACH: Cabbage Looper, Imported Cabbage Worm, Leafhopper: 1 quart per acre. Do not apply within 3 days of harvest on Broccoli, Brussels Sprouts, Cabbage, Cauliflower or Kohlrabi or within 7 days of harvest on Spinach.

LETTUCE: Cabbage Looper, Imported Cabbage Worm, 2 quarts per acre. Do not apply within 4 day of harvest.

WARNING

Harmful if swallowed, inhaled or in contact with skin or eyes.

Avoid breathing of spray or spray mist. Avoid contact with skin.

Wash thoroughly after using.

Do not contaminate or store near feed or foodstuffs.

AVOID CONTAMINATION OF FEED AND FOODSTUFFS.

Do not allow product to drift onto adjoining food or feed crops as residues may contaminate these crops

making them unfit for sale, feed, or human consumption. Keep clear of food or dairy animals, and streams.

HIGHLY TOXIC TO FISH AND WILDLIFE

Birds feeding on treated areas may be killed. Do not apply when weather conditions favor drift from areas treated. Do not apply where runoff is likely to occur.

Do not contaminate any body of water, by direct application, cleaning of equipment, or disposals of wastes and containers. Do not apply this product to any area or to any crop not specifically indicated on this label.

This product is toxic to bees and should not be applied when bees are actively visiting the area.

NOTICE ON CONDITIONS OF SALE

Our recommendations for use of this product are based upon tests believed to be reliable. The use of this product being beyond the control of the manufacturer, no guarantee, expressed or implied, is made as to the effects or such or the results to be obtained if not used in accordance with directions or established safe practice. The buyer must assume all responsibility including injury or damage resulting from its misuse as such or in combination with other materials.