

# PROKIL METHYL PARATHION 5

## — DIRECTIONS FOR USE —

PROKIL METHYL PARATHION-5 is an emulsible solution containing 5 pounds methyl parathion per gallon, for use in the control of certain insects and mites as specified. This product is for use only by thoroughly trained operators familiar with the handling of hazardous insecticides and is to be applied only by power sprayers or by aircraft. When applied use sufficient water for good coverage: By Aircraft 5-10 gallons per acre; and by ground equipment 20 or more gallons per acre.

**READ ENTIRE LABEL.** Use strictly in accordance with dangers, cautions, warnings, and directions, and with applicable State and Federal Regulations. Timing and dosage vary greatly between districts, as a consequence of local weather or conditions. Always follow the recommendations of State Agricultural Extension Service concerning specific usage.

**ALFALFA:** Spotted alfalfa aphid—Apply  $\frac{3}{5}$  pint per acre in sufficient water for good coverage. DO NOT APPLY within 15 days of cutting or pasturing.

**SEED ALFALFA:** Lygus bugs—Apply 1 to 2 pints per acre in sufficient water to cover. DO NOT apply within 20 days of harvest.

**BEANS, DRY:** Aphids, Armyworms (up to 3rd instar), Flea Beetles, Mites — Apply  $1\frac{3}{5}$  -  $2\frac{2}{5}$  pints per acre. DO NOT apply within 15 days of harvest.

**BEANS, GREEN:** Aphids, Armyworms (up to 3rd instar), Flea Beetles, Mites— Apply  $1\frac{3}{5}$  -  $2\frac{2}{5}$  pints per acre. DO NOT apply within 21 days of harvest.

**BEETS:** Armyworms (up to 3rd instar), Leafhoppers, Red spider mites—Apply  $\frac{3}{4}$  to  $1\frac{1}{2}$  pints per acre in sufficient water to cover. DO NOT apply within 15 days of harvest or within 21 days if tops are to be used for feed or food.

**BROCCOLI, BRUSSEL SPROUTS, CABBAGE:** Aphids, Armyworms (up to 3rd instar), Fleabeetles, Thrips, Red spider mites—Apply 1 to 2 pints per acre in sufficient water to cover. Loopers (Immature) — Apply  $1\frac{3}{5}$  to 2 pints per acre in sufficient water to cover. DO NOT apply within 21 day of harvest.


**SUGAR BEETS:** Armyworms (up to 3rd instar), Fleabeetles, Red spider mites, Leafhoppers—Apply 0.6 pint per acre in sufficient water to cover. DO NOT apply within 20 days of harvest. DO NOT treat within 60 days of harvest if tops are to be fed to Livestock.

**COTTON:** Stink bugs, Lygus bugs, Red spider mites—Apply  $\frac{3}{4}$  to 2 pints per acre in sufficient water to cover. DO NOT apply within 15 days of harvest. Workers entering treated fields within 24 hours after applications should wear protective clothing.

**CARROTS:** Aphids—Apply  $\frac{3}{4}$  to  $1\frac{1}{2}$  pints per acre in sufficient water to cover. DO NOT apply within 15 days of harvest or use treated tops for food or feed.

**CELERY, PEAS, PEPPERS, LETTUCE, SPINACH:** Aphids, Armyworms (up to 3rd instar), Fleabeetles, Leafhoppers, Red spider mites - Apply  $\frac{3}{4}$  to  $1\frac{1}{2}$  pints per acre in sufficient water to cover. DO NOT APPLY to Celery, Peas, Peppers within 15 days of harvest. DO NOT APPLY to Lettuce and Spinach within 21 days of harvest.

**TOMATOES:** Aphids, Armyworms (up to 3rd instar), Fleabeetles — Apply 0.8 to 2.4 pts. per acre. DO NOT apply within 10 days of harvest when



Apply 1 3/5 - 2 2/5 pints per acre. DO NOT apply within 15 days of harvest.

**BEANS, GREEN:** Aphids, Armyworms (up to 3rd instar), Flea Beetles, Mites—Apply 1 3/5 - 2 2/5 pints per acre. DO NOT apply within 21 days of harvest.

**BEETS:** Armyworms (up to 3rd instar), Leafhoppers, Red spider mites—Apply 3/4 to 1 1/2 pints per acre in sufficient water to cover. DO NOT apply within 15 days of harvest or within 21 days if tops are to be used for feed or food.

**BROCCOLI, BRUSSEL SPROUTS, CABBAGE:** Aphids, Armyworms (up to 3rd instar), Fleabeetles, Thrips, Red spider mites—Apply 1 to 2 pints per acre in sufficient water to cover. Loopers (Immature) — Apply 1 3/5 to 2 pints per acre in sufficient water to cover. DO NOT apply within 21 day of harvest.

**SUGAR BEETS:** Armyworms (up to 3rd instar), Fleabeetles, Red spider mites, Leafhoppers—Apply 0.6 pint per acre in sufficient water to cover. DO NOT apply within 20 days of harvest. DO NOT treat within 60 days of harvest if tops are to be fed to Livestock.

**COTTON:** Stink bugs, Lygus bugs, Red spider mites—Apply 3/4 to 2 pints per acre in sufficient water to cover. DO NOT apply within 15 days of harvest. Workers entering treated fields within 24 hours after applications should wear protective clothing.

**CARROTS:** Aphids—Apply 3/4 to 1 1/2 pints per acre in sufficient water to cover. DO NOT apply within 15 days of harvest or use treated tops for food or feed.

**CELERY, PEAS, PEPPERS, LETTUCE, SPINACH:** Aphids, Armyworms (up to 3rd instar), Fleabeetles, Leafhoppers, Red spider mites - Apply 3/4 to 1 1/2 pints per acre in sufficient water to cover. DO NOT APPLY to Celery, Peas, Peppers within 15 days of harvest. DO NOT APPLY to Lettuce and Spinach within 21 days of harvest.

**TOMATOES:** Aphids, Armyworms (up to 3rd instar), Fleabeetles. Apply 0.8 to 2.4 pts. per acre. DO NOT apply within 10 days of harvest when applying 0.8 pts. DO NOT apply within 15 days of harvest when applying more than 0.8 pts. per acre.

PROKIL METHYL PARATHION-5 IS COMPATIBLE WITH THE FOLLOWING: Pyrethrum, Oils, Dinitros, Methoxychlor, Perthane, Toxaphene, Tedion, Thiodan, Kelthane, Chlorobenzilate, Malathion, Guthion, Phosdrin, Dibrom, Bidrin, Phosphamidon, Vapona, Cygon, Dilox, Ethion, Trithion, Delnav, Diazinon, Sevin

**DESTROY THIS CONTAINER WHEN EMPTY:** Wash out thoroughly with water. Punch drum full of holes or crush so that it cannot be used for any purpose. Dispose of it where it cannot be salvages for any use. Do not contaminate any body of water nor apply to any area not specified on the label.

**THIS PRODUCT IS HIGHLY TOXIC TO BEES** exposed to direct treatment or residues on crops. Protective information may be obtained from your Cooperative Agricultural Extension Service.

**FLAMMABLE:** Do not use or store near heat or open flame. Keep container closed.

#### NOTICE ON CONDITIONS OF SALE

Our recommendations for use of this product are based upon tests believed to be reliable. The use of this product being beyond the control of the manufacturer no guarantee, expressed or implied, is made as to the effects of such or the results to be obtained if not used in accordance with directions or established safe practice. The buyer must assume all responsibility, including injury or damage, resulting from its misuse as such, or in combination with other materials.

1-74 106 - PROOF

EPA Reg. No. 10162-7 AA

ACCEPTED

10/16/3-17