

P R O K I L

SUGGESTED USE

READ ENTIRE LABEL. USE STRICTLY IN ACCORDANCE WITH CAUTIONS, WARNINGS AND DIRECTIONS AND WITH APPLICABLE STATE AND FEDERAL REGULATIONS. TIMING AND DOSAGE MAY VARY AS A CONSEQUENCE OF LOCAL WEATHER OR CONDITIONS. WE RECOMMEND REFERENCE TO AGRICULTURAL EXPERIMENT STATIONS ON SPECIFIC USAGE.

Apply thoroughly. Do not apply to food crops within 30 days of harvest and to other crops intended for use remaining in harvest by washing, brushing or other effective means.

CUCUMBERS, SQUASH, MELONS: Aphids, Spider Mites, Pickleworm, Cucumber Beetles, Flea Beetles; **CANTALOUPE** — Leafhoppers — 30 to 35 lbs. per acre. Repeat as necessary. **SQUASH:** Squash Vine Borers — 30 lbs. per acre. **PUMPKINS:** Squash Vine Borers — 35 lbs. per acre. Apply twice a week to stems and vines at base of plants. Do not apply to Pumpkins within 3 days of harvest, and to Cucumbers, Cantaloupes, Squash and Melons within 1 day of harvest. Do not apply unless plants are dry.

Malathion - Cryolite 4-35 Dust

Active Ingredient	By Weight
Malathion	4%
Sodium Fluoroborate	35%
Inert Ingredients	61%
100% contains 4.35 lbs. per 100 lbs. of diethyl mercaptophosphate	
Net Weight	19.0

READ THE LABEL
KEEP OUT OF REACH OF CHILDREN
CAUTION

Avoid prolonged breathing of dust.

Avoid repeated or prolonged contact with skin, wash thoroughly after use. Avoid contamination of feed and food. Do not apply under conditions involving possible drift to food, forage or other plantings that might be damaged or the crop thereof rendered unfit for sale, use or consumption.

SEE RIGHT PANEL FOR ADDITIONAL CAUTIONS!

NET WEIGHT 50 LBS.

LETTUCE: Aphids, Mites, Cabbage Looper, Leafhoppers, Climbing Cutworms, Alfalfa Looper, Corn Earworms — 20 to 30 lbs. per acre. Do not apply after the seedling stage.

TOMATOES: Aphids, Spider Mites, Tomato Russet Mites, Drosophila, Armyworms, Flea Beetles — 35 to 45 lbs. per acre. Do not apply within 1 day of harvest.

WARNING

This product will kill fish. Keep out of any body of water. This product is highly toxic to bees exposed to treatment or residues on crops. Protective information may be obtained from your Cooperative Agricultural Extension Service.

NOTICE ON CONDITIONS OF SALE

The Dune Company makes No Warranty, expressed or implied concerning the use of this product other than indicated on the label. Buyer assumes all risk of use and/or handling of this material when such use and/or handling is contrary to label instructions.

U.S.D.A. Reg. No. 10163-14

CALIFORNIA (Plant)
P. O. Box 406
340 East Main Street
Calipatria, Calif. 92233
Telephone: 348-5150

ACCEPTED
10-28-69
10163-14

The Dune Company
AGRICULTURAL CHEMICALS

ARIZONA
Route 1, Box 33
Somerton, Arizona 85350
Telephone: 697-2479

P R O

SUGGESTED USE

READ ENTIRE LABEL. USE STRICTLY IN ACCORDANCE WITH CAUTIONS, WARNINGS AND DIRECTIONS, AND WITH APPLICABLE STATE AND FEDERAL REGULATIONS. TIMING AND DOSAGE MAY VARY AS A CONSEQUENCE OF LOCAL WEATHER OR CONDITIONS; WE RECOMMEND REFERENCE TO AGRICULTURAL EXPERIMENT STATIONS ON SPECIFIC USAGE

Apply to the undersides of leaves and food crops within 10 days after emergence of insects as specified. Remove insects from plants by washing, brushing or other means.

CUCUMBERS SQUASH MELONS: Aphids, Spider Mites, Pickleworm, Cucumber Beetles, Flea Beetles; **CANTALOUPE** — Leafhoppers — 30 to 35 lbs. per acre. Repeat as necessary. **SQUASH:** Squash Vine Borers — 30 lbs. per acre. **PUMPKINS:** Squash Vine Borers — 30 lbs. per acre. Apply 7 to 10 weeks before harvest. Do not apply to flowers. Do not apply to Cucumbers, cantaloupes, melons, or pumpkins. Malathion will injure plants if plants are dry.

Malathion Cryolite D

Active Ingredients:
Malathion*
Sodium Fluoroborate
Inert Ingredients:
*O,O dimethyl phosphorodithioic acid
Fluorine oxide

READ THE LABEL
KEEP OUT OF REACH OF CHILDREN

CAUTION

Avoid contact with eyes, nose, mouth, and skin. Do not drink or eat food that might be damaged. Do not use on food crops unless possible drift to food crops is avoided. Do not use on food crops unless possible drift to food crops is avoided. Do not use on food crops unless possible drift to food crops is avoided. Do not use on food crops unless possible drift to food crops is avoided.

NET WEIGHT

CALIFORNIA (Plant)
P. O. Box 406
340 East Main Street
Calipatria, Calif. 92233
Telephone: 348-5150

ACCEPTED

10-28-69

The Dune

AGRICULTURAL

10163-14