

Prokil Dibrom^(R) Cryolite 3-50 Dust

ACTIVE INGREDIENTS	% by Wt
Naled*	3
Sodium Fluoaluminate	50
INERT INGREDIENTS	47
	100
*1 2-dibromo-2, 2-dichloroethyl dimethyl phosphate	
U.S. Pat. No. 2,971,882	
Flourine expressed as elemental	27.1

USE WITHIN 60 DAYS OF MANUFACTURE ONLY, AS THIS PRODUCT DECOMPOSES

DATE OF MANUFACTURE 19...

(R) Dibrom is a Registered Trademark of Chevron Chemical Company

CAUTION

KEEP OUT OF REACH OF CHILDREN

CAUTION: Harmful if swallowed. May cause irritation of eyes, nose, and throat. Avoid breathing dust. Avoid contamination of skin and eyes. Wash thoroughly after using. Avoid contamination of feed and foodstuffs. Do not use on cherries as injury may result.

THIS PRODUCT IS HIGHLY TOXIC TO BEES exposed to treatment or residues on crops. Protective information may be obtained from your Cooperative Agricultural Extension Service.

This product will kill fish and wildlife, keep out of lakes, streams, or ponds. Birds feeding on treated areas may be killed. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment, or disposal of wastes. Apply this product only as specified on this label.

May corrode metal dust equipment, clean equipment after use.

STORE IN COOL DRY PLACE — OUT OF DIRECT SUNLIGHT.

BURN BAG IMMEDIATELY WHEN EMPTY. STAY OUT OF SMOKE.

NET WEIGHT 50 LBS.

U.S.D.A. Reg. No. 10163-18 — Calif Reg. No. 10163 50016 AA

READ ENTIRE LABEL USE STRICTLY IN ACCORDANCE WITH CAUTIONS, DIRECTIONS, AND WITH APPLICABLE STATE AND FEDERAL REGULATIONS. USES, TIMING AND DOSAGE MAY VARY AS A CONSEQUENCE OF WEATHER OR CONDITIONS, WE RECOMMEND REFERENCE TO STATE AGRICULTURAL EXTENSION SERVICE CONCERNING SPECIFIC USAGE.

Apply PROKIL DIBROM CRYOLITE 3-50 DUST at the rate of 25-50 lbs. Use the higher dosage when treating for aphids, lygus bugs, hornworms, and miners. This product is formulated for application by ground or air. For Airplane Application, use heavier dosages. Begin applications at first insects. Repeat as necessary.

LETTUCE Aphids, Climbing Cutworms, Alfalfa Looper, Corn Earworm — when worms first appear while still small and before they get down head. Do Not Apply after seedling stage.

BROCCOLI, CABBAGE Aphids, Climbing Cutworms, Cabbage Looper — when first worms appear while still small. Do Not Apply after edibles begin to form.

MELONS, WINTER SQUASH Aphids, Armyworms, Lygus Bugs, Cabbage Looper — Do Not Apply within 1 day of harvest. Remove excess residues by brushing or other effective means.

CUCUMBERS, SUMMER SQUASH Aphids, Armyworms, Lygus Bugs, Looper — May be applied up to harvest. Remove excess residues by brushing or other effective means.


TOMATOES Leaf Beetles, Leafminers, Tomato Fruitworms, Hornworms — when plants are first set out and repeat as necessary. Do Not apply day of harvest. If applied after fruit starts to set remove residues at harvest by washing, wiping or other effective means.

NOTICE ON CONDITIONS OF SALE

This product is sold on the basis of its use as described on the label. The user assumes all responsibility for the results obtained. The manufacturer is not responsible for the results obtained when the product is used in a manner not specified on the label. The user should read the label carefully and follow the directions. The manufacturer is not responsible for the results obtained when the product is used in a manner not specified on the label.

371-128-Proof

CALIFORNIA (Plant)
P. O. Box 406
340 East Main Street
Calipatria, Calif. 92233
Telephone: 348-5150



The Dune Company
AGRICULTURAL CHEMICALS



Route 1,
Somerton, Arizona
Telephone: 61

Prokil Dibrom^(R) Cryolite 3-50 Dust

ACTIVE INGREDIENTS	% by Wt
Naled*	3
Sodium Fluoaluminate	50
INERT INGREDIENTS	47
	100
* 1,2-dibromo-2,2-dichloroethyl dimethyl phosphate	
U.S. Pat. No. 2,971,882	
Flourine expressed as elemental	27.1

WITHIN 60 DAYS OF MANUFACTURE ONLY, AS THIS PRODUCT DECOMPOSES

DATE OF MANUFACTURE . . . 19 . . .

(R) Dibrom is a Registered Trademark of Chevron Chemical Company.

CAUTION

KEEP OUT OF REACH OF CHILDREN

WARNING: Harmful if swallowed. May cause irritation of eyes, nose, and throat. Avoid breathing dust. Avoid contamination of skin and eyes. Wash thoroughly after using. Avoid contamination of feed and foodstuffs. Do not use on cherries if injury may result.

PRODUCT IS HIGHLY TOXIC TO BEES exposed to treatment or residues on plants. Protective information may be obtained from your Cooperative Agricultural Extension Service.

This product will kill fish and wildlife, keep out of lakes, streams, or ponds. Do not feed on treated areas may be killed. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment, or disposal of wastes. Apply this product only as specified on this label.

Do not corrode metal dust equipment, clean equipment after use.

STORE IN COOL DRY PLACE — OUT OF DIRECT SUNLIGHT.

BURN BAG IMMEDIATELY WHEN EMPTY. STAY OUT OF SMOKE.

NET WEIGHT 50 LBS.

F.A. Reg. No. 10163-18 — Calif. Reg. No. 10163 50016 AA

READ ENTIRE LABEL. USE STRICTLY IN ACCORDANCE WITH CAUTIONS, WARNINGS AND DIRECTIONS, AND WITH APPLICABLE STATE AND FEDERAL REGULATIONS. USES, TIMING AND DOSAGE MAY VARY AS A CONSEQUENCE OF LOCAL WEATHER OR CONDITIONS, WE RECOMMEND REFERENCE TO STATE AGRICULTURAL EXTENSION SERVICE CONCERNING SPECIFIC USAGE.

Apply PROKIL DIBROM CRYOLITE 3-50 DUST at the rate of 25-50 lbs. per acre. Use the higher dosage when treating for aphids, lygus bugs, hornworms, and leaf miners. This product is formulated for application by ground or air equipment. For Airplane Application, use heavier dosages. Begin applications at first sign of insects. Repeat as necessary.

LETTUCE Aphids, Climbing Cutworms, Alfalfa Looper, Corn Earworm. Apply when worms first appear while still small and before they get down into the head. Do Not Apply after seedling stage.

BROCCOLI, CABBAGE Aphids, Climbing Cutworms, Cabbage Looper — Apply when first worms appear while still small. Do Not Apply after edible parts begin to form.

MELONS, WINTER SQUASH. Aphids, Armyworms, Lygus Bugs, Cabbage Looper. Do Not Apply within 1 day of harvest. Remove excess residues by washing, brushing or other effective means.


CUCUMBERS, SUMMER SQUASH. Aphids, Armyworms, Lygus Bugs, Cabbage Loopers — May be applied up to harvest. Remove excess residues by washing, brushing or other effective means.

TOMATOES Flea Beetles, Leafminers, Tomato Fruitworms, Hornworms — Apply when plants are first set out and repeat as necessary. Do Not apply within 1 day of harvest. If applied after fruit starts to set remove residues at harvest by washing, wiping or other effective means.

NOTICE ON CONDITIONS OF SALE

3-71-128-- Proof

CALIFORNIA (Plant)
P.O. Box 406
100 East Main Street
Imperial, Calif. 92233
Telephone: 348-5150



The Dune Company
AGRICULTURAL CHEMICALS



ARIZONA
Route 1, Box 33
Somerton, Arizona 85350
Telephone: 627-2474

Prokil Dibrom^(R)

ACTIVE INGREDIENTS	% by Wt.
Naled*	3
Sodium Fluoaluminate	50
INERT INGREDIENTS	47
	100

*1 2-dibromo-2, 2-dichloroethyl dimethyl phosphate
U.S. Pat. No. 2,971,882

Flourine expressed as elemental 27.1

USE WITHIN 60 DAYS OF MANUFACTURE ONLY, AS THIS PRODUCT DECOMPOSES

DATE OF MANUFACTURE 19__

(R) Dibrom is a Registered Trademark of Chevron Chemical Company.

CAUTION

KEEP OUT OF REACH OF CHILDREN

CAUTION: Harmful if swallowed. May cause irritation of eyes, nose, and throat. Avoid breathing dust. Avoid contamination of skin and eyes. Wash thoroughly after using. Avoid contamination of feed and foodstuffs. Do not use on cherries as injury may result.

THIS PRODUCT IS HIGHLY TOXIC TO BEES exposed to treatment or residues on crops. Protective information may be obtained from your Cooperative Agricultural Extension Service.

This product will kill fish and wildlife, keep out of lakes, streams, or ponds. Birds feeding on treated areas may be killed. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment, or disposal of wastes. Apply this product only as specified on this label.

May corrode metal dust equipment, clean equipment after use.


STORE IN COOL DRY PLACE -- OUT OF DIRECT SUNLIGHT.

BURN BAG IMMEDIATELY WHEN EMPTY. STAY OUT OF SMOKE.

NET WEIGHT 50 LBS.

U.S.D.A. Reg. No. 10163-18 - Calif. Reg. No. 10163 50016 AA

CALIFORNIA (Plant)
P. O. Box 406
340 East Main Street
Calipatria, Calif. 92233
Telephone: 348-5150



The Dunc

AGRICULTURE