



4 / 16 / 2014
UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
WASHINGTON, DC 20460

1/13

OFFICE OF CHEMICAL SAFETY
AND POLLUTION PREVENTION

April 22, 2015

Ms. Emily Foley
Registration Specialist
Gowan Company
P.O. Box 5569
Yuma, AZ 85366-5569

Subject: Clarification of the Subject Line of the Letter for Label Stamped April 16, 2014
Product Name: Malathion 8
EPA Registration Number: 10163-21
Associated Decision Numbers: 483078; 485256

Dear Ms. Foley:

On April 16, 2014, the Agency sent Gowan Company a letter that accepted the product label for 10163-21 which included only minor changes from the previously accepted label dated July 14, 2010. However, the 4/16/14 Agency letter subject line stated "Submission of amended labeling adding me-too uses for aerial applications and crops (asparagus, carrots, chestnuts, melon, papaya, pineapple pumpkin, turnips and watermelon)". Although these me-too uses were what was originally requested, the Agency would like to clarify that none of the above mentioned me-too uses were granted.

The request to add aerial application and additional crops to the label were reviewed, and found not acceptable at that time due to data compensation issues. Therefore, during the review process, a revised label was submitted without the aerial and additional crop uses that was deemed acceptable. This was the label stamped on April 16, 2014.

If you have any questions, please feel free to contact me at (703) 308-8045.

Regards,

Venus Eagle, Product Manager 1
Invertebrate & Vertebrate Branch 3
Registration Division (7505P)

2/13

UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
WASHINGTON, D.C. 20460



OFFICE OF
CHEMICAL SAFETY AND
POLLUTION PREVENTION

APR 16 2014

Tracey Mixon
Registration Specialist
Gowan Company
P.O. Box 5569
Yuma, AZ 85366-5569

Dear Ms. Mixon:

Subject: Submission of amended labeling adding me-too uses for aerial applications and crops (asparagus, carrots, chestnuts, melon, papaya, pineapple, pumpkin, turnips & watermelon)
EPA Reg. No. 10163-21
Submissions dated September 9, 2013 and November 20, 2013

The proposed labeling referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide, and Rodenticide Act (FIFRA), is acceptable provided that you submit two copies of your final printed label before you release the product for shipment. Products shipped after 18 months from the date of this amendment or the next printing of the label which ever occurs first, must bear the new revised label. Your release for shipment of the product bearing the amended label constitutes acceptance of these conditions. If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA sec. 6(e).

Additional label corrections may be needed pending submittal and review of your responses to the Malathion RED.

A stamped copy is enclosed for your records. If you have any questions, please contact Marianne Lewis at (703) 308-8043 or lewis.marianne@epa.gov.

Regards,

Venus Eagle
Insecticide-Rodenticide Branch
Registration Division (7505P)

Enclosure

GROUP 1B INSECTICIDE

GOWAN MALATHION 8

AGRICULTURAL INSECTICIDE

ACTIVE INGREDIENT:		% By Wt.
Malathion (O,O-dimethyl phosphorodithioate of diethyl mercaptosuccinate):	79.5%
INERT INGREDIENTS	20.5%
		TOTAL 100.0%

Contains Petroleum Distillates
 Contains 8 lbs. Malathion per gallon
KEEP OUT OF REACH OF CHILDREN
CAUTION

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle.
 (If you do not understand the label, find someone to explain it to you in detail.)

FIRST AID	
Organophosphate Insecticide	
If swallowed	<ul style="list-style-type: none"> • Immediately call a poison control center or doctor. • Do not induce vomiting unless told to by a poison control center or doctor. • Do not give any liquid to the person. • Do not give anything by mouth to an unconscious person.
If in eyes	<ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing. • Call a poison control center or doctor for treatment advice.
If on skin or clothing	<ul style="list-style-type: none"> • Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15-20 minutes. • Call a poison control center or doctor for treatment advice.
If inhaled	<ul style="list-style-type: none"> • Move person to fresh air. • If person is not breathing, call 911 or ambulance, then give artificial respiration, preferably mouth-to-mouth if possible. • Call a poison control center or doctor for further treatment advice.
HOT LINE NUMBER	
Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-888-478-0798 for emergency medical treatment information.	
NOTE TO PHYSICIAN	
Malathion upon use may cause cholinesterase inhibition. Atropine is antidotal. May pose an aspiration pneumonia hazard. Contains petroleum distillates.	

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION

Harmful if swallowed. Avoid breathing of spray mist. Avoid contact with skin.

PERSONAL PROTECTIVE EQUIPMENT (PPE)

Some materials that are chemical-resistant to this product are barrier laminate, butyl rubber, nitrile, or viton. If you want more options, follow the instructions for category F on an EPA chemical resistance category selection chart.

Mixers, Loaders, Applicators, Flaggers, and other Handlers must wear:

- Long-sleeved shirt and long pants
- Chemical-resistant gloves
- Shoes plus socks

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

NET CONTENTS _____ GALLONS ACCEPTED

4/16/14

Under the Federal Insecticide, Fungicide,
 and Rodenticide Act, as amended, for the
 pesticide registered under:

Produced For:
 Gowan Company
 P. O. Box 5569
 Tulsa, AZ 85366-5569
 800-883-1844

EPA Reg. No. 10163-21
 EPA Est. No.



EPA. Reg. No: 10163-21

USER SAFETY RECOMMENDATIONS

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

ENGINEERING CONTROLS

Pilots must use an enclosed cockpit in a manner that is consistent with the WPS for Agricultural Pesticides [40 CFR 170.240(d)(6)]. Pilots must wear the PPE required on this labeling for applicators.

ENVIRONMENTAL HAZARDS

This pesticide is toxic to aquatic organisms, including fish and invertebrates. This product may contaminate water through drift of spray in wind. This product has a high potential for runoff after application. Use care when applying in or to an area which is adjacent to any body of water, and do not apply when weather conditions favor drift from target area. Poorly draining soils and soils with shallow water tables are more prone to produce runoff that contains this product.

A level, well maintained vegetative buffer strip between areas to which this product is applied and surface water features such as ponds, streams, and springs will reduce the potential for contamination of water from rainfall-runoff. Runoff of this product will be reduced by avoiding applications when rainfall is forecasted to occur within 48 hours.

Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwater or rinsate.

This pesticide is highly toxic to bees exposed to direct treatment on blooming crops or weeds. Do not apply this product or allow it to drift to blooming crops or weeds **when bees are foraging** the treatment area.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, and nurseries, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), and restricted-entry intervals. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted-entry interval (REI). The REI for each crop is listed in the directions for use associated with each crop.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water is:

- Coveralls
- Chemical-resistant gloves, made out of any waterproof material
- Shoes plus socks

NON-AGRICULTURAL USE REQUIREMENTS

The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard (WPS) for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, or nurseries.

Do not enter or allow others to enter until sprays have dried.

PRECAUTIONS AND RESTRICTIONS

In order that pesticide residues on food and forage crops will not exceed tolerances established by the Federal Food and Drug Administration, use only the specified rates and intervals, and do not apply closer to harvest than prescribed.

Unless otherwise specified, apply at the first sign of infestation and repeat as needed observing the use limitations listed for each specified crop in the application tables. Consult your State Agriculture Experiment Station or the State Agricultural Extension Service for additional information as the timing of applications needed will vary with local conditions.

Applications may be made by aircraft or by ground equipment according to the DIRECTIONS FOR DILUTION below. The amount of water needed to treat an acre varies, therefore the following directions are given to cover a broad range of applications.

5/13

Buffer Zones for Aerial Application:

When making a Non-ULV application with aerial application equipment, a minimum buffer zone of 25 feet must be maintained along any water body.

Do not use in greenhouses.

PHYTOTOXICITY ADVISORY STATEMENT

As is common with most emulsifiable concentrate formulations adverse effects, such as spotting or discoloration of the fruit or foliage can occur. Some conditions known to contribute to phytotoxicity include, but are not limited to : high temperatures, poor spray drying conditions, excessive spray runoff, certain spray mixtures, stage of crop development or tank mixes with other pesticides.

SPRAY DRIFT REQUIREMENTS

Observe the following requirements when spraying in the vicinity of aquatic areas such as, but not limited to lakes; reservoirs; rivers; permanent streams; marshes or natural ponds; estuaries and commercial fish ponds.

Droplet Size: Use the largest droplet size consistent with acceptable efficiency. Formation of very small droplets may be minimized by appropriate nozzle selection, by orienting nozzles away from the air stream as much as possible, and by avoiding excessive spray boom pressure.

For groundboom and aerial applications, use only medium or coarser spray nozzles according to ASAE (S572) definition for standard nozzles, or a volume mean diameter (VMD) of 300 microns or greater for spinning atomizer nozzles. In conditions of low humidity and high temperatures, applicators should use a coarser droplet size.

Wind Direction and Speed: Make aerial or ground applications when the wind velocity favors on target product deposition (approximately 3 to 10 mph). Do not apply when wind velocity exceeds 15 mph. Avoid applications when wind gusts approach 15 mph. For all non-aerial applications, wind speed must be measured adjacent to the application site on the upwind side, immediately prior to application.

Temperature Inversion: Do not make aerial or ground applications into areas of temperature inversions. Inversions are characterized by stable air and increasing temperatures with increasing distance above the ground. Mist or fog may indicate the presence of an inversion in humid areas. Where permissible by local regulations, the applicator may detect the presence of an inversion by producing smoke and observing a smoke layer near the ground surface. In conditions of low humidity and high temperatures, applicators should use a coarser droplet size.

Additional Requirements for Ground Applications: Spray should be released at the lowest height consistent with pest control and flight safety. Applications more than 10 feet above the crop canopy should be avoided. For groundboom applications, apply with nozzle height no more than 4 feet above the ground or crop canopy. For airblast applications, turn off outward pointing nozzles at row ends and when spraying the outer two rows. To minimize spray loss over the top in orchard applications, spray must be directed into the canopy.

Additional Requirements for Aerial Applications: For aerial applications, the spray boom should be mounted on the aircraft as to minimize drift caused by wingtip or rotor vortices. The minimum practical boom length should be used and must not exceed 75% of wing span or 90% rotor diameter. Aerial applicators must consider flight speed and nozzle orientation in determining droplet size. When applications are made with a cross-wind, the swath will be displaced downwind. The applicator must compensate for this displacement at the downwind edge of the application area by adjusting the path of the aircraft upwind.

MIXING DIRECTIONS

Pour specified amount of product into spray tank nearly filled with water. Add balance of water to fill tank. Keep agitator running during filling and spraying operations. If mixture does not mix readily, but tends to separate as an oily layer, do not use as injury to plants may result. Do not combine with wettable powders unless previous use of the mixture has proven physically compatible and safe to plants. Always thoroughly emulsify this product with at least half of total water before adding wettable powders.

PREHARVEST INTERVAL

Minimum days between last application and harvest are given in () after each crop name.

TREES AND VINES

Rates are given in terms of pints of Malathion 8 Flowable per acre. Under heavy pest pressure, use higher rates.

Dilute Application: Use specified rate in 100 to 800 gallons of water per acre.

Aerial Application: Use a minimum of 10 gallons of water per acre.

CROP	REI (HRS)	RATE (PTS/ACRE)	PESTS	COMMENTS
APRICOTS (7)	12	1.5	Aphid, Codling moth, European Lecanium scale, Orange tortrix, Soft brown scale, Terrapin scale	The maximum application rate is 1.5 pints of product per acre; the maximum number of applications per year is 2; and the minimum retreatment interval is 7 days.
AVOCADOS (7)	48	4.7	Green house thrips, Latania scale, Omnivorous looper, Soft brown scale, Orange tortrix	The maximum application rate is 4.7 pints of product per acre; the maximum number of applications per year is 2; and the minimum retreatment interval is 30 days.

CROP	REI (HRS)	RATE (PTS/ACRE)	PESTS	COMMENTS
BLACKBERRIES (1), BOYSENBERRIES (1), DEWBERRIES (1), LOGANBERRIES (1), RASPBERRIES (1)	12	2	Japanese beetle, Leafhoppers, Mites, Thrips	The maximum application rate is 2.0 pints of product per acre; the maximum number of applications per year is 3; and the minimum retreatment interval is 7 days.
		2	Aphid, Rose scale	
BLUEBERRIES (1)	12	1.25	Aphids, Blueberry maggot, Blueberry tip borer, Cherry fruitworm, Cranberry fruitworm, Japanese beetle, Plum curculio, Leafrollers, Sharp-nosed leafhopper, White Tussock moth	The maximum application rate is 1.25 pints of product per acre; the maximum number of applications per year is 3; and the minimum retreatment interval is 5 days.
CHERRIES (3)	12	1.75	Black cherry aphid, Bud moth, Cherry fruit fly, Fruittree leafroller, Lesser peach twig borer, Forbes and San Jose scale	For Lesser peach twig borer, apply to trunk and scaffold limbs at 21 day intervals beginning with emergence (Do not exceed 4 applications per year). May cause injury on certain varieties of sweet cherries. The maximum application rate is 1.75 pints of product per acre; the maximum number of applications per year is 4; and the minimum retreatment interval is 3 days.
CITRUS [GRAPEFRUIT, LEMONS, LIMES, ORANGES, TANGELOS, TANGERINES [Mandarin or Mandarin Oranges, Tangors, and other hybrids of tangerines with other citrus] (7)	72	CA: 7.5 All Other States: 4.5	Aphids, Black scale (single and off-brooded), California red scale, Citricola scale, Orange worm, Purple scale, Soft scale, Thrips, Yellow scale	Do not apply when trees are in bloom. FOR CALIFORNIA: The maximum application rate is 7.5 pints of product per acre; the maximum number of applications per year is 1. ALL OTHER STATES: The maximum application rate is 4.5 pints of product per acre; the maximum number of applications per year is 1.
		CA: 1 - 7.5 All Other States: 1 - 4.5	Mediterranean fruit fly	
KUMQUATS (7)	48	4.5	Aphids, Black scale (single and off-brooded), California red scale, Citricola scale, Orange worm, Purple scale, Soft scale, Thrips, Yellow scale	Do not apply when trees are in bloom. The maximum application rate is 4.5 pints product per acre; the maximum number of applications per year is 1.
CURRANTS (1)	12	1.25	Japanese beetle, Mites	The maximum application rate is 1.25 pints product per acre; the maximum number of applications per year is 3; the minimum retreatment interval is 7 days.
GOOSEBERRIES (3)	12	2	Currant aphid, Imported currantworm	The maximum application rate is 2.0 pints product per acre; the maximum number of applications per year is 3; the minimum retreatment interval is 7 days.
FIGS (5)	12	1.5	Dried fruit beetles, Vinegar flies	Apply with 1 - 2 gallons sulfured molasses per acre. The maximum application rate is 1.5 pints product per acre; the maximum number of applications per year is 2; the minimum retreatment interval is 5 days.
GRAPES (3)	72 girdling and tying 24 other activities	1.88	Drosophila, European fruit lecanium, Grape leafhopper, Japanese beetle, Leafhopper, Mealybug, Spider mites, Terrapin scale	Injury may occur to grape berries when applications are made after bloom. The maximum application rate is 1.88 pints product per acre; the maximum number of applications per year is 2; and the minimum retreatment interval is 14 days.
GUAVA (2)	12	.75 - 1.25	Fruit flies	Apply with 1 pound partially hydrolyzed yeast protein or enzymatic yeast hydrolyzate. The maximum application rate is 1.25 pints product per acre; the maximum number of applications per year is 13; and the minimum retreatment interval is 3 days.
MANGO (1)	12	0.9375	Fruit flies	The maximum application rate is 0.9375 pints product per acre; the maximum number of applications per year is 10; and the minimum retreatment interval is 7 days.
PASSION FRUIT (3)	12	1	Fruit flies	The maximum application rate is 1.0 pints product per acre; the maximum number of applications per year is 8; and the minimum retreatment interval is 7 days.

CROP	REI (HRS)	RATE (PTS/ACRE)	PESTS	COMMENTS
MACADAMIA NUTS (1)	12	0.94	Green Stink bug	The maximum application rate is 0.94 pints product per acre; the maximum number of applications per year is 6; and the minimum retreatment interval is 7 days.
NECTARINES (7)	24	3	Black cherry aphid, Black peach aphid, Green peach aphid, Japanese beetle, Rusty plum aphid	May be mixed with spray oil for dormant and delayed dormant applications. Follow spray oil manufacturer's directions. The maximum application rate is 3.0 pints product per acre; the maximum number of applications per year is 3; and the minimum retreatment interval is 7 days.
PEACHES (7)	24	1.25	Cottony peach scale, Lesser peach tree borer, Plum curculio, Oriental fruit moth, San Jose scale, Terrapin scale	The maximum application rate is 1.25 pints product per acre; the maximum number of applications per year is 3; and the minimum retreatment interval is 11 days.
PECANS (7)	24	2.5	Aphid, Mites, Pecan bud moth, Pecan leaf casebearer, Pecan nut casebearer, Pecan phylloxera	The maximum application rate is 2.5 pints product per acre; the maximum number of applications per year is 3; and the minimum retreatment interval is 7 days.
WALNUTS (7)	24	1.5 - 2.5	Aphid, Mites, Walnut husk fly	The maximum application rate is 2.5 pints product per acre; the maximum number of applications per year is 3; and the minimum retreatment interval is 7 days.

FIELD AND ROW CROPS

Rates are given in terms of pints of Malathion 8 Flowable per acre. Under heavy pest pressure, use higher rates.

Dilute Application: Use specified rate in 20 to 60 gallons of water per acre.

Aerial Application: Use a minimum of 2 gallons of water per acre.

CROP	REI (HRS)	RATE (PTS/ACRE)	PESTS	COMMENTS
ALFALFA, BIRDSFOOT TREFOIL, CLOVER, LESPEDEZA, VETCH (0)	12	1 - 1.25	Alfalfa weevil larvae, Aphids, Armyworms, Clover leaf weevil, Grasshoppers, Lygus bugs, Pea aphid, Potato leafhoppers, Spider mites, Spittlebug, Vetch bruchid	Use higher rate for Armyworm control. Apply to alfalfa in bloom only in the evening or early morning when bees are not working in the fields or are not hanging on the outside of hives. The maximum application rate is 1.25 pints product per acre; the maximum number of applications is 2 per cutting; and the minimum retreatment interval is 14 days.
LEAFY VEGETABLES (EXCEPT BRASSICA VEGETABLES) CROP GROUPING: AMARANTH (LEAFY AMARANTH, CHINESE SPINACH, TAMPALA) (7), ARRUGULA (ROQUETTE) (7), CELTUCE (7), CHERVIL (7), CHRYSANTHEMUM-Edible-leafed, Garland (7), CORN SALAD (7), DOCK (SORREL) (7), FLORENCE FENNEL (7), ORACH (7), PURSLANE-Garden and Winter (7)	24	1 - 1.25	Aphids	The maximum application rate is 1.25 pints product per acre; the maximum number of applications per year is 2; and the minimum retreatment interval is 7 days.
DANDELIONS (7)	24	1.25	Aphids	The maximum application rate is 1.25 pints product per acre; the maximum number of applications per year is 2; and the minimum retreatment interval is 7 days.
PARSLEY (7)	24	1.5	Aphids	The maximum application rate is 1.5 pints product per acre; the maximum number of applications per year is 2; and the minimum retreatment interval is 7 days.
SWISS CHARD (14)	12	1.0	Aphids	The maximum application rate is 1.0 pints product per acre; the maximum number of applications per year is 2; and the minimum retreatment interval is 7 days.
CELERY (7)	24	1.0 - 1.5	Aphids, spider mite	The maximum application rate is 1.5 pints product per acre; the maximum number of applications per year is 2; and the minimum retreatment interval is 7 days.

CROP	REI (HRS)	RATE (PTS/ACRE)	PESTS	COMMENTS
LETTUCE, FIELD HEAD (14)	24	1.88	Aphids, Alfalfa loopers, Leafhoppers, Mites	The maximum application rate is 1.88 pints product per acre; the maximum number of applications per year is 2; and the minimum retreatment interval is 6 days.
LETTUCE, FIELD LEAF (14)	24	1.88	Aphids, Alfalfa loopers, Leafhoppers, Mites	The maximum application rate is 2.0 pints product per acre; the maximum number of applications per year is 2; and the minimum retreatment interval is 5 days.
ENDIVE, FIELD (7)	24	1.25	Aphids, Alfalfa loopers, Leafhoppers, Mites	The maximum application rate is 1.25 pints product per acre; the maximum number of applications per year is 2; and the minimum retreatment interval is 7 days.
SPINACH (7)	12	1.0	Aphids	The maximum application rate is 1.0-5 pints product per acre; the maximum number of applications per year is 2; and the minimum retreatment interval is 7 days.
BEETS, Table (7)	12	1.25	Aphids, Beet armyworm, Blister beetles, Flea beetles	Do not use on Sugar Beets. The maximum application rate is 1.25 pints product per acre; the maximum number of applications per year is 3; and the minimum retreatment interval is 7 days.
COLE CROPS (Brassica (cole) Leafy Vegetable crop group: BROCCOLI (2), BROCCOLI RAAB (RAPINI) (2), BRUSSELS SPROUTS (2), CAULIFLOWER (2), CAVALO BROCCOLO (2), CHINESE BROCCOLI (2), CHINESE MUSTARD CABBAGE (7), MIZUNA (7), MUSTARD SPINACH (7), RAPE GREENS (7)	48	1.25	Aphids, Cabbage loopers, Flea beetles, Imported cabbage worms	The maximum application rate is 1.25 pints product per acre; the maximum number of applications per year is 2; and the minimum retreatment interval is 7 days.
CABBAGE (7)	48	1.25	Aphids, Cabbage loopers, Flea beetles, Imported cabbage worms	The maximum application rate is 1.25 pints product per acre; the maximum number of applications per year is 6; and the minimum retreatment interval is 7 days.
CHINESE CABBAGE (BOK CHOY, NAPA) (7)	48	1.25	Aphids, Cabbage loopers, Flea beetles, Imported cabbage worms	The maximum application rate is 1.25 pints product per acre; the maximum number of applications per year is 2; and the minimum retreatment interval is 7 days.
COLLARDS (7)	12	1	Aphids, Cabbage loopers, Flea beetles, Imported cabbage worms	The maximum application rate is 1.0 pints product per acre; the maximum number of applications per year is 3; and the minimum retreatment interval is 7 days.
KALE (7), MUSTARD GREENS (7),	12	1	Aphids, Cabbage loopers, Flea beetles, Imported cabbage worms	The maximum application rate is 1.0 pints product per acre; the maximum number of applications per year is 3; and the minimum retreatment interval is 5 days.
KOHLRABI (7)	24	1.25	Aphids, Cabbage loopers, Flea beetles, Imported cabbage worms	The maximum application rate is 1.25 pints product per acre; the maximum number of applications per year is 2; and the minimum retreatment interval is 7 days.
CORN-Field (7)	72 hours for detasseling 12 hours for all other activities	0.61	Aphids, Corn rootworm adults, Sap beetles, Thrips, Young grasshoppers	CAUTION: Injury may occur in whorl and silk stages. The maximum application rate is 0.61 pints product per acre; the maximum number of applications per year is 2; and the minimum retreatment interval is 7 days.
COTTON (7)	48	2.5	Aphids, Brown cotton leafworm, Cotton leaf perforator, Leafhoppers, Spider mites, Whitefly, Boll weevils, Cotton fleahoppers, Fall armyworms, Grasshoppers, Garden webworms and Lygus	Do not graze or feed forage to livestock. The maximum application rate is 2.5 pints product per acre; the maximum number of applications per year is 3; and the minimum retreatment interval is 7 days.
CUCUMBERS (1)	24	1.75	Aphids, Cucumber beetles, Cutworms, Darkling ground beetles, Leafhoppers, Pickleworm, Spider mites, Squash vine borer, Thrips	Do not apply unless plants are dry. For vine borer apply to stems and vines at base of plant. The maximum application rate is 1.75 pints product per acre; the maximum number of applications per year is 2; and the minimum retreatment interval is 7 days.

9/13

CROP	REI (HRS)	RATE (PTS/ACRE)	PESTS	COMMENTS
SQUASH, Summer (1)	24	1.75	Aphids, Cucumber beetles, Cutworms, Darkling ground beetles, Leafhoppers, Pickleworm, Spider mites, Squash vine borer, Thrips	The maximum application rate is 1.75 pints product per acre; the maximum number of applications per year is 3; and the minimum retreatment interval is 7 days.
SQUASH, Winter (1)	12	1	Aphids, Cucumber beetles, Cutworms, Darkling ground beetles, Leafhoppers, Pickleworm, Spider mites, Squash vine borer, Thrips	The maximum application rate is 1.0 pints product per acre; the maximum number of applications per year is 3; and the minimum retreatment interval is 7 days.
EGGPLANT (3)	12	1.56	Aphids, Spider mites, Lace bugs	The maximum application rate is 1.56 pints product per acre; the maximum number of applications per year is 4; and the minimum retreatment interval is 7 days.
FLAX (52)	12	0.5	Grasshoppers	The maximum application rate is 0.5 pints product per acre; the maximum number of applications per year is 3; and the minimum retreatment interval is 7 days.
GARLIC (3),	24	1 -1.56	Aphids, Thrips	The maximum application rate is 1.56 pints product per acre; the maximum number of applications per year is 3; and the minimum retreatment interval is 7 days.
LEEKs (3), SHALLOTS (3)	24	1 - 1.56	Aphids, Thrips	The maximum application rate is 1.56 pints product per acre; the maximum number of applications per year is 2; and the minimum retreatment interval is 7 days.
GRASSES (Forage, Hay) (0)	12	1 - 1.25	Aphids, Grasshoppers, Leafhoppers	The maximum application rate is 1.25 pints product per acre; the maximum number of applications per year is 1.
HOPS (10)	12	0.63	Aphids	The maximum application rate is 0.63 pints product per acre; the maximum number of applications per year is 3; and the minimum retreatment interval is 7 days.
HORSERADISH (7), PARSNIPS (7), SALSIFY (7)	24	1.25	Aphids, Diamondback moths, Flea beetles, Leafhoppers	The maximum application rate is 1.25 pints product per acre; the maximum number of applications per year is 3; and the minimum retreatment interval is 7 days.
RADISHES (7)	12	1	Aphids, Diamondback moths, Flea beetles, Leafhoppers	The maximum application rate is 1.0 pints product per acre; the maximum number of applications per year is 3; and the minimum retreatment interval is 7 days.
MUSHROOMS (1)	12	1.7	Phorid flies, Sciarid flies	Apply in 130 gallons of water per acre, or 1 tablespoon per 3 gallons of water per 1000 square foot bed. Make thorough application as soon as possible after picking. The maximum application rate is 1.7 pints product per acre; the maximum number of applications per year is 4; and the minimum retreatment interval is 3 days.
OKRA (1)	12	1.2	Aphids, Japanese beetles	The maximum application rate is 1.2 pints product per acre; the maximum number of applications per year is 5; and the minimum retreatment interval is 7 days.
ONIONS- Bulb and Green (3)	12	1 -1.56	Thrips	The maximum application rate is 1.56 pints product per acre; the maximum number of applications per year is 2; and the minimum retreatment interval is 7 days.
		1.5	Onion maggots	
PEAS, DRIED (3)	12	1	Aphids, Pea weevils	Do not graze or feed forage to livestock. Dried peas can be treated by ground and foliar applications only. The maximum application rate is 1.0 pints product per acre; the maximum number of applications per year is 2; and the minimum retreatment interval is 7 days.
PEAS, GREEN (3)	12	1	Aphids, Pea weevils	Do not graze or feed forage to livestock. Green peas can be treated by ground, foliar and aerial applications. The maximum application rate is 1.0 pints product per acre; the maximum number of applications per year is 2; and the minimum retreatment interval is 7 days.

10/13

CROP	REI (HRS)	RATE (PTS/ACRE)	PESTS	COMMENTS
PEPPERMINT (7), SPEARMINT (7)	12	0.94	Adult flea beetles, Leafhoppers	The maximum application rate is 0.94 pints product per acre; the maximum number of applications per year is 3; and the minimum retreatment interval is 7 days.
PEPPERS (Field) (3)	12	1.5	Aphids, Pepper maggots	The maximum application rate is 1.5 pints product per acre; the maximum number of applications per year is 2; and the minimum retreatment interval is 5 days.
POTATOES (0)	12	1	False chinch bugs, Leafhoppers, Mealybugs	The maximum application rate is 1.5 pints product per acre; the maximum number of applications per year is 2; and the minimum retreatment interval is 7 days.
		1.5	Aphids, Blister beetles	
RICE-Domestic, Grain or Wild (7)	12	1.25	Rice leaf miners, Rice stink bugs	Do not apply Propanil within 15 days of Malathion treatment. Broadcast use only over intermittently flooded areas. Application may not be made around bodies of water where fish or shellfish are grown and/or harvested commercially. The maximum application rate is 1.25 pints product per acre; the maximum number of applications per year is 2; and the minimum retreatment interval is 7 days.
RUTABAGAS (7)	12	1	Aphids	The maximum application rate is 1.0 pints product per acre; the maximum number of applications per year is 3; and the minimum retreatment interval is 7 days.
SMALL GRAINS (BARLEY) (7)	12	1 - 1.25	Armyworms, English grain aphids, Grasshoppers, Greenbugs	The maximum application rate is 1.25 pints product per acre; the maximum number of applications per year is 3; and the minimum retreatment interval is 7 days.
SMALL GRAINS (OATS, RYE, WHEAT[spring and summer]) (7)	12	1	Armyworms, English grain aphids, Grasshoppers, Greenbugs	The maximum application rate is 1.0 pints product per acre; the maximum number of applications per year is 2; and the minimum retreatment interval is 7 days.
SORGHUM-Grain (7)	12	1.0	Greenbugs	Do not graze or feed forage to livestock. The maximum application rate is 1.0 pints product per acre; the maximum number of applications per year is 2; and the minimum retreatment interval is 7 days.
STRAWBERRIES (3)	12	1.5 - 2	Aphids, Field crickets, Lygus bugs, Potato leafhoppers, Spider mites, Spittlebugs, Strawberry leafrollers, Strawberry root weevils, Thrips, Whiteflies	The maximum application rate is 2.0 pints product per acre; the maximum number of applications per year is 4; and the minimum retreatment interval is 7 days.
SWEET CORN (Field) (5)	72 detasseling 12 other activities	1	Japanese beetles	CAUTION: Injury may occur in whorl and silk stages. The maximum application rate is 1.0 pints product per acre; the maximum number of applications per year is 5; and the minimum retreatment interval is 5 days.
SWEET POTATOES (3)	12	1 - 1.5	Leafhoppers	The maximum application rate is 1.5 pints product per acre; the maximum number of applications per year is 2; and the minimum retreatment interval is 7 days.
		1.5	Morning Glory leafminers	
TOMATOES (Field) (1)	12	1.5	Aphids, Spider mites, Drosophila flies	Apply a full coverage application to fruit and foliage. The maximum application rate is 1.5 pints product per acre; the maximum number of applications per year is 4; and the minimum retreatment interval is 5 days.
WATERCRESS (7)	12	1	Aphids	The maximum application rate is 1.0 pints product per acre; the maximum number of applications per year is 5; and the minimum retreatment interval is 3 days.

11/13

OUTDOOR ORNAMENTALS

Note: Before treating a large number of ornamental plants with Gowen Malathion 8 alone or as a tank mixture with any other material, make a test application on a few plants and observe for 7-10 days prior to treating large areas to reduce the possibility of plant injury.

CROP	REI (HRS)	RATE	PESTS	COMMENTS
FLOWERS, SHADE TREES and SHRUBS	12	1 pint in 100 gals of water as a dilute spray	Aphids, Euonymus scales, European pine shoot moths, Four-lined leaf bugs, Japanese beetle adults, Lace scales, Mealybugs, Millipedes, Oyster shell scales, Potato leafhoppers, Rose leafhoppers, Scurfy scales, Spider mites, Springtails, Sowbugs, Tarnished plant bugs, Thrips, Whiteflies	<p>CAUTION: Avoid use on certain ferns including Boston, Maidenhair and Pteris, as well as some species of Crassula and Canaetri Juniper.</p> <p>For Oyster shell, Fletch, Juniper, Oak kermes and Pine needle scales apply when scale crawlers have settled on foliage. The maximum number of applications per year is 2; and the minimum retreatment interval is 10 days.</p>
		1.25 pints in 100 gals of water as a dilute spray	Azalea scales, Bagworms, Birch leafminers, Boxwood leafminers, Fletch scales, Florida-red scales, Juniper scales, Magnolia scales, Oak kermes, Pine leaf scales, Tent caterpillars	
		1.6 pints in 100 gals of water	Black scale crawlers, Monterey pine scales	
		2 pints in 100 gals of water	Pine needle scales, Wax scales	

12/13

SLASH PINE, PINE SEED ORCHARDS, and CHRISTMAS TREE PLANTATIONS

CROP	REI (HRS)	PESTS	RATE	COMMENTS
SLASH PINE, and PINE SEED ORCHARDS	12	Slash pine flower thrips, European pine sawfly	For ground application, mix 0.4 gallons of Malathion 8 in 100 gallons of water.	Apply 3/4 gallon of the mixture per tree on the smallest flowering trees. Mist blowers or airblast sprays may be used. The maximum application rate is 3.2 pints product per acre; the maximum number of applications per year/growing season is 2; and the minimum retreatment interval is 7 days.
			For air application, mix 0.4 gallons of Malathion 8 in at least 10 gallons of water.	Apply a minimum of 10 gallons of mixture per acre. Make two applications, the first when female flowers are in twig bud stage, the second one week prior to maximum flower receptivity to pollen. The maximum application rate is 3.2 pints product per acre; the maximum number of applications per year/growing season is 2; and the minimum retreatment interval is 7 days.
CHRISTMAS TREE PLANTATIONS	12	Slash pine flower thrips, European pine sawfly	For ground application, mix 0.4 gallons of Malathion 8 in 100 gallons of water.	Apply 3/4 gallon of the mixture per tree on the smallest flowering trees. Mist blowers or airblast sprays may be used. The maximum application rate is 3.2 pints product per acre; the maximum number of applications per year is 2.
			For air application, mix 0.4 gallons of Malathion 8 in at least 10 gallons of water.	Apply a minimum of 10 gallons of mixture per acre. Make two applications, the first when female flowers are in twig bud stage, the second one week prior to maximum flower receptivity to pollen. The maximum application rate is 3.2 pints product per acre; the maximum number of applications per year is 2.

MOSQUITO CONTROL

AROUND THE OUTSIDE OF BUILDINGS

Around lower outside foundations of homes, yards - spot treatment only. Outdoor garbage cans, and garbage dumps: Apply 0.2439 gallons of Malathion 8 undiluted per 1000 sq. ft. on painted surfaces. Apply 0.2439 gallons of Malathion 8 undiluted per 1000 sq. ft. on unpainted surfaces.

CULL FRUIT AND VEGETABLE DUMP

Around cull fruit and vegetable dumps: Apply 6.857 pounds of Malathion 8 undiluted per 1000 sq. ft. on painted surfaces. Apply 2 gallons of Malathion 8 undiluted per 1000 sq. ft. on unpainted surfaces.

APPLICATION THROUGH IRRIGATION SYSTEMS - CHEMIGATION

Apply this product only through sprinkler, including center pivot, lateral move, end tow side (wheel) roll, traveler, big gun, solid set, or hand move, or drip (including surface and subsurface) irrigation systems. Do not apply this product through any other type of irrigation system.

Crop injury, lack of effectiveness, or illegal pesticide residues in the crop can result from nonuniform distribution of treated water.

If you have questions about calibration, you should contact State Extension Service specialists, equipment manufacturers or other experts.

Do not connect an irrigation system (including greenhouse systems) used for pesticide application to a public water system unless the pesticide label prescribed safety devices for public water systems are in place.

A person knowledgeable of the chemigation system and responsible for its operation or under the supervision of the responsible person, shall shut the system down and make necessary adjustments should the need arise.

Mix in clean supply tank the specified amount of this product for acreage to be covered, and needed quantity of water.

This product should not be tank-mixed with other pesticides, surfactants or fertilizers unless prior use has shown the combination noninjurious under your conditions of use. Follow precautionary statements and directions for all tank-mix products.

On all crops, use sufficient gallonage of water to obtain thorough and uniform coverage, but not cause runoff or excessive leaching. This will vary depending on equipment, pest problem and stage of crop growth. Application of more or less than optimal quantity of water may result in decreased chemical performance, crop injury or illegal pesticide residues.

Meter this product into the irrigation water uniformly during the period of operation. Do not overlap application. Follow specified label rates, application timing, and other directions and precautions for crop being treated. Continuous mild agitation of pesticide mixture may be needed to assure a uniform application, particularly if the supply tank requires a number of hours to empty.

Do not apply when wind speed favors drift beyond the area intended for treatment.

CHEMIGATION SYSTEMS CONNECTED TO PUBLIC WATER SYSTEMS

Note: Gowan Company does not encourage connecting chemigation systems to public water supplies. The following information is provided for users who have diligently considered all other application and water supply options before electing to make such a connection.

Public water systems means a system for the provision to the public of piped water for human consumption if such system has at least 15 service connections or regularly serves an average of at least 25 individuals daily at least 60 days out of the year. Chemigation systems connected to public water systems must contain a functional reduced-pressure zone, backflow preventer (RPZ) or the functional equivalent in the water supply line upstream from the point of pesticide introduction. As an option to the RPZ, the water from the public water system should be discharged into a reservoir tank prior to pesticide introduction. There shall be a complete physical break (air gap) between the flow outlet end of the fill pipe and the top or overflow rim of the reservoir tank of at least twice the inside diameter of the fill pipe.

The pesticide injection pipeline must contain a functional, automatic, quick-closing check valve to prevent the flow of fluid back toward the injection.

13/
13

The pesticide injection pipeline must contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shutdown.

The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops, or in cases where there is no water pump, when the water pressure decreases to the point where pesticide distribution is adversely affected. Systems must use a metering pump, such as a positive displacement injection pump (e.g., diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock. Do not apply when wind speed favors drift beyond the area intended for treatment.

SPRINKLER CHEMIGATION (FOLIAR SPRAY USES)

The system must contain a functional check valve, vacuum relief valve and low pressure drain appropriately located on the irrigation pipeline to prevent water source contamination from backflow. The pesticide injection pipeline must contain a functional, automatic, quick-closing check valve to prevent the flow of fluid back toward the injection pump.

The pesticide injection pipeline must also contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down. The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops. The irrigation line or water pump must include a functional pressure switch which will stop the water pump motor when the water pressure decreases to the point where pesticide distribution is adversely affected.

Systems must use a metering pump, such as a positive displacement injection pump (e.g., diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.

Do not apply when wind speed favors drift beyond the area intended for treatment.

DRIP (INCLUDING SURFACE AND SUBSURFACE) CHEMIGATION

The system must contain a functional check valve, vacuum relief valve and low pressure drain appropriately located on the irrigation pipeline to prevent water source contamination from backflow.

The pesticide injection pipeline must contain a functional, automatic, quick-closing check valve to prevent the flow of fluid back toward the injection pump.

The pesticide injection pipeline must also contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down.

The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops. The irrigation line or water pump must include a functional pressure switch which will stop the water pump motor when the water pressure decreases to the point where pesticide distribution is adversely affected.

Systems must use a metering pump such as a positive displacement injection pump (e.g., diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.

STORAGE AND DISPOSAL

DO NOT CONTAMINATE water, food or feed by storage or disposal.

PESTICIDE STORAGE: Gowan Malathion 8 should be stored in the original unopened container in a secure, dry place. Do not contaminate with other pesticides or fertilizers. The product should never be heated above 55°C (131°F), and should not be stored for long periods of time at a temperature in excess of 25°C (77°F).

PESTICIDE DISPOSAL: To avoid wastes, use all materials in this container by application according to label directions. If wastes cannot be avoided, offer remaining product to a waste facility or pesticide disposal program (often such programs are run by state or local governments or by industry).

CONTAINER DISPOSAL: Nonrefillable container. Do not reuse or refill this container. Offer for recycling, if available. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. After cleaning, if recycling is not available, puncture and dispose of in a sanitary landfill.

FOR 24-HOUR EMERGENCY ASSISTANCE (SPILL, LEAK OR FIRE), CALL CHEMTREC® (800) 424-9300
For other product information, contact Gowan Company or see Material Safety Data Sheet.

NOTICE OF CONDITIONS OF SALE AND WARRANTY AND LIABILITY LIMITATIONS

Important: Read the entire Directions for Use and Notice of Conditions of Sale and Warranty and Liability Limitations before using this product. If terms are not acceptable return the unopened container for a full refund.

Our directions for use of this product are based on tests believed to be reliable. However, it is impossible to eliminate all risk associated with the use of this product. Crop injury, inadequate performance, or other unintended consequences may result due to soil or weather conditions, off target movement, presence of other materials, method of use or application, and other factors, all of which are beyond the control of Gowan Company. All such risks shall be assumed by the Buyer and User.

Gowan Company warrants that this product conforms to the specifications on the label when used in strict conformance with Direction for Use, subject to the above stated risk limitations. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, GOWAN COMPANY MAKES NO OTHER EXPRESS OR IMPLIED WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE NOR ANY OTHER EXPRESS OR IMPLIED WARRANTY.

TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, GOWAN COMPANY'S EXCLUSIVE LIABILITY FOR ANY AND ALL LOSSES, INJURIES OR DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT WHETHER IN CONTRACT, WARRANTY, TORT, NEGLIGENCE, OR ANY OTHER LEGAL THEORY IS STRICTLY LIMITED TO THE PURCHASE PRICE PAID OR REPLACEMENT OF PRODUCT, AT GOWAN COMPANY'S SOLE DISCRETION.

Chemtrek® is a registered trademark of American Chemistry Council, Inc.