



DKIL Cryolite 50 Dust

Granulate*	% by Wt
	50
	50
	100
Expressed as elemental	27.1

CAUTION
 OUT OF REACH OF CHILDREN
 Avoid Inhalation of Dust

DANGEROUS TO BEES exposed to treatment or residues on
 may be obtained from your Cooperative Agricultural

WEIGHT 50 LBS.

USE STRICTLY IN ACCORDANCE WITH CAUTIONS, WARNINGS
 WITH APPLICABLE STATE AND FEDERAL REGULATIONS.
 DOSAGE MAY VARY AS A CONSEQUENCE OF LOCAL
 WEATHER. WE RECOMMEND REFERENCE TO STATE AGRICULTURAL
 CONCERNING SPECIFIC USES.

CONTAINER!

USE ONLY WHEN EMPTY. STAY OUT OF SMOKE.

Apply application by ground or air equipment.

Apply pounds per acre depending on plant size and insect
 as necessary. Remove residues remaining at
 or other effective means. Do not apply under
 drift to food, forage or other planting that might
 be rendered unfit for sale, use or consumption.

U.S.D.A. Reg. No. 10163-22

BEANS: Beetles, Bean Leafcutter -- Repeat at two week intervals..

CABBAGE, CAULIFLOWER, BROCCOLI: Flea Beetles, Cabbage Caterpillars -- DO NOT APPLY to Broccoli after the edible part starts to form. DO NOT APPLY to Cabbage after the heads start to form. DO NOT APPLY to Cauliflower after the seedling stage.

CARROTS: Vegetable Weevil.

CUCUMBERS: Striped Cucumber Beetle, Twelve Spotted Cucumber Beetle, Flea Beetle -- Dust both sides of foliage at first signs of Beetles.

EGGPLANT: Flea Beetle -- Dust thoroughly.

LETTUCE: Climbing Cutworms, Alfalfa Looper, Corn Earworm -- Apply when worms first appear. DO NOT APPLY after the seedling stage.

MELONS: Cucumber Beetle -- Dust both sides of foliage. Repeat if necessary.

PEPPERS: Pepper Weevil -- Dust thoroughly at first set of fruit. Repeat every 5 days until about 6 applications have been made.

STRAWBERRIES: Leaf Beetle -- Dust both sides of leaves.

TOMATOES: Flea Beetle -- Apply when plants are first set out and repeat as necessary. If applied after fruit starts to set, remove residue at harvest by washing bushes or other effective means.

NOTICE ON CONDITIONS OF SALE

This is a general statement of the product and is based upon tests believed to be reliable. The use of this product keeps beyond the control of the manufacturer. No guarantee is made as to the results to be obtained from its use, or as to the results of its use in conjunction with other materials.

ACCEPTED
 July 14, 1971
 U.S. DEPARTMENT OF AGRICULTURE
 PESTICIDE REGISTRATION DIVISION

The Dune Company
 AGRICULTURAL CHEMICALS

Route 1, Box 33
 Somerton, Arizona 85350
 Telephone: 627-2479
 ARIZONA

PROKIL Cryo

Active Ingredients	by Wt
Sodium Fluoroaluminate*	50
Inert Ingredients	50
	100
*Fluorine expressed as elemental	27.1

CAUTION

KEEP OUT OF REACH OF CHILDREN
Avoid Inhalation of Dust

THIS PRODUCT IS HIGHLY TOXIC TO BEES exposed to treatment or residues on crops. Protective information may be obtained from your Cooperative Agricultural Extension Service.

NET WEIGHT 50 LBS.

READ ENTIRE LABEL. USE STRICTLY IN ACCORDANCE WITH CAUTIONS, WARNINGS AND DIRECTIONS, AND WITH APPLICABLE STATE AND FEDERAL REGULATIONS. USES, TIMING AND DOSAGE MAY VARY AS A CONSEQUENCE OF LOCAL WEATHER OR CONDITIONS. WE RECOMMEND REFERENCE TO STATE AGRICULTURAL EXTENSION SERVICE CONCERNING SPECIFIC USES.

KEEP PESTICIDE IN ORIGINAL CONTAINER!

BURN BAG IMMEDIATELY WHEN EMPTY. STAY OUT OF SMOKE.

This product is formulated for application by ground or air equipment.

Use at rate of 20 to 40 pounds per acre depending on plant size and insect population. Repeat application as necessary. Remove residues remaining at harvest by washing, brushing or other effective means. Do not apply under conditions involving possible drift to food crops or other planting that might be damaged by the crop, the root, or the fruit for sale or consumption.

U.S. Pat. 2,113,000

U.S.D.A. Reg. No. 10163-22

CALIFORNIA (Plant)

P. O. Box 406

310 East Main Street

Calipatria, Calif. 92232

Telephone 348-5150

ACCEPTED

12/19/71

The Dune

AGRICULTURAL

