

# P R O K I L



## SUGGESTED USE

READ ENTIRE LABEL. USE STRICTLY IN ACCORDANCE WITH WARNINGS AND DIRECTIONS, AND WITH APPLICABLE STATE AND FEDERAL REGULATIONS. TIMING AND DOSAGE VARY GREATLY BETWEEN DISTRICTS, AS A CONSEQUENCE OF LOCAL WEATHER AND OTHER CONDITIONS. ALWAYS FOLLOW THE RECOMMENDATIONS OF STATE AGRICULTURAL EXPERIMENT STATION CONCERNING SPECIFIC USAGE.

BURN BAG IMMEDIATELY WHEN EMPTY. STAY OUT OF SMOKE

Apply thoroughly. Do not apply on food crops within 3 days of harvest unless otherwise specified. Remove residues remaining at harvest by washing, brushing or other effective means.

**CUCUMBERS, SQUASH, MELONS:** Aphids, Spider Mites, Pickleworm, Cucumber Beetles, Flea Beetles; **CANTALOUPE** — Leafhoppers — 25 to 40 lbs. per acre. Repeat as necessary. **SQUASH:** Squash Vine Borers — 40 lbs. per acre. **PUMPKINS:** Squash Vine Borers — 45 lbs. per acre. Apply twice a week to stems and vines at base of plants. Do not apply to Pumpkins within 3 days of harvest, and to Cucumbers, Cantaloupes, Squash and Melons within 1 day of harvest. Do not apply unless plants are dry.

**LETTUCE:** Aphids, Mites, Cabbage Looper, Leafhoppers, Climbing Cutworms, Alfalfa Looper, Corn Earworms — 30 to 40 lbs. per acre. Do not apply after the seedling stage.

## Malathion Cryolite 4.50 Dust

Active Ingredients	By Wt.
Malathion*	4%
Sodium Fluoaluminate	46%
Inert Ingredients	50%
*O,O-dimethyl dithiophosphate of diethyl mercaptosuccinate.	
Fluorine (expressed as elemental)	25%

**READ THE LABEL  
KEEP OUT OF REACH OF CHILDREN  
CAUTION**

Avoid prolonged breathing of dust.

Avoid repeated or prolonged contact with skin; wash thoroughly after using. Avoid contamination of feed and foodstuffs. Do not apply under conditions involving possible drift to food, forage or other plantings that might be damaged or the crop thereof rendered unfit for sale, use or consumption.

SEE RIGHT PANEL FOR ADDITIONAL CAUTIONS!

NET WEIGHT 50 LBS.

**CABBAGE:** Aphids, Cabbage Looper, Climbing Cutworms and Alfalfa Looper — 30 per acre. Do not apply after head starts to

**TOMATOES:** Aphids, Spider Mites, Tomato Russetworm, Armyworms, Flea Beetles — 30 per acre. Do not apply within 1 day of har

## CAUTION

This product will kill fish. Keep out of an water. Do not apply where runoff is likely to not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as directed on this label.


This product is highly toxic to bees exposed to it on flowers or residues on crops. Protective information should be obtained from your Cooperative Agricultural Extension Service.

## NOTICE ON CONDITIONS OF SALE

The user assumes all liability for any loss of crop or other damage resulting from the use of this product. The use of this product is beyond the control of the manufacturer and no warranty is made as to the effects of the product when used in accordance with the established practice. The manufacturer assumes no liability for injury or damage resulting from its misuse as such or in combination with other materials.

U.S.D.A. Reg. No.

CALIFORNIA (Plant)  
P. O. Box 406  
340 East Main Street  
Calipatria, Calif. 92233  
Telephone: 348-5150



## The Dune Company

AGRICULTURAL CHEMICALS

Route 1,  
Somerton, Arizona  
Telephone: 62

# P R O

## SUGGESTED USE

READ ENTIRE LABEL. USE STRICTLY IN ACCORDANCE WITH WARNINGS AND DIRECTIONS, AND WITH APPLICABLE STATE AND FEDERAL REGULATIONS. TIMING AND DOSAGE VARY GREATLY BETWEEN DISTRICTS, AS A CONSEQUENCE OF LOCAL WEATHER AND OTHER CONDITIONS. ALWAYS FOLLOW THE RECOMMENDATIONS OF STATE AGRICULTURAL EXPERIMENT STATION CONCERNING SPECIFIC USAGE.

DO NOT APPLY TO FOOD CROPS WITHIN 3 DAYS OF HARVEST UNLESS OTHERWISE SPECIFIED.

Apply thoroughly. Do not apply on food crops within 3 days of harvest unless otherwise specified. Remove residues remaining at harvest by washing, brushing or other effective means.

**CUCUMBERS, SQUASH, MELONS:** Aphids, Spider Mites, Pickleworm, Cucumber Beetles, Flea Beetles; **CANTALOUPE** — Leafhoppers — 25 to 40 lbs. per acre. Repeat as necessary. **SQUASH:** Squash Vine Borers— 40 lbs. per acre. **PUMPKINS:** Squash Vine Borers — 45 lbs. per acre. Apply twice a week to stems and vines at base of plants. Do not apply to Pumpkins within 3 days of harvest and to Cucumbers, Cantaloupes, Squash and Melons within 1 day of harvest. Do not apply unless plants are dry.

**LETTUCE:** Aphids, Mites, Cabbage Looper, Leafhoppers, Climbing Cutworms, Alfalfa Looper, Corn Earworms— 30 to 40 lbs. per acre. Do not apply after the seedling stage.

# Malathion Cryolite D

Active Ingredients  
Malathion\*  
Sodium Fluoroaluminate  
Inert Ingredients  
\*O,O-dimethyl dithiophosphoric  
succinate.  
Fluorine expressed as

READ THE LABEL  
KEEP OUT OF REACH OF CHILDREN

## CAUTION

Avoid

Avoid repeated or prolonged contact with skin. Wash thoroughly after using. Do not use on or near food and foodstuffs. Do not allow drift to food crops. Avoid possible drift to food crops that might be damaged or rendered unfit for sale, use or consumption.

SEE RIGHT PANEL FOR DIRECTIONS

NET WEIGHT

CALIFORNIA (Plant)  
P. O. Box 406  
310 East Main Street  
Calipatria, Calif. 92233  
Telephone 348-5150

1/23/70

# The Dunes

AGRICULTURAL CHEMICALS