

Agricultural Poison Bait

ACTIVE INGREDIENT:	% by Wt.
Carbaryl (1-naphthyl N-methylcarbamate)	5
INERT INGREDIENTS	95
	100

*Registered Trade-mark of Union Carbide Corp. for the active ingredient, 1-naphthyl N-methylcarbamate.

Do not reuse container. DESTROY WHEN EMPTY!

CAUTION

KEEP OUT OF REACH OF CHILDREN

HARMFUL IF SWALLOWED OR INHALED

Do Not Breathe Dust. Do Not Take Internally.
Skin Contact May Be Harmful. Avoid Contact.
Wash Hands Before Eating. Take Shower or Bath After Work.
Wear Regular Long Sleeved Work Clothing. Change to Clean Clothing Daily.
NOTE TO PHYSICIAN: Carbaryl is a moderate, reversible, Cholinesterase inhibitor.
Atropine is antidotal.
AVOID CONTAMINATION OF FOOD feed, feeding troughs and watering receptacles

USE PRECAUTIONS

Toxic to honey bees. For protection to honey bees avoid use, if possible, during periods when honey bees are visiting the crop. When necessary to use during such periods, warn beekeepers well in advance to locate hives at a safe distance until 1 week after application.

Avoid drift to adjoining food and forage crops.

In order that pesticidal residues on food and forage crops will not exceed tolerances established by the Federal Food and Drug Administration use only at recommended rate, and intervals, and do not apply closer to harvest than specified. Do not apply or allow to drift to areas occupied by unprotected humans or beneficial animals or onto adjoining food, fiber or pasture crops. The grower is responsible for residues on his crops, as well as for damages caused by drift from his property to that of others.

Consult State Agricultural Experiment Stations or State Agricultural Extension Service for additional information, as the timing, number and rate of applications needed will vary with local conditions.

NOTICE ON CONDITIONS OF SALE:

Our recommendations for use of this product are based upon tests believed to be reliable. The use of this product being beyond the control of the manufacturer, no guarantee, expressed or implied, is made as to the effects of such or the results to be obtained if not used in accordance with directions or established safe practice. The buyer must assume all responsibility, including injury or damage, resulting from its misuse as such, or in combination with other materials.

RECOMMENDATIONS

Prokil Sevin Bait is recommended for control of armyworms, cutworms, crickets, darkling ground beetle, grasshoppers and sowbugs on crops listed.

DIRECTIONS FOR USE (May be used only in California and Arizona)

Recommended dosage rates refer to pounds of Prokil Sevin Bait per acre unless otherwise specified. Use suitable ground or aircraft equipment which provides good distribution. Apply when pests appear in damaging numbers and repeat 7 to 14 days later if necessary, but observe use limitations given for specific crops. Evening applications are preferred.
DO NOT USE WHEN BEES ARE PRESENT.

ALFALFA, PEAS, COTTON—Cutworms, Armyworms, Sowbugs, Grasshoppers, Darkling Ground Beetles, Crickets. Apply 30 lbs per acre. May be used on alfalfa and peas up to 1 day of harvest. Do not apply within 7 days of harvest or grazing of cotton.

CUCUMBERS, SQUASH, MELONS—Cutworms, Armyworms, Sowbugs, Grasshoppers, Darkling Ground Beetles, Crickets. Apply 20 lbs. per acre. May be used up to 1 day of harvest.

BEANS, CARROTS, CORN FORAGE AND SWEET CORN, EGG PLANT, OKRA, PEPPER, POTATO, TOMATO—Cutworms, Armyworms, Sowbugs, Grasshoppers, Darkling Ground Beetles, Crickets. Apply 40 lbs. per acre. Do not apply within 1 day of harvest.

ASPARAGUS, STRAWBERRIES—Cutworms, Armyworms, Sowbugs, Grasshoppers, Darkling Ground Beetles Crickets. Apply 40 lbs. per acre. Do not apply within 1 day of harvest.

BROCCOLI, BRUSSELS SPROUTS, CABBAGE, CAULIFLOWER, HEAD LETTUCE, GARDEN BEET ROOTS, HORSERADISH, PARSNIP, RADISH, RUTABAGA, TURNIP—Cutworms, Armyworms, Sowbugs, Grasshoppers, Darkling Ground Beetles, Crickets. Apply 40 lbs. per acre. Do not apply within 5 days of harvest.

SUGAR BEET, COLLARDS, ENDIVE, GARDEN BEET TOPS, KALE, LEAF LETTUCE, PARSLEY, SPINACH, SWISS CHARD, TURNIP TOPS—Cutworms, Armyworms, Sowbugs, Grasshoppers, Darkling Ground Beetles, Crickets. Apply 40 lbs. per acre. Do not apply within 14 days of harvest.

NET WEIGHT 50 LBS.

EPA Reg. No. 10163-32 AA

8-71 139

CALIFORNIA (Plant)

P. O. Box 406
340 East Main Street
Calipatria, Calif. 92233
Telephone: 348-5150

The Dune Company

AGRICULTURAL CHEMICALS

ARIZONA

Route 1, Box 33
Somerton, Arizona 85350