

# P R O K I L



## RECOMMENDED USE

USE STRICTLY IN ACCORDANCE WITH DIRECTIONS, AND WITH APPLICABLE FEDERAL REGULATIONS. TIMING Varies GREATLY BETWEEN DISTRICTS OF LOCAL WEATHER AND CLIMATIC CONDITIONS. ALWAYS FOLLOW THE RECOMMENDATIONS OF AGRICULTURAL EXPERIMENT STATION SPECIFIC USAGE.

KEEP EMPTY CONTAINER OUT OF REACH OF CHILDREN. Do not apply on food crops within 1 day of harvest unless otherwise specified. Remove residue from plants by washing, brushing or spraying.

**CANTON:** Aphids, Spider Mites, Flea Beetles, Flea Beetles; **CANTON:** 25 to 40 lbs. per acre. **FLORIDA:** Squash Vine Borers—25 to 40 lbs. per acre. **FLORIDA:** Squash Vine Borers—25 to 40 lbs. per acre. Do not apply to Pumpkins and to Cucumbers, Cantons within 1 day of harvest. Do not apply to Pumpkins and to Cucumbers, Cantons within 1 day of harvest. Do not apply to Pumpkins and to Cucumbers, Cantons within 1 day of harvest.

**CABBAGE:** Cabbage Looper, Leafhoppers, Cabbage Looper, Corn Earworms—Do not apply after the seed-

## Malathion Cryolite 4.50 Dust

Active Ingredients:	By Weight
Malathion*	45%
Sodium Fluoaluminate	46%
Inert Ingredients	50%
*O,O-dimethyl dithiophosphate of diethyl mercaptosuccinate.	
Fluorine expressed as elemental	25%

**READ THE LABEL  
KEEP OUT OF REACH OF CHILDREN**

### CAUTION

Avoid prolonged breathing of dust.

Avoid repeated or prolonged contact with skin; wash thoroughly after using. Avoid contamination of feed and foodstuffs. Do not apply under conditions involving possible drift to food, forage or other plantings that might be damaged or the crop thereof rendered unfit for sale, use or consumption.

SEE RIGHT PANEL FOR ADDITIONAL CAUTIONS!

NET WEIGHT 50 LBS.

**CABBAGE:** Aphids, Cabbage Looper, Leafhopper, Climbing Cutworms and Alfalfa Looper — 30 to 40 lbs. per acre. Do not apply after head starts to form.

**TOMATOES:** Aphids, Spider Mites, Tomato Russet Mites, Drosophila, Armyworms, Flea Beetles — 30 to 45 lbs. per acre. Do not apply within 1 day of harvest.

### CAUTION

This product will kill fish. Keep out of any body of water. Do not apply where runoff is likely to occur. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on this label.

This product is highly toxic to bees exposed to treatment or residues on crops. Protective information may be obtained from your Cooperative Agricultural Extension Service.

#### NOTICE ON CONDITIONS OF SALE

Our recommendations for use of this product are based on tests believed to be reliable. The use of this product beyond the control of the manufacturer no guarantee, expressed or implied, is made as to the effects of such or the results to be obtained if not used in accordance with the label or established safe practices. The buyer must assume all responsibility including injury or damage, resulting from its misuse, as such or in combination with other materials.

U.S.D.A. Reg. No. 10163-25

ACCEPTED  
10/23/70  
FOR REGISTRATION UNDER FIFTEEN SUBJECT TO ATTACHED COMMENTS.

## The Dune Company

AGRICULTURAL CHEMICALS

ARIZONA

Route 1, Box 33

Somerton, Arizona 85350

Telephone: 627-2479